

2024-07-31  
2023-3760

(Container Label)



GROUP	2	HERBICIDE
-------	---	-----------

## PRISM™ HERBICIDE

25% DRY FLOWABLE

FOR SALE FOR USE IN WEED CONTROL FOR POTATOES IN EASTERN CANADA, IRRIGATED POTATOES IN WESTERN CANADA, SEED POTATOES AND TRANSPLANTED PROCESSING TOMATOES AND FRESH MARKET FIELD TOMATOES ONLY AND WEED CONTROL/SUPPRESSION IN HIGHBUSH BLUEBERRY

COMMERCIAL (AGRICULTURAL)

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

**ACTIVE INGREDIENT:** Rimsulfuron 25.0%  
Wettable granules

WARNING - EYE IRRITANT

REGISTRATION NO. 23983 PEST CONTROL PRODUCTS ACT

**NET CONTENTS:** 480 GRAMS

### **Corteva Agriscience Canada Company**

Suite 240, 115 Quarry Park Rd. SE  
Calgary, AB  
T2C 5G9  
1-800-667-3852

™Trademark of Corteva Agriscience and its affiliated companies.

## PRECAUTIONS

- KEEP OUT OF REACH OF CHILDREN.
- AVOID BREATHING SPRAY MIST.
- AVOID CONTACT WITH SKIN, EYES AND CLOTHING.
- DO NOT CONTAMINATE ANY BODY OF WATER.
- DO NOT APPLY BY AIR.
- Mixers, loaders and applicators must wear a long-sleeved shirt, long pants, and chemical resistant gloves (gloves not required for groundboom sprayers). Wear gloves and protective goggles for mixing/loading /clean -up operations and when making sprayer and nozzle repairs and adjustments. Do not handle pesticides with bare hands. Chemical-resistant gloves significantly reduce hand exposure. Do not use leather or cloth gloves.

## ENVIRONMENTAL PRECAUTIONS

- TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.
- This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.
- Avoid application when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

## FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**IF SWALLOWED:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**TOXICOLOGICAL INFORMATION:** Treat symptomatically.

## STORAGE

Store this product away from food or feed. To prevent contamination, store product in original container in a secure, dry area, away from other pesticides, fertilizer, seed, food or feed. Not for use or storage in or around the home. Keep container tightly closed.

## DISPOSAL

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/ territorial regulatory agency. Contact the manufacturer and the provincial/ territorial regulatory agency in case of a spill, and for clean-up of spills.

**NOTICE TO USER**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use a control product in a way that is inconsistent with the directions on the label.

(Booklet)



GROUP	2	HERBICIDE
-------	---	-----------

## **PRISM™ HERBICIDE**

25% DRY FLOWABLE

FOR SALE FOR USE IN WEED CONTROL FOR POTATOES IN EASTERN CANADA, IRRIGATED POTATOES IN WESTERN CANADA, SEED POTATOES AND TRANSPLANTED PROCESSING TOMATOES AND FRESH MARKET FIELD TOMATOES ONLY AND FOR WEED CONTROL/SUPPRESSION IN HIGHBUSH BLUEBERRY

COMMERCIAL (AGRICULTURAL)

READ THE LABEL AND BOOKLET BEFORE USING

**ACTIVE INGREDIENT:** Rimsulfuron                      25.0%  
Wettable granules

WARNING - EYE IRRITANT

REGISTRATION NO. 23983 PEST CONTROL PRODUCTS ACT

**NET CONTENTS:** 0.25 – 5 KILOGRAMS

### **Corteva Agriscience Canada Company**

Suite 240, 115 Quarry Park Rd. SE

Calgary, AB

T2C 5G9

1-800-667-3852

™Trademark of Corteva Agriscience and its affiliated companies.

## PRECAUTIONS

- KEEP OUT OF REACH OF CHILDREN.
- AVOID BREATHING SPRAY MIST.
- AVOID CONTACT WITH SKIN, EYES AND CLOTHING.
- DO NOT CONTAMINATE ANY BODY OF WATER.
- DO NOT APPLY BY AIR.
- Mixers, loaders and applicators must wear a long-sleeved shirt, long pants, and chemical resistant gloves (gloves not required for groundboom sprayers). Wear gloves and protective goggles for mixing/loading /clean -up operations and when making sprayer and nozzle repairs and adjustments. Do not handle pesticides with bare hands. Chemical-resistant gloves significantly reduce hand exposure. Do not use leather or cloth gloves.

## FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**IF SWALLOWED:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**TOXICOLOGICAL INFORMATION:** Treat symptomatically.

## STATUTORY CONDITIONS RELATING TO USE:

Mode of Action:	ALS inhibitor
Crops:	potatoes, transplanted processing tomatoes, fresh market field tomatoes
Maximum Rate:	60 g/ha for potatoes and 140 g/ha for transplanted processing tomatoes and fresh market field tomatoes
Maximum No. of Treatments:	One per year
PHI:	30 days before harvest
Application Timing:	post emergent

Because potato varieties differ in their tolerance to herbicides, limit first use of Prism Herbicide 25% Dry Flowable to a small area of each variety prior to adoption as a field practice.

DO NOT APPLY BY AIR.

Carefully observe sprayer cleanup instructions, as spray tank residue may damage other crops.

Do not apply Prism Herbicide 25% Dry Flowable during periods of intense rainfall or to soils saturated with water. Do not apply directly to standing or running water. Do not apply in areas where surface water from the treatment site can run off to adjacent cropland, either planted or to be planted, or into streams, irrigation water or wells. Applications should only be made when there is no hazard of spray drift contaminating non-target land areas since very small quantities of the Prism Herbicide 25% Dry Flowable spray solution may severely injure susceptible crops during both growing and dormant periods.

Application of Prism Herbicide 25% Dry Flowable to potatoes may result in TEMPORARY foliar symptoms (discolouration of younger leaves and pinching of the terminal leaf) that may be confused with symptoms of a viral disease. Early application timing may reduce the likelihood of foliar symptoms.

As with any herbicide, overlaps or starting, stopping, slowing and turning while spraying may result in crop injury.

Under extreme weather conditions, such as hot, dry weather, excessive moisture, or frost, weed control may be reduced.

Rainfall within 2-4 hours after application of Prism Herbicide 25% Dry Flowable may reduce weed control.

CAUTION: Do not graze the treated crop or cut for hay; sufficient data are not available to support such use.

#### **ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY**

- A rapid fluctuation in temperature (greater than 20 deg C difference within 24-36 hours) will stress the crop. For maximum crop safety, allow 48-72 hours for the crop to acclimatize before spraying Prism Herbicide 25% Dry Flowable.
- Only apply Prism Herbicide 25% Dry Flowable when the temperature in the 24 hours before AND after the application ranges between 5 deg C and 28 deg C. Temperatures beyond this range increase the potential for crop injury.
- **WARNING:** Crop injury may result if application is made to potatoes that have been stressed by abnormally hot, humid or cold weather conditions, frost, low fertility, drought, water saturated soil, compacted soil, previous pesticide applications, disease or insect damage. If potatoes have been injured by frost, wait 48-72 hours before applying Prism Herbicide 25% Dry Flowable.
- Prism Herbicide 25% Dry Flowable rapidly stops growth of susceptible species; typical symptoms usually appear within 5-7 days, but may not be noticeable for 2-3 weeks after application, depending on growing conditions and weed susceptibility.
- Warm, moist conditions following application promote the activity of Prism Herbicide 25% Dry Flowable, while cool and/or dry conditions may reduce or delay activity. Weeds hardened off by cold weather or drought stress may not be controlled.
- Prism Herbicide 25% Dry Flowable is a contact herbicide, and will not give residual control of grass or broadleaf weed seedlings that may germinate after application. Weed control may be reduced if the crop canopy has closed in over the weeds, intercepting the spray.

#### **ENVIRONMENTAL PRECAUTIONS**

- TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.
- This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.
- Avoid application when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

#### **DIRECTIONS FOR USE**

Prism Herbicide 25% Dry Flowable is a dry flowable granular formulation to be mixed in water and applied post-emergence to potatoes for control of quackgrass, annual grasses and redroot pigweed. Prism Herbicide 25% Dry Flowable is non-corrosive, non-flammable, non-volatile, and does not freeze.

This jug will treat 8 ha.

Control will not be achieved if surfactant is not applied with Prism Herbicide 25% Dry Flowable.

Make only one application per year.

- Do not re-enter treated fields until 12 hours after application.

- Apply only to agricultural crops when the potential for drift to areas of human habitation and areas of human activity (houses, cottages, schools and recreational areas) is minimal. Take into consideration wind speed, wind direction, temperature, application equipment and sprayer settings.
- As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.
- DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

### Field sprayer application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

DO NOT apply by air.

### SPRAY BUFFER ZONES

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, rangelands, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop	Spray Buffer Zones (metres) Required for the Protection of:		
		Terrestrial habitat	Aquatic habitat of water depths:	
			Less than 1 m	Greater than 1 m
Field sprayer	Transplanted processing tomatoes Potatoes Highbush blueberries	5	1	1
	Field tomatoes	15	1	1

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

### SPRAY ADJUVANTS

Prism Herbicide 25% Dry Flowable must be applied with a non-ionic surfactant, either Citowett Plus, Agral 90 or Ag-Surf at 2 L per 1000 L spray solution (0.2% v/v).

### RATES AND TIMING

Apply 60 grams/ha of Prism Herbicide 25% Dry Flowable as a broadcast spray, with a surfactant, to potatoes prior to initiation of flowering.

Apply to weeds according to the leaf staging in the chart below:

<b>WEED</b>	<b>LEAF STAGING AT APPLICATION</b>
Annual grasses	1 to 6 leaf stage (up to early tillering, –two 2-leaf tillers)
Quackgrass	Apply to actively growing shoots in the 3 to 6 leaf (< 25 cm leaf extended) stage
Lamb's-quarters, Redroot Pigweed	4 to 6 leaf stage (< 10 cm tall or =across)

Application to control ANNUAL GRASSES and QUACKGRASS must be made before the crop canopy can interfere with spray coverage of the target weeds.

Cultivation is NOT recommended within 7-10 days prior to or after application of Prism.

#### **WEEDS CONTROLLED AT 60 GRAMS PER HECTARE**

Barnyard Grass	Fall Panicum	Green Foxtail
Lamb's-Quarters <sup>†</sup>	Redroot Pigweed	Quackgrass <sup>1</sup>
Yellow Foxtail <sup>2</sup>	Witchgrass	

<sup>†</sup> Suppression only. Weed suppression is a visual reduction in weed competition (reduced population and/or vigour) as compared to an untreated check. Degree of suppression will vary with size of weed and environmental conditions prior to and following treatment.

NOTE: Weeds which emerge after application of Prism Herbicide 25% Dry Flowable will not be controlled. Application should be made when the majority of weeds have emerged.

Annual grass and broadleaf weeds are most sensitive when small and actively growing.

Early crop establishment and a good crop stand are important in providing competition for weeds and effective postemergence control of quackgrass and annual weeds when using Prism Herbicide 25% Dry Flowable.

#### **Specific Weed Control Recommendations:**

<sup>1</sup> Quackgrass - Apply Prism Herbicide 25% Dry Flowable with a non-ionic surfactant postemergence when the majority of the quackgrass shoots are actively growing and in the 3 to 6 leaf stage (<25cm - leaf extended). Annual grasses not emerged at the time of Prism Herbicide 25% Dry Flowable application to quackgrass will not be controlled. Application to quackgrass prior to the 3 to 6 leaf stage may result in reduced weed control since the majority of shoots may not have emerged.

<sup>2</sup> Yellow foxtail present at the time of Prism Herbicide 25% Dry Flowable application will be controlled. Any yellow foxtail emerging after application will not be controlled.

#### **Tank Mixtures**

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Corteva Agriscience Canada Company at 1-800-667-3852 or [www.corteva.ca](http://www.corteva.ca) for information before applying any tank mix that is not specifically recommended on this label.

**NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR SPECIAL USE APPLICATIONS**

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Corteva Agriscience Canada Company under the User Requested Minor Use Label Expansion program. For these uses, Corteva Agriscience Canada Company has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

**FOR USE ON IRRIGATED POTATOES IN WESTERN CANADA (MB, SK, AB, BC)**

Apply 60 grams/ha of Prism Herbicide 25% Dry Flowable as a postemergent broadcast spray, with a non-ionic surfactant, to potatoes prior to initiation of flowering. Apply after crop emergence, to young actively growing annual grasses and broadleaf weeds listed in this booklet.

Make only one application per growing season.

For additional application and recropping instructions and/or use precautions, see appropriate headings within this booklet.

NOTE: Annual grass and broadleaf weeds are most sensitive when small and actively growing (2.5 cm in height/diameter).

**TRANSPLANTED PROCESSING TOMATOES**

Apply Prism Herbicide 25% Dry Flowable at a rate of 60 grams/hectare + 0.2% v/v non-ionic surfactant (Citowett Plus, Agral 90 or Ag-Surf) to transplanted processing tomatoes.

Weeds controlled or suppressed include all weeds listed on the Prism Herbicide 25% Dry Flowable label plus hairy nightshade.

NOTE: Prism Herbicide 25% Dry Flowable may be applied up to the 4 leaf stage of hairy nightshade.

Weeds which emerge after application of Prism Herbicide 25% Dry Flowable will not be controlled. Application should be made when the majority of weeds have emerged.

Early crop establishment and a good crop stand are important in providing competition for weeds and effective postemergence control of quackgrass and annual weeds when using Prism Herbicide 25% Dry Flowable.

Apply only one application of Prism Herbicide 25% Dry Flowable per year with ground equipment. Do not harvest crop within 30 days of treatment. Refer to the appropriate sections of this label for recropping information, and for additional application instructions, specific weed recommendations, mixing instructions and/or use precautions.

**FOR USE ON FRESH MARKET FIELD TOMATOES**

For control of hairy nightshade in fresh market field tomatoes, apply Prism Herbicide 25% Dry Flowable at a rate of 60 grams/hectare + 0.2% v/v non-ionic surfactant (Citowett Plus, Agral 90 or Ag-Surf).

Weeds which emerge after application of Prism Herbicide 25% Dry Flowable will not be controlled. Application should be made when the majority of weeds have emerged.

Apply only one application of Prism Herbicide 25% Dry Flowable per year with ground equipment. Do not harvest crop within 30 days of treatment. Refer to the appropriate sections of the label for recropping information, and for additional application instructions, specific weed recommendations and/or use precautions.

### **FIELD TOMATOES**

For control of triazine resistant lambs quarters, hairy nightshade plus other broadleaf weeds listed on the Prism Herbicide label, apply Prism Herbicide as a single post-emergent spray at a rate of 140 grams/hectare + 0.2% v/v non-ionic surfactant (Agral 90) to field tomatoes, at least 21 days after transplanting the crop.

Weeds which emerge after application of Prism Herbicide will not be controlled. Application should be made when the majority of weeds have emerged.

Use ground equipment only. Use in a minimum of 200 litres of water/ha.

Do not harvest crop within 30 days of treatment

### **FOR USE ON Highbush BLUEBERRIES**

For control/suppression of weeds listed on the Prism Herbicide 25% Dry Flowable, make a single application early postemergence to actively growing weeds at a rate of 60 g/ha plus a recommended non-ionic surfactant such as Citowett Plus, Agral 90, or Ag-Surf at 2 L per 1000 L spray solution (0.2 % v/v). Use a directed spray application adjusted to provide complete coverage of the weeds while minimizing the amount of spray coming into contact with the blueberry plants.

When applied as a banded treatment (50% band or less), Prism Herbicide 25% Dry Flowable may be applied twice per year. However, do not apply more than 60 grams per hectare on a broadcast application basis per year.

Use on highbush blueberries that have gone through at least one growing season and are in good health and vigor.

Use ground equipment only. Use a minimum of 100 litres of water/ha (recommended spray volume of 140-190 L/ha).

Do not apply within 21 days of harvest.

Refer to the appropriate sections of the Prism Herbicide 25% Dry Flowable booklet for a complete list of weeds, their stages of control, as well as additional application and/or use precautions instructions, and/or mixing instructions.

### **MIXING INSTRUCTIONS:**

1. Fill clean tank about 1/4 full with fresh water.
2. Turn on full agitation.
3. Add the proper amount of Prism Herbicide 25% Dry Flowable to the spray tank with the agitator running. Continuous agitation is required to keep the product in suspension.
4. After Prism Herbicide 25% Dry Flowable has been well mixed and is in suspension, add a recommended non-ionic surfactant such as Citowett Plus, Agral 90, or Ag-Surf at 2 L per 1000 L spray solution (0.2% v/v).
5. Fill the remainder of the spray tank with water.

On repeat tank loads, ensure that the amount of spray solution left in the tank from the previous load is less than 10% of the volume about to be mixed. Then follow mixing instructions above.

Agitation is required for uniform mixing and application. Apply Prism Herbicide 25% Dry Flowable in a minimum of 100 litres of water per hectare. Flat fan nozzles are recommended. Use a spray pressure of 175 - 275 kPa. Use 50 mesh screens or larger. Use spray preparation of Prism Herbicide 25% Dry Flowable within 24 hours or product degradation may occur resulting in a loss of weed control. Use vigorous agitation to thoroughly disperse spray mixtures that have been allowed to stand in the tank.

NOTE: Prism Herbicide 25% Dry Flowable will degrade in acidic or highly alkaline water. Mix no more than can be used in one day. If spraying is interrupted, thoroughly re-agitate the spray mixture before resuming spraying.

**DO NOT MIX PRISM HERBICIDE 25% DRY FLOWABLE WITH SUBSTANCES THAT CONTAIN BORON OR SUBSTANCES THAT RELEASE FREE CHLORINE.**

Calibrate sprayers only with clean water away from well sites. Make scheduled checks of spray equipment. Assure accurate measurement of products. Mix only enough product for the job at hand. Avoid overfilling spray tank.

### **MANAGEMENT OF SPRAY DRIFT**

When applying Prism Herbicide 25% Dry Flowable near adjacent and/or sensitive plants, do everything possible to reduce the potential for spray drift.

High temperatures, drought and low relative humidity increase the potential for spray drift. Extreme caution must be used when these conditions exist.

Do not apply when a temperature inversion exists characterized by low air movement and an increase in temperature with an inversion in altitude.

### **GROUND AND SURFACE WATER PROTECTION**

Do not mix or load within 10 m of a body of water or wetland.

Do not apply Prism Herbicide 25% Dry Flowable during periods of intense rainfall or to areas where surface water could run off to adjacent land or water bodies, including ditches and streams.

Observe provincial/ territorial regulations and guidelines that regulate treatments around water bodies and well sites.

### **SPRAYER CLEANUP**

To avoid subsequent injury to desirable crops, thoroughly clean all mixing and spray equipment immediately following applications of Prism Herbicide 25% Dry Flowable as follows:

1. Drain tank; thoroughly hose down the interior surfaces of the tank; then flush tank, boom and hoses with clean water for a minimum of 5 minutes.
2. Fill the tank with clean water while adding one litre of household ammonia (containing a minimum of 3% ammonia) for every 100 litres of water. Flush the hoses, boom and nozzles with the cleaning solution and then add more water to completely fill the tank. Circulate the cleaning solution through the tank and hoses for at least 15 minutes. Again flush the hoses, boom and nozzles with the cleaning solution and then drain the tank.
3. Remove the nozzles and screens and clean separately in a bucket containing cleaning agent and water.
4. Repeat step 2.
5. Thoroughly rinse the tank with clean water for a minimum of 5 minutes, flushing the water through the hoses and boom.
6. PRIOR to using the sprayer for the next application, flush the tank, boom and hoses for 5 minutes with fresh water.

## **RESISTANCE-MANAGEMENT RECOMMENDATIONS**

For resistance management, Prism Herbicide 25% Dry Flowable is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to Prism Herbicide 25% Dry Flowable and other Group 2 herbicides. The resistance biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of Prism Herbicide 25% Dry Flowable or other Group 2 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact your local Corteva Agriscience Canada Company representative or the Corteva Agriscience Canada Company hotline at 1-800-667-3852 for further information.

## **STORAGE**

Store this product away from food or feed. To prevent contamination, store product in original container in a secure, dry area, away from other pesticides, fertilizer, seed, food or feed. Not for use or storage in or around the home. Keep container tightly closed.

## **DISPOSAL**

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
  2. Make the empty, rinsed container unsuitable for further use.
- If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

## WILDLIFE AND HABITAT PROTECTION

Injury to or loss of desirable trees or vegetation may result from failure to observe the following: Do not apply or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts, or similar areas. Do not contaminate any body of water, including irrigation water.

Overspray or drift to important wildlife habitats such as ponds, wetlands, streams, woodlots and shelterbelts should be avoided.

## REPLANTING TO OTHER CROPS

Prism Herbicide 25% Dry Flowable is degraded by natural soil processes, and field tests have shown that the following crops may safely be planted at the prescribed interval following potatoes that have been treated with Prism Herbicide 25% Dry Flowable.

### ROTATIONAL CROPS REPLANT INTERVAL:

Winter wheat	4 months
Spring barley, Soybeans, White beans, Red clover, Sorghum, Field corn, Potatoes	10 months

For other crops, a field bioassay is recommended before planting. A successful field bioassay means growing to maturity a test strip of the crop(s) intended for production the following year.

## FIELD BIOASSAY

Select a representative area or areas of the field previously treated with Prism Herbicide 25% Dry Flowable to plant your bioassay crop(s). Be sure to consider factors such as size of field, soil texture, drainage and turn-around areas when selecting the site(s) that are most representative of the soil conditions in the field. On large fields, more than one site may be needed in order to obtain reliable results.

Plant the test strips perpendicular to the direction in which the field was sprayed. The strips should be long enough to cross the width of several spray swaths. Large test strip areas are more reliable than small ones.

Use standard tillage and seeding equipment to plant the bioassay. Prepare a seed bed and plant the crops and varieties you want the option of growing the following year. It is important to use the same planting time, conditions, techniques and cultural practices you normally use to plant and grow the bioassay crop(s). Also plant into an adjacent area not treated with Prism Herbicide 25% Dry Flowable to use as a comparison.

As the crop(s) emerges and grows, examine these key points in Prism Herbicide 25% Dry Flowable treated and non-treated areas:

crop stand	root development	rate of growth
plant colour	vigour -yield	

Allow the bioassay crop(s) to grow to maturity while making your observations. Do not overspray the test strips with herbicides that may damage the bioassay crop(s). If the bioassay indicates that Prism Herbicide 25% Dry Flowable residues are still present, continue cropping only to those crops listed on the label and do not rotate to other crops until bioassay results indicate that susceptible crops are growing normally.

## NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product in a way that is inconsistent with the directions on the label.

™Trademark of Corteva Agriscience and its affiliated companies.

All other products mentioned are trademarks of their respective companies.