

2023-6567  
2024-12-04

# FERRAMOL®

SLUG AND SNAIL BAIT

COMMERCIAL

READY TO USE

Granules

CONTROLS: Slugs and Snails

KEEP OUT OF REACH OF CHILDREN AND PETS.

EASY TO USE

ACTIVE INGREDIENT: Iron (present as ferric phosphate) 0.28%

Warning: Contains the allergen wheat

READ THE LABEL BEFORE USING

REG NO. 27085 P.C.P. ACT

NET CONTENTS 5-50 kg

W. Neudorff GmbH KG, An der Mühle 3,  
31860 Emmerthal, Germany

**Canadian Agent:** Neudorff North America  
PO Box 178  
Brentwood Bay, B.C., V8M 1R3  
250-652-5888

Ferramol® is a registered trademark of W. Neudorff GmbH KG

2023-6567  
2024-12-04

Neudorff's FERRAMOL is a unique product containing ferric phosphate, which is a compound commonly found in soils, in a specially formulated bait. Any uneaten bait will degrade and become part of the soil.

After ingesting this bait (even in small amounts) slugs and snails cease feeding, which provides immediate protection to plants. Affected slugs and snails die within three to six days. Dead slugs and snails may not be visible as they often crawl away to secluded places to die. Plant protection will be observed in the decrease in plant damage.

PESTS: Controls Deroceras species of slugs such as *D. reticulatum* (field slug), and *D. laeve* (smooth slug); Arion species of slugs such as *A. hortensis* (black field slug), *A. rufus* (large red slug), and *A. ater* (large black slug); and Helix species of snails such as *H. aspersa* (garden snail).

Preharvest Interval: The harvest of crops treated with Ferramol Slug and Snail Bait is permitted on the day of application.

Restricted Entry Interval (REI): DO NOT enter or allow worker entry into treated area for 4 hours or until dusts have settled.

#### **PRECAUTIONS**

KEEP OUT OF REACH OF CHILDREN AND PETS. May cause eye irritation. Avoid contact with eyes. Wash hands after handling. Avoid direct application to ponds, streams and lakes.

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair.

#### **FIRST AID:**

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**If in eyes:** hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

#### **TOXICOLOGICAL INFORMATION**

Treat symptomatically

#### **ENVIRONMENTAL PRECAUTIONS**

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

#### **STORAGE**

2023-6567  
2024-12-04

Store this product away from food or feed.

### DISPOSAL

1. Thoroughly empty the contents of the container into the application device.
2. Follow provincial/territorial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial/territorial requirements.
5. For information on disposal of unused, unwanted product contact the manufacturer or provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for cleanup of spills.

### DIRECTIONS FOR USE

DO NOT apply by air. DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes. As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

Apply the bait around the plants to be protected. For broadcast application, use standard broadcast spreaders. For row application, use standard granular spreaders. Apply bait evenly at 12 – 50 kg per hectare (1.2 – 5.0 grams per square metre) unless otherwise indicated. Apply at the higher rates within the recommended rate range when slug and/or snail pressure is high, if the infestation is severe, if the area is heavily watered or after long periods of heavy rain. Reapply as the bait is consumed or at least every two weeks if slugs and snails continue to cause plant damage. Do not place in piles. If the ground is dry, wet it before applying bait. The soil should be moist but with little or no standing water. See table for specific application directions for crops, greenhouse uses and application types.

WHEN TO APPLY: Evening is the best time to apply the bait, as slugs and snails travel and feed mostly by night or early morning.

WHERE TO APPLY: All likely areas of infestation (i.e., damp places) should be treated. Also treat around the perimeter of crop areas to intercept any slugs or snails travelling into these areas from daytime refuge sites.

Crops		Application Directions
Root and Tuber Vegetables (greenhouse or field)	Beet, Carrot, Celeriac, Ginseng, Horseradish, Jerusalem Artichoke, Parsnip, Potato, Radish, Daikon Radish, Rutabaga, Salsify, Sugar Beets, Sweet Potato, Turnip, Yam	Use broadcast or granular spreader to apply evenly around or over the plants to be protected. At the seedling, transplant and later stages, apply the bait between the rows
Leaves of Root and Tuber Vegetables (greenhouse or	Beet, Carrot, Turnip	

field)		and around the perimeter of the field. For greenhouse grown container plants, scatter the bait in or around the plant pots. Apply about 2.5 g (½ teaspoon) per 23 cm (9") pot.
Crop Group 3 Bulb Vegetables  (greenhouse or field)	Great-headed garlic, Garlic, Dry bulb and Green onion, Leek, Welch onion, Shallot	Use broadcast or granular spreader to apply evenly around or over the plants to be protected. For greenhouse grown container plants, scatter the bait in or around the plant pots. Apply about 2.5 g (½ teaspoon) per 23 cm (9") pot.
Leafy Vegetables (except Brassicas)  (greenhouse or field)	Amaranth, Arugula, Cardoon, Celery, Chervil, Chinese broccoli, Corn salad, Cress, Dandelion, Dock, Endive, Florence Fennel, Lettuce, Orach, Purslane, Radicchio, Rhubarb, Spinach, Swiss chard	Use broadcast or granular spreader to apply evenly around or over the plants to be protected. At the seedling, transplant and later stages, apply the bait between the rows and around the perimeter of the field.
Brassica Leafy Vegetables  (greenhouse or field)	Broccoli, Broccoli raab (rapini), Brussel sprouts, Cabbage, Chinese cabbage (bok choy), Chinese mustard cabbage (gai choy), Cauliflower, Collards, Kale, Kohlrabi, Mizuna, Mustard greens	
Legume Vegetables  (greenhouse or field)	Beans (including dry seed and snap beans), Broad beans, Chickpea, Pea (including snow, snap and shelling peas), Soybean	For greenhouse grown container plants, scatter the bait in or around the plant pots.
Fruiting Vegetables (except cucurbits)  (greenhouse or	Tomato, Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper), Eggplant, Groundcherry, Okra,	Apply about 2.5 g (½ teaspoon) per 23 cm (9") pot.

field)	Pepino, Tomatillo	
Cucurbit Vegetables (greenhouse or field)	Cucumber, Gourds, Muskmelon (includes cantaloupe, casaba, crenshaw and honeydew melons), Pumpkin, Summer squash, Watermelon, Winter squash	
Crop Group 11 Pome Fruits	Apple, Crabapple, Loquat, Mayhaw, Pear, Oriental Pear, Quince	Use broadcast or granular spreader to apply evenly around or over the plants to be protected. For saplings, bait can be applied selectively in a band 15 cm wide around the base of the stem.
Crop Group 12 Stone Fruits	Apricot, Sweet cherry, Tart cherry, Nectarine, Peach, Plum, Chickasaw plum, Damson plum, Japanese plum, Plumcot, Prune	
Berries (greenhouse or field)	Blackberry (including boysenberry), Blueberry, Currant, Cranberry, Elderberry, Gooseberry, Grape, Huckleberry, Kiwifruit, Loganberry, Raspberry, Strawberry	Use broadcast or granular spreader to apply evenly around or over the plants to be protected. The bait can be applied selectively in a 15cm wide band around the base of berry bushes. For greenhouse grown container plants, scatter the bait in or around the plant pots. Apply about 2.5 g (½ teaspoon) per 23 cm (9") pot or 5g (1tsp) for larger pots.
Tree Nuts	Butternut, Chestnut, Filbert, Hickory nut, Pecan, Walnut	Use broadcast or granular spreader to apply evenly around or over the plants to be protected. For

		saplings, bait can be applied selectively in a band 15 cm wide around the base of the stem.
Cereal Grains	Barley, Buckwheat, Corn, Pearl millet, Proso millet, Oats, Popcorn, Rye, Triticale, Wheat, Wild rice	Use broadcast or granular spreader to apply evenly around or over the plants to be protected. At the seedling and later stages, apply the bait between the rows and around the perimeter of the field.
Herbs and Spices (greenhouse or field)	Angelica, Anise, Balm (lemon), Basil, Borage, Burnet, Chamomile, Caraway, Catnip, Celery seed, Chervil, Chive, Chinese Chive, Clary, Coriander, Cilantro, Dillweed, Dill seed, Fennel, Horehound, Hyssop, Juniper berry, Lavender, Lovage, Marigold (calendula), Marjoram, Mustard, Nasturtium, Parsley, Pennyroyal, Poppy (seed), Rosemary, Rue, Sage, Savory, Sweet bay, Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood	Use broadcast or granular spreader to apply evenly around or over the plants to be protected. For greenhouse grown container plants, scatter the bait in or around the plant pots. Apply about 2.5 g (½ teaspoon) per 23 cm (9") pot or 5g (1tsp) for larger pots.
Oilseeds	Canola (rape) seed, Mustard seed, Flax seed, Sunflower seed	Use broadcast or granular spreader to apply evenly around or over the plants to be protected. At the seedling and later stages, apply the bait between the rows and around the perimeter of the field.

Tropical and Subtropical Fruits	Fig	Use broadcast or granular spreader to apply evenly around or over the plants to be protected. For vines and/or saplings, bait can be applied selectively in a 15cm wide band around the base of the stem.
	Hops	
Stalk, Stem and Leaf Petioles	Asparagus	Use broadcast or granular spreader to apply evenly around or over the plants to be protected.
	Globe Artichoke	
Outdoor Ornamentals Including Container-Grown Plants		Use broadcast or granular spreader to apply evenly around or over the plants to be protected. For susceptible plants, shrubs, trees and vines, bait can be applied selectively in a 15cm band around the base of the plant. For container grown plants, scatter the bait in or around the plant pots. Apply about 2.5g (½ teaspoon) per 23 cm (9") pot or 5 g (1 tsp) for larger pots.
Lawns and Turf		Use broadcast or granular spreader

2023-6567  
2024-12-04

		to apply evenly over the area.
--	--	--------------------------------

**NOTICE TO USER**

This pest control product is to be used only in accordance with the directions on the label. It is an offense under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.