

(Container Label)

GROUP	6	HERBICIDE
-------	----------	-----------

IPCO[®]
BROTEX[®]240

**EMUSIFIABLE CONCENTRATE
HERBICIDE**

CONTAINS BROMOXYNIL

For use in WHEAT (Spring and Winter), BARLEY, OATS, FLAX (including low linolenic acid varieties), CORN, FALL RYE, CANARY SEED (for seed production), TRITICALE, GARLIC, ONION (dry bulb only), SEEDLING ALFALFA, ESTABLISHED ALFALFA (for seed production only), and SEEDLING GRASSES, FORAGE SORGHUM, FORAGE MILLET and ZERO TILL

COMMERCIAL (AGRICULTURAL)

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

ACTIVE INGREDIENT: BROMOXYNIL (present as the octanoate ester): 240 g/L

REGISTRATION NO: 28519

PEST CONTROL PRODUCTS ACT

NET CONTENTS: **9.7 litres**



WARNING

POISON

EYE IRRITANT

INTERPROVINCIAL COOPERATIVE LIMITED
P.O. BOX 1050, SASKATOON, SASKATCHEWAN S7K 3M9
1-204-233-3461

[®]T.M. Reg'd.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Causes eye irritation, **DO NOT** get in eyes. Avoid contact with skin and clothing. Harmful or fatal if swallowed. Store the container tightly closed away from seeds, fertilizer, plants and foodstuffs. Wash concentrate from skin or eyes immediately.

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear, and protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair.

Ground applications: Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and chemical-resistant footwear. Gloves are not required during application within a closed cab.

Aerial applications: the field crew and the mixer/loaders: Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, clean-up and repair.

Aerial applicators: Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during application. Gloves are not required during application within a closed cockpit.

DO NOT enter or allow worker entry into treated areas during the restricted-entry intervals (REIs) specified in the following table:

CROP	POST-APPLICATION ACTIVITY	REI
Garlic	Irrigation, handset	2 days
	All other activities	24 hours
Sweet Corn	Irrigation, handset	5 days
	Harvest, hand	20 days
	All other activities	24 hours

DO NOT apply this product in a way that this product will contact workers or other persons, either directly or through drift. Only handlers wearing personal protective equipment may be in the area being treated during application.

Discard clothing or other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. DO NOT reuse them.

Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Avoid breathing spray mist. After use wash hands and other exposed skin. Avoid spray drift onto crops other than those recommended.

Do not use in residential areas, which are defined as sites where bystanders may be present during or after spraying, including homes, schools, parks, playgrounds, playing fields, and public buildings.

Apply only to agricultural crops when the potential for drift to areas of human habitation and areas of human activity such as houses, cottages, schools, and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

Pre-harvest grazing intervals (wheat, barley, oats, forage sorghum, forage millet and seedling alfalfa): Do not use treated crops for grazing of livestock or green feed until 30 days after application of IPCO BROTEX 240 Liquid Herbicide unless otherwise stated on the label. Do not cut treated crops for forage until 30 days after application of IPCO BROTEX 240 Liquid Herbicide.

CAUTION: Do not graze other treated crops or cut for feed unless specified above; sufficient data are not available to support such use.

ENVIRONMENTAL PRECAUTIONS:

Toxic to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under **DIRECTIONS FOR USE**. This product contains an active ingredient and aromatic petroleum distillates, which are toxic to aquatic organisms. To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

FIRST AID:

If swallowed, call a poison control centre or doctor IMMEDIATELY for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin IMMEDIATELY with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

This product contains petroleum distillates. Vomiting may cause aspiration pneumonia. Treat symptomatically.

STORAGE CONDITIONS:

Store this product away from food or feed. This IPCO BROTEX 240 Liquid Herbicide formulation will solidify at temperatures below -20°C but will become useable again at temperatures above 0°C. Insecticides and fungicides should be segregated from herbicides so as to prevent the possibility of cross- contamination.

SHAKE WELL BEFORE USING.

RECYCLABLE CONTAINER DISPOSAL:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

REFILLABLE CONTAINERS DISPOSAL:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

DISPOSAL OF UNUSED, UNWANTED PRODUCT:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

IPCO[®]
BROTEX[®] 240

**EMUSIFIABLE CONCENTRATE
HERBICIDE**

CONTAINS BROMOXYNIL

For use in WHEAT (Spring and Winter), BARLEY, OATS, FLAX (including low linolenic acid varieties), CORN, FALL RYE, CANARY SEED (for seed production), TRITICALE, GARLIC, ONION (dry bulb only), SEEDLING ALFALFA, ESTABLISHED ALFALFA (for seed production only), and SEEDLING GRASSES, FORAGE SORGHUM, FORAGE MILLET and ZERO TILL

COMMERCIAL (AGRICULTURAL)
READ THE LABEL AND BOOKLET BEFORE USING

ACTIVE INGREDIENT: BROMOXYNIL (present as the octanoate ester): 240 g/L

REGISTRATION NO: 28519
PEST CONTROL PRODUCTS ACT

NET CONTENTS: 9.7 litres

WARNING  **POISON**
EYE IRRITANT

INTERPROVINCIAL COOPERATIVE LIMITED
P.O. BOX 1050, SASKATOON, SASKATCHEWAN S7K 3M9
1-204-233-3461

[®]T.M. Reg'd.

LABEL INDEX – IPCO BROTEX 240 LIQUID HERBICIDE

General Information and Directions for Use	Go to Section 1	Triticale	Go to Section 11
Weeds Controlled	Go to Section 2	Garlic	Go to Section 12
Wheat (spring & winter) – not under-seeded to legumes	Go to Section 3	Onions	Go to Section 13
Barley – not under-seeded to legumes	Go to Section 4	Seed and Established Alfalfa	Go to Section 14
Oats – not under-seeded to legumes	Go to Section 5	Minimum or Zero Tillage	Go to Section 15
Corn – field and sweet	Go to Section 6	Summer Fallow	Go to Section 16
Field corn (do not use on sweet corn)	Go to Section 7	Seedling Grasses (not under-seeded to legumes)	Go to Section 17
Fall Rye – not under-seeded to legumes	Go to Section 8	Forage & Sorghum Millet	Go to Section 18
Flax (including low linolenic acid varieties)	Go to Section 9	Mixing Table	Go to Section 19
Canary Seed (for seed production)	Go to Section 10		

SECTION 1: GENERAL INFORMATION AND DIRECTIONS FOR USE

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Timing: For best results spray when weeds are in the seedling stage. Apply in good growing conditions. Application **must** be made before the crop shields the weeds.

Sprayer: Wash the tank and clean all filters, screens and tips. Select nozzle tips to apply the recommended volume of water per hectare. Flood jet type tips are not recommended. Adjust boom height to ensure uniform coverage of weeds. For ground application, spray at 5–10 km/h. Ensure that all tips are in good condition and spraying the same volume.

Mixing: Half fill the tank with clean water. Add the required amount of IPCO BROTEX 240 Liquid Herbicide and agitate thoroughly. Fill the tank and agitate again before use.

Field Sprayer Application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

AERIAL APPLICATION (wheat and barley only):

Aerial Application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length MUST NOT exceed 65% of the wing or rotorspan.

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rate and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Use Precautions: Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the *National Aerial Pesticide Application Manual*, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Do not apply to any body of water. Avoid drifting of spray onto any body of water or other non-target areas. Specified spray buffer zones should be observed.

Coarse sprays are less likely to drift, therefore, avoid combinations of pressure and nozzle type that will result in fine particles (mist). Do not apply during periods of dead calm or when wind velocity and direction pose a risk of spray drift. Do not spray when the wind is blowing towards a nearby sensitive crop, garden, terrestrial habitat (such as shelter-belt) or aquatic habitat.

Operator Precautions: Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders: Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, goggles or face shield, socks and shoes during mixing, loading, clean-up and repair. Aerial applicators: Wear a long-sleeved shirt, long pants, socks and shoes during application. Gloves are not required during application within a closed cockpit. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions: Read and understand the entire label before opening this product. If you have questions, call Interprovincial Cooperative Limited at 1-204-233-3461 or obtain technical advice from the distributor or your provincial agricultural representative.

Application of this specific product must meet and/or conform to the following:

Volume: IPCO BROTEX 240 Liquid Herbicide plus either MCPA or 2,4-D must be applied in volume rates of not *less than* 20 L/ha. For best results when there is a heavy crop canopy, or when the majority of the weeds are cow cockle, green or pale smartweed, hemp-nettle, redroot pigweed or Canada thistle, a volume rate of 40 L/ha is recommended.

Spray Buffer Zones to Protect Sensitive Habitat

A spray buffer zone is NOT required for:

- Uses with hand-held application equipment permitted on this label, low-clearance hooded or shielded sprayers that prevent spray contact with crop, fruit or foliage

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands), and estuarine/marine habitats.

Method of Application	Crop		Spray Buffer Zones (metres) Required for the Protection of:				Terrestrial Habitat
			Freshwater Habitat of Depths:		Estuarine/Marine Habitats of Depths:		
			Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field Sprayer	All crops		1	1	1	1	1
Aerial	Barley and wheat	Fixed wing	20	5	1	1	55
		Rotary wing	20	3	1	1	45

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone calculator on the Pesticides portion of the Canada.ca website.

When tank mixes are permitted, consult the labels of the other tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products included in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

HERBICIDE RESISTANCE MANAGEMENT RECOMMENDATIONS:

For resistance management, IPCO BROTEX 240 Liquid Herbicide is a Group 6 herbicide. Any weed population may contain or develop plants naturally resistant to IPCO BROTEX 240 Liquid Herbicide and other Group 6 herbicides. The resistant

biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of IPCO BROTEX 240 Liquid Herbicide or other Group 6 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact your local Interprovincial Cooperative Limited representative, or call Interprovincial Cooperative Limited at 1-204-233-3461.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Causes eye irritation, **DO NOT** get in eyes. Avoid contact with skin and clothing. Harmful or fatal if swallowed. Store the container tightly closed away from seeds, fertilizer, plants and foodstuffs. Wash concentrate from skin or eyes immediately.

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear, and protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair.

Ground applications: Wear coveralls over a long-sleeved shirt, long pants, chemical-

resistant gloves, socks and chemical-resistant footwear. Gloves are not required during application within a closed cab.

Aerial applications: the field crew and the mixer/loaders: Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, clean-up and repair.

Aerial applicators: Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during application. Gloves are not required during application within a closed cockpit.

DO NOT enter or allow worker entry into treated areas during the restricted-entry intervals (REIs) specified in the following table:

CROP	POST-APPLICATION ACTIVITY	REI
Garlic	Irrigation, handset	2 days
	All other activities	24 hours
Sweet Corn	Irrigation, handset	5 days
	Harvest, hand	20 days
	All other activities	24 hours

DO NOT apply this product in a way that this product will contact workers or other persons, either directly or through drift. Only handlers wearing personal protective equipment may be in the area being treated during application.

Discard clothing or other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. DO NOT reuse them.

Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Avoid breathing spray mist. After use wash hands and other exposed skin. Avoid spray drift onto crops other than those recommended.

Do not use in residential areas, which are defined as sites where bystanders may be present during or after spraying, including homes, schools, parks, playgrounds, playing fields, and public buildings. Apply only to agricultural crops when the potential for drift to areas of human habitation and areas of human activity such as houses, cottages, schools, and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

Pre-harvest grazing intervals (wheat, barley, oats, forage sorghum, forage millet and seedling alfalfa): Do not use treated crops for grazing of livestock or green feed until 30 days after application of IPCO BROTEX 240 Liquid Herbicide unless otherwise stated on the label. Do not cut treated crops for forage until 30 days after application of IPCO

BROTEX 240 Liquid Herbicide.

CAUTION: Do not graze other treated crops or cut for feed unless specified above; sufficient data are not available to support such use.

ENVIRONMENTAL PRECAUTIONS:

Toxic to non-target terrestrial plants. Observe buffer zones specified under **DIRECTIONS FOR USE**. This product contains an active ingredient and aromatic petroleum distillates, which are toxic to aquatic organisms. To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

FIRST AID:

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

This product contains petroleum distillates. Vomiting may cause aspiration pneumonia. Treat symptomatically.

STORAGE CONDITIONS:

Store this product away from food or feed. This IPCO BROTEX 240 Liquid Herbicide formulation will solidify at temperatures below -20°C but will become useable again at temperatures above 0°C. Insecticides and fungicides should be segregated from herbicides so as to prevent the possibility of cross contamination.

SHAKE WELL BEFORE USING.

RECYCLABLE CONTAINER DISPOSAL:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

REFILLABLE CONTAINERS DISPOSAL:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

DISPOSAL OF UNUSED, UNWANTED PRODUCT:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

SECTION 2: Weeds Controlled and Application Timing

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label. In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Interprovincial Cooperative Ltd. at 1-204-233-3461 for information before applying any tank mix that is not specifically recommended on this label.

APPLICATION TIMING:

For best results, spray up to 4-leaf stage of weeds. Apply in good growing conditions. Application must be made before the crop shields the weeds.

WEEDS CONTROL BY IPCO Brotex 240 Liquid Herbicide ALONE

Seedlings up to 4-leaf stage:

Green Smartweed	Common Ragweed
Pale Smartweed	Pigweed****
Lady's-thumb	Cocklebur
Bluebur	Russian thistle**
Kochia**	Wild mustard*
Cow Cockle*	Stinkweed*
Velvet leaf***	American nightshade

Seedlings up to 8-leaf stage:

Wild buckwheat	Lamb's-quarters
Common buckwheat	Common groundsel

Tartary buckwheat

- * Under normal conditions will be controlled up to the 4-leaf stage. Plants beyond this stage are unlikely to be controlled. The higher rate generally gives better control.
- ** Spray before plants are 5 cm high.
- *** Spray before plants are 8 cm high.
- **** Including triazine resistant pigweed

WEEDS CONTROL BY IPCO Brotex 240 Liquid Herbicide + MCPA

Weeds listed for IPCO Brotex 240 Liquid Herbicide alone **plus** these additional weeds:

Seedlings up to 4-leaf stage:

Ball mustard	Canada thistle****
Flixweed	Hemp nettle***
Night flowering catchfly	Perennial sowthistle****
Redroot pigweed	Scentless chamomile*
Shepherd's purse	Volunteer rapeseed/canola**
Volunteer sunflower	

Seedlings up to 8 leaf stage:

Common ragweed	Stinkweed
Wild mustard	Wormseed mustard

- * Spring annuals only. Will not control overwintered weeds.
- ** For high infestations tank-mix 550 g active ingredient MCPA per ha.
- *** Tank-mix 550 g active ingredient MCPA per ha. Plants beyond the 4-leaf stage are not likely to be controlled. Plants emerging after application, which is often the case on peat-type soils, will not be controlled
- **** Top growth control.

WEEDS CONTROL BY IPCO Brotex 240 Liquid Herbicide + 2,4-D

Weeds listed for IPCO Brotex 240 Liquid Herbicide alone plus these additional weeds:

Seedlings up to 4-leaf stage:

Redroot pigweed	Shepherd's purse
Night flowering catchfly	Volunteer sunflower
Flixweed*	Ball mustard

Seedlings up to 8 leaf stage:

Stinkweed*	Wild mustard
------------	--------------

*The higher rate of 2,4-D is recommended for larger over winter weeds (fall rosettes).

SECTION 3 - WHEAT (spring and winter — not underseeded to legumes)

IPCO BROTEX 240 Liquid Herbicide used ALONE	
Weeds Controlled	See Section 2
Spray volume and pressure	Apply in 100 L water per hectare at a pressure of 275 kPa.
Timing	Spring wheat may be treated from the 2-leaf until the early flag leaf stage. Winter wheat may be treated from the 2- to 4-leaf stage in the fall or from the time growth begins to the early flag leaf stage in the spring.
Application rate	1.2–1.4 L/ha. (9.7 L treats 6.9–8.1 ha.) If weeds are beyond the leaf stages indicated on under adverse growing conditions, use of the higher recommended rate will improve control.

IPCO BROTEX 240 Liquid Herbicide + MCPA	
Weeds Controlled	See Section 2 under IPCO Brotex 240 Liquid Herbicide + MCPA
Spray volume and pressure	Apply in 50–100 L water per hectare at a pressure of 275 kPa.
Timing	Spring wheat may be treated from the 2-leaf until the early flag leaf stage. Winter wheat may be treated from the 2- to 4-leaf stage in the fall or from the time growth begins to the early flag leaf stage in the spring.
Application rate	Spring wheat: IPCO BROTEX 240 Liquid Herbicide at 1.2 L/ha (9.7 L treats 8.1 ha) tank-mixed with 275–550 g active ingredient MCPA per hectare where indicated. Winter wheat: IPCO BROTEX 240 Liquid Herbicide at 1.2 L/ha (9.7 L treats 8.1 ha) can be tank-mixed with 275 g active ingredient MCPA per hectare. Refer to table at end of booklet for correct volume of MCPA. Note: MCPA-K is preferred for hemp-nettle although other formulations may be used. Add MCPA to the spray tank first, agitate, then add IPCO BROTEX 240 Liquid Herbicide.

IPCO BROTEX 240 Liquid Herbicide + 2,4-D	
Weeds Controlled	See Section 2
Spray volume and pressure	Apply in 100 L water per hectare at a pressure of 275 kPa.
Timing	Spring and winter wheat may be treated from the 4-leaf until the early flag leaf stage. Application before the 4-leaf stage may injure the crop.
Application rate	IPCO BROTEX 240 Liquid Herbicide at 1.2 L/ha (9.7 L treats 8.1 ha) tank-mixed with 275 - 420 g active ingredient 2,4-D per hectare. Refer to the table at end of booklet for correct volume of 2,4-D.

Note: 2,4-D ester is preferred although other formulations may be used. Add 2,4-D to the spray tank first, agitate and then add IPCO BROTEX 240 Liquid Herbicide.

IPCO BROTEX 240 Liquid Herbicide + Liquid ACHIEVE SC Herbicide	
Weeds Controlled:	See Section 2, plus weeds indicated on the Liquid ACHIEVE SC Herbicide Label
Spray volume and pressure	Apply in 100 L water per hectare at a pressure of 275 kPa.
Timing	Durum wheat, spring wheat and winter wheat may be treated from the 2-leaf until the early flag leaf stage of growth.
Application rate	IPCO BROTEX 240 Liquid Herbicide at 1.2 L/ha (9.7 L treats 8.1 ha) tank mixed with Liquid ACHIEVE SC Herbicide at 0.5 L/ha. When using this tank-mixture, add Turbocharge to the spray tank at a concentration of 0.5% v/v (i. e. 0.5 L of Turbocharge per 100 L of spray solution).

Note: IPCO BROTEX 240 Liquid Herbicide and Liquid ACHIEVE SC Herbicide tank-mixtures can be used in all varieties of spring wheat (including Canada Western red spring, amber durum, soft white spring, extra strong and Canada prairie spring) and red winter wheat.

SECTION 4 - BARLEY (not under-seeded to legumes)

IPCO BROTEX 240 Liquid Herbicide used ALONE	
Weeds Controlled	See Section 2.
Spray volume and pressure	Apply in 100 L water per hectare at a pressure of 275 kPa.
Timing	The crop may be treated from the 2-leaf until the early flag leaf stage.
Application rate	1.2–1.4 L/ha. (9.7 L treats 6.9–8.1 ha.) If weeds are beyond the leaf stages indicated or under adverse growing conditions, use of the higher recommended rate will improve control.

IPCO BROTEX 240 Liquid Herbicide + MCPA	
Weeds Controlled	See Section 2 under IPCO Brotex 240 Liquid Herbicide + MCPA
Spray volume and pressure	Apply in 50–100 L water per hectare at a pressure of 275 kPa.
Timing	The crop may be treated from the 2-leaf until the early flag leaf stage.
Application rate	IPCO BROTEX 240 Liquid Herbicide at 1.2 L/ha (9.7 L treats 8.1 ha) tank-mixed with 275–550 g active ingredient MCPA per hectare where indicated. Refer to the table at end of booklet for correct volume of MCPA.

Note: MCPA-K is preferred for hemp-nettle although other formulations may be used. Add MCPA to the spray tank first, agitate, then add IPCO BROTEX 240 Liquid Herbicide.

IPCO BROTEX 240 Liquid Herbicide + 2,4-D	
Weeds Controlled	See Section 2 under IPCO Brotex 240 Liquid Herbicide + 2,4-D
Spray volume and pressure	Apply in 100 L water per hectare at a pressure of 275 kPa.

Timing	The crop may be treated from the 4-leaf until the early flag leaf stage. Application before the 4-leaf stage may result in injury to the crop.
Application rate	IPCO BROTEX 240 Liquid Herbicide at 1.2 L/ha (9.7 L treats 8.1 ha) tank-mixed with 275–420 g active ingredient 2,4-D per hectare. Refer to table at end of booklet for correct volume of 2,4-D.

Note: 2,4-D ester is preferred although other formulations may be used. Add 2,4-D to the spray tank first, agitate and then add IPCO BROTEX 240 Liquid Herbicide.

IPCO BROTEX 240 Liquid Herbicide + Liquid ACHIEVE SC Herbicide	
Weeds Controlled	See Section 2, plus weeds indicated on the Liquid ACHIEVE SC Herbicide Label.
Spray volume and pressure	Apply in 100 L water per hectare at a pressure of 275 kPa.
Timing	Barley may be treated from the 2-leaf until the early flag leaf stage of growth. Note: IPCO BROTEX 240 Liquid Herbicide and Liquid ACHIEVE SC Herbicide tank-mixtures can be used in all 2 or 6 row varieties of barley (malting and feed varieties).
Application rate	IPCO BROTEX 240 Liquid Herbicide at 1.2 L/ha (9.7 L treats 8.1 ha) tank-mixed with Liquid ACHIEVE SC Herbicide at 0.5 L/ha. When using this tank-mixture, add Turbocharge to the spray tank at a concentration of 0.5% v/v (i. e. 0.5 L of Turbocharge per 100 L of spray solution).

SECTION 5 - OATS (not under-seeded to legumes)

IPCO BROTEX 240 Liquid Herbicide used ALONE	
Weeds Controlled	See Section 2.
Spray volume and pressure	Apply in 100 L water per hectare at a pressure of 275 kPa. DO NOT apply using aerial application equipment.
Timing	The crop may be treated from the 2-leaf until the early flag leaf stage.
Application rate	1.2–1.4 L/ha. (9.7 L treats 6.9–8.1 ha.) If weeds are beyond the leaf stages indicated or under adverse growing conditions, use of the higher recommended rate will improve control.

IPCO BROTEX 240 Liquid Herbicide + MCPA	
Weeds Controlled	See Section 2 under IPCO Brotex 240 Liquid Herbicide + MCPA
Spray volume and pressure	Apply in 50–100 L water per hectare at a pressure of 275 kPa. DO NOT apply using aerial application equipment.
Timing	The crop may be treated from the 2-leaf until the early flag leaf stage.
Application rate	IPCO BROTEX 240 Liquid Herbicide at 1.2 L/ha (9.7 L treats 8.1 ha) tank-mixed with 275–550 g active ingredient MCPA per hectare where indicated. Refer to the table at end of booklet for correct volume of MCPA.

Note: MCPA-K is preferred for hemp-nettle although other formulations may be used. Add MCPA to the spray tank first, agitate, then add IPCO BROTEX 240 Liquid Herbicide.

SECTION 6 - CORN (field and sweet)

IPCO BROTEX 240 Liquid Herbicide used ALONE	
Weeds Controlled	See Section 2
Spray volume and pressure	Apply in 200–300 L water per hectare at a pressure of 275 kPa. DO NOT apply by air.
Timing	Corn may be treated with a broadcast post-emergence application at the recommended rate from the 4-leaf stage onward. To ensure adequate coverage of weeds, drop pipes should be used when corn is beyond the 8-leaf stage or for a second application for later germinating weeds such as cocklebur and velvetleaf. Minimum re-treatment interval for the second application is 21 days. The pre-harvest interval (PHI) is 20 days.
Application rate	1.2–1.4 L/ha. (9.7 L treats 6.9–8.1 ha.) Use of the higher recommended rate will improve control when heavy infestations of weeds are present. IPCO BROTEX 240 Liquid Herbicide is a contact herbicide thus good coverage of the weeds is essential.

Note: Temporary crop injury in the form of leaf scorching may occur in adverse growing conditions (especially if applied during or after periods of cool and wet, or hot and humid weather conditions). **DO NOT ADD OIL OR SURFACTANT.**

IPCO BROTEX 240 Liquid Herbicide + ATRAZINE	
Weeds Controlled	See Section 2, plus weeds indicated on the atrazine label.
Spray volume and pressure	Apply in 200–300 L water per hectare at a pressure of 275 kPa. DO NOT apply by air.
Timing	Corn should be treated from the 4- to 8-leaf stage. Refer to the ATRAZINE label for limitations on grassy weeds.
Application rate	IPCO BROTEX 240 Liquid Herbicide at 1.2 L/ha (9.7 L treats 8.1 ha) tank-mixed with ATRAZINE at 1.1–1.5 kg active ingredient per hectare. Add ATRAZINE to the spray tank first, agitate well, and then add IPCO BROTEX 240 Liquid Herbicide.

Note: Temporary crop injury in the form of leaf scorching may occur in adverse growing conditions (especially if applied during or after periods of cool and wet, or hot and humid weather conditions). DO NOT ADD OIL OR SURFACTANT, OR USE ATRAZINE FORMULATIONS CONTAINING OIL.

IPCO BROTEX 240 Liquid Herbicide + LOW RATE ATRAZINE	
Weeds Controlled	See Section 2, plus seedlings up to 6-leaf stage: redroot pigweed (triazine susceptible), velvet leaf*; seedlings up to 8-leaf stage: common ragweed. * Spray before weed reaches 10 cm in height.
Spray volume and pressure	Apply in 200–300 L water per hectare at a pressure of 275 kPa. DO NOT apply by air.
Timing	Corn may be treated with a broadcast post-emergence application at the recommended rate from the 4-leaf stage onward. To ensure adequate coverage of weeds, drop pipes should be used when corn is beyond the 8-leaf stage or for a second application for later germinating weeds such as cocklebur and velvetleaf. Minimum re-treatment interval for the second application is 21 days. The pre-harvest interval (PHI) is 20 days.
Application rate	IPCO BROTEX 240 Liquid Herbicide at 1.2–1.4 L/ha (9.7 L treats 6.9–8.1 ha) tank-mixed with ATRAZINE at 0.5 kg active ingredient per hectare. Add ATRAZINE to the spray tank first, agitate well and then add IPCO BROTEX 240 Liquid Herbicide.

Note: Temporary crop injury in the form of leaf scorching may occur in adverse growing conditions (especially if applied during or after periods of cool and wet, or hot and humid weather conditions). DO NOT ADD OIL OR SURFACTANT, OR USE ATRAZINE FORMULATIONS CONTAINING OIL.

SECTION 7 - FIELD CORN (do not use on sweet corn)

IPCO BROTEX 240 Liquid Herbicide + BANVEL or BANVEL II (Do not use on sweet corn)	
Weeds Controlled	See Section 2, plus weeds indicated on the BANVEL II OR BANVEL labels.
Spray volume and pressure	Apply in 200–300 L water per hectare at a pressure of 275 kPa. DO NOT apply using aerial application equipment.
Timing	Field corn should be treated from the 4- to 6-leaf stage as an overall broadcast treatment. Drop pipes should be used when applying to corn over the 6-leaf stage and up to 50 cm; direct drop pipe nozzles on the weeds beneath the corn leaves. Apply no later than 2 weeks prior to tassel emergence and do not apply to corn over 50 cm in height.
Application rate	IPCO BROTEX 240 Liquid Herbicide at 1.2 L/ha (9.7 L treats 8.1 ha) tank-mixed with BANVEL or BANVEL II at 290 mL/ha. Add IPCO BROTEX 240 Liquid Herbicide to the spray tank first, agitate well and then add BANVEL or BANVEL II.

Note: Temporary crop injury in the form of leaf scorching may occur in adverse growing conditions (especially if applied during or after periods of cool and wet, or hot and humid weather conditions). DO NOT ADD OIL OR SURFACTANT.

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Interprovincial Cooperative Ltd. under the User Requested Minor Use Label Expansion program. For these uses, Interprovincial Cooperative Ltd. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

IPCO BROTEX 240 LIQUID HERBICIDE + ACCENT 75 DF + NON-IONIC SURFACTANT - FIELD CORN (EASTERN CANADA AND PRAIRIE PROVINCES)

Weeds Controlled	See Section 2, plus weeds indicated on the ACCENT 75 DF label.
Spray volume and pressure	Apply in a minimum of 100 L water per hectare at a pressure of 275 kPa by ground only. DO NOT apply by air.
Timing	Apply as a single post-emergent spray. Apply post-emergence to corn when at the 4- to 8-leaf stage (2–6 visible collars). Do not apply prior to the 4-leaf (2 visible collars) or after the 8-leaf (6 visible collars) stage of corn. Observe a PHI of 30 days.
Application rate	IPCO BROTEX 240 Liquid Herbicide at 1.2 L/ha (9.7 L treats 8.1 ha) tank-mixed with ACCENT 75 DF at 33.4 g/ha and a recommended non-ionic surfactant such as AG-SURF® at 2 L per 1000 L spray solution (0.2% v/v).

SECTION 8 - FALL RYE (not under-seeded to legumes)

IPCO BROTEX 240 Liquid Herbicide used ALONE	
Weeds Controlled	See Section 2.
Spray volume and pressure	Apply in 100 L water per hectare at a pressure of 275 kPa. DO NOT apply by air.
Timing	Fall rye may be treated from the time growth begins to the early flag leaf stage in the spring.
Application rate	1.2–1.4 L/ha. (9.7 L treats 6.9–8.1 ha.) If weeds are beyond the leaf stages indicated or under adverse growing conditions, use of the higher recommended rate will improve control

IPCO BROTEX 240 Liquid Herbicide + MCPA	
Weeds Controlled	See Section 2 under IPCO Brotex 240 Liquid Herbicide + MCPA
Spray volume and pressure	Apply in 50–100 L water per hectare at a pressure of 275 kPa. DO NOT apply by air.
Timing	Fall rye may be treated from the time growth begins to the early flag leaf stage in the spring.
Application rate	IPCO BROTEX 240 Liquid Herbicide at 1.2 L/ha (9.7 L treats 8.1 ha) tank-mixed with 275 g active ingredient MCPA per hectare where indicated. Refer to the mixing tables at end of book for correct volume of MCPA.

Note: MCPA ester is preferred although other formulations may be used. Add MCPA to the spray tank first, agitate, then add IPCO BROTEX 240 Liquid Herbicide.

SECTION 9 - FLAX (including low linolenic acid varieties)

IPCO BROTEX 240 Liquid Herbicide used ALONE	
Weeds Controlled	See Section 2
Spray volume and pressure	Apply in 100 L water per hectare at a pressure of 275 kPa. DO NOT apply by air.
Timing	Flax may be treated from the time it is 5 cm high up to the early flower bud stage but for best results apply IPCO BROTEX 240 Liquid Herbicide when flax is 5–10 cm high.
Application rate	1.2 L/ha. (9.7 L treats 8.1 ha.)

Note: Spraying in the evening may reduce risk of flax injury.

Warning: Do not spray unthrifty crops or when plants are under stress. Do not spray in periods of hot, humid weather. Observe a minimum interval to harvest of 60 days after application of IPCO BROTEX 240 Liquid Herbicide.

Flax is less tolerant of this product than are the cereal crops. Some leaf burn and retarded growth may delay maturity 2–3 days. Avoid using IPCO BROTEX 240 Liquid Herbicide when flax is under stress or during hot (over 29°C) humid weather as severe crop injury may occur.

IPCO BROTEX 240 Liquid Herbicide + MCPA	
Weeds Controlled	See Section 2 under IPCO Brotex 240 Liquid Herbicide + MCPA mixture
Spray volume and pressure	Apply in 100 L water per hectare at a pressure of 275 kPa. DO NOT apply by air.
Timing	Flax may be treated from the time it is 5 cm high up to the early flower bud stage but for best results apply IPCO BROTEX 240 Liquid Herbicide when flax is 5–10 cm high.
Application rate	IPCO BROTEX 240 Liquid Herbicide at 1.2 L/ha (9.7 L treats 8.1 ha) tank-mixed with MCPA at 275 g a.i./ha. The amine or ester formulations of MCPA can be used in tank-mixtures.

Note: Spraying in the evening may reduce risk of flax injury.

Warning: Do not spray unthrifty crops or when plants are under stress. Do not spray in periods of hot, humid weather. Observe a minimum interval to harvest of 60 days after application of IPCO BROTEX 240 Liquid Herbicide.

Flax is less tolerant of this product than are the cereal crops. Some leaf burn and retarded growth may delay maturity 2–3 days. Avoid using IPCO BROTEX 240 Liquid Herbicide when flax is under stress or during hot (over 29°C) humid weather as severe crop injury may occur.

SECTION 10 - CANARY SEED (*Phalaris canariensis*; for seed production)

IPCO BROTEX 240 Liquid Herbicide used ALONE	
Weeds Controlled	See section 2.
Spray volume and pressure	Apply in 100 L water per hectare at a pressure of 275 kPa. DO NOT apply by air.
Timing	The crop may be treated from the 3- to 5-leaf stage.
Application rate	1.2 L/ha. (9.7 L treats 8.1 ha.)

IPCO BROTEX 240 Liquid Herbicide + MCPA	
Weeds Controlled	See Section 2 under IPCO Brotex 240 Liquid Herbicide + MCPA mixture
Spray volume and pressure	Apply in 50–100 L water per hectare at a pressure of 275 kPa. DO NOT apply by air.
Timing	The crop may be treated from the 3- to 5-leaf stage.
Application rate	IPCO BROTEX 240 Liquid Herbicide at 1.2 L/ha (9.7 L treats 8.1 ha) tank-mixed with 275 g active ingredient MCPA per hectare. Refer to the table at end of book for correct volume of MCPA.

Note: MCPA ester is preferred although other formulations may be used. Add MCPA to the spray tank first, agitate, then add IPCO BROTEX 240 Liquid Herbicide.

SECTION 11 – TRITICALE

IPCO BROTEX 240 Liquid Herbicide used ALONE	
Weeds Controlled	See section 2.
Spray volume and pressure	Apply in 100 L water/ha at a pressure of 275 kPa. DO NOT apply by air.
Timing	The crop may be treated from the 2-leaf until the early flag leaf stage.
Application rate	1.2–1.4 L/ha. (9.7 L treats 6.9–8.1 ha.) If weeds are beyond the leaf stage indicated or under adverse growing conditions, use of the higher recommended rate will improve control.

SECTION 12 – GARLIC

IPCO BROTEX 240 Liquid Herbicide used ALONE	
Weeds Controlled	See Section 2.
Spray volume and pressure	Apply in 200–300 L water per hectare at a pressure of 275 kPa. DO NOT apply by air.
Timing	Apply early post emergent to weeds by ground application only. One application per year. Observe a PHI of 58 days.
Application rate	1.2 L/ha. (9.7 L treats 8.1 ha.)

SECTION 13 – ONION (dry bulb only)

<p>The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Interprovincial Cooperative Ltd. under the User Requested Minor Use Label Expansion program. For these uses, Interprovincial Cooperative Ltd. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.</p>			
	Spray Volume and Pressure	Timing	Application Rate
IPCO BROTEX 240 EC used alone	Apply in 200 L water per hectare at a pressure of 170 kPa.	Make two applications per season, at an interval of 10 to 18 days. The first application should be made when onions are at the 2- to 3-leaf stage, and the second application made when onions are at the 4- to 5-leaf stage. DO NOT harvest within 75 days of application.	Apply IPCO BROTEX 240 EC twice per season, each time at a rate of 0.6 L/ha Application Precaution: IPCO BROTEX 240 EC may cause severe leaf burn in onions if weather conditions have not been conducive to the development of the outer waxy layer of the onion leaf.
ONION (Dry Bulb Only) – WEEDS CONTROLLED			
WEEDS CONTROLLED			
IPCO BROTEX 240 EC used alone	Seedlings up to 4-leaf stage: Redroot pigweed, common groundsel.		

**SECTION 14 - SEEDLING and ESTABLISHED ALFALFA (seed production only)
PROVINCES OF ALBERTA, SASKATCHEWAN, MANITOBA ONLY**

IPCO BROTEX 240 Liquid Herbicide used ALONE	
Weeds Controlled	See Section 2.
Spray volume and pressure	Apply in 100 L water/ha at a pressure of 275 kPa. Ground application only. DO NOT apply by air.
Timing	Seedling alfalfa – From the 2 to 6 trifoliolate leaf stage. Do not apply to seedlings under stress . Established alfalfa may be treated until alfalfa is 25 cm tall. Maximum of 2 applications per year. Minimum re-treatment interval for the second application is 21 days.
Application rate	Seedling - 1.2 L/ha. (9.7 L treats 8.1 ha.) Established for seed – 1.2 – 1.4 L/ha (9.7 L treats 6.9–8.1 ha.) If weeds are beyond the leaf stage indicated or under adverse growing conditions, use of the higher recommended rate will improve control.

Note: Use of this product may cause temporary leaf scorch or foliar burn in adverse growing conditions, especially if applied during or after periods of cool and wet, or hot and humid weather conditions. Do not apply if crop is under stress. Do not apply to alfalfa seedlings in the unifoliolate or 1st trifoliolate leaf stage. Avoid overlapping as severe crop injury may occur. In established stands - crop will recover and yield will not be affected.

SECTION 15 - MINIMUM OR ZERO TILL

IPCO BROTEX 240 Liquid Herbicide + GLYPHOSATE HERBICIDE*	
Weeds Controlled	See Section 2, plus weeds controlled on glyphosate labels
Spray volume and pressure	Apply in 100 L of clean water per hectare at a pressure of 275 kPa. DO NOT apply by air.
Timing	This recommendation applies only to minimum or zero till situations before emergence of the crop. Application can be made just prior to or immediately after seeding. Under no circumstances should application be made after crop emergence. Weeds in the seedling stage are most susceptible. This treatment should only be used for minimum or zero-tillage cropping systems in wheat, barley and oats.
Application rate	IPCO BROTEX 240 Liquid Herbicide at 1.2 L/ha (9.7 L treats 8.1 ha) tank-mixed with glyphosate.. A non-ionic surfactant is recommended for use with glyphosate. IPCO BROTEX 240 Liquid Herbicide should be added to the spray tank first, agitate then add glyphosate, agitate and then add surfactant.

***Note:** Glyphosate herbicide (present as potassium salt, isopropylamine salt, ammonium salt, mono-ammonium salt, diammonium salt, dimethylamine salt or trimethylsulfonium

salt) may be tank-mixed with IPCO BROTEX 240 Liquid Herbicide for use in minimum or zero tillage systems.

SECTION 16 – SUMMER FALLOW

IPCO BROTEX 240 Liquid Herbicide + GLYPHOSATE HERBICIDE*	
Weeds Controlled	See Section 2, plus weeds controlled on glyphosate labels
Spray volume and pressure	Apply in 100 L of clean water per hectare at a pressure of 275 kPa. DO NOT apply by air.
Timing	Ensure weeds are at the desired growth stage and actively growing at application for best results. Reduced control may result if weeds are drought stressed. Weeds will continue to germinate from seed throughout the growing season. Repeat treatments may be necessary to control later germinating weeds.
Application rate	IPCO BROTEX 240 Liquid Herbicide at 1.2 L/ha (9.7 L treats 8.1 ha) tank-mixed with glyphosate. A non-ionic surfactant is recommended for use with glyphosate. IPCO BROTEX 240 Liquid Herbicide should be added to the spray tank first, agitate then add glyphosate, agitate and then add surfactant.

***Note:** Glyphosate herbicide (present as potassium salt, isopropylamine salt, ammonium salt, mono-ammonium salt, diammonium salt, dimethylamine salt or trimethylsulfonium salt) may be tank-mixed with IPCO BROTEX 240 Liquid Herbicide for use in summer fallow.

SECTION 17 - SEEDLING GRASSES (not under-seeded to legumes)

IPCO BROTEX 240 Liquid Herbicide used ALONE	
Weeds Controlled	See Section 2.
Spray volume and pressure	Apply in 150 L water per hectare at a pressure of 275 kPa. DO NOT apply by air.
Timing	Seedling grasses may be treated from the 2- to 4-leaf stage.
Application rate	1.2–1.4 L/ha. (9.7 L treats 6.9–8.1 ha.) If weeds are beyond the leaf stage indicated or under adverse growing conditions, use of the higher recommended rate will improve control.

Note: Grasses grown for seed production in the year of establishment only.

Crops: Brome grass, crested wheat grass, intermediate wheat grass, slender wheat grass, tall wheat grass, Russian wild rye, timothy, orchard grass, creeping red fescue, meadow fescue, reed canary grass.

IPCO BROTEX 240 Liquid Herbicide + MCPA	
Weeds Controlled	See Section 2 under IPCO Brotex 240 Liquid Herbicide + MCPA mixture
Spray volume and pressure	Apply in 150 L water per hectare at a pressure of 275 kPa. DO NOT apply by air.
Timing	Seedling grasses may be treated from the 2- to 4-leaf stage.
Application rate	IPCO BROTEX 240 Liquid Herbicide at 1.2 L/ha (9.7 L treats 8.1 ha) tank-mixed with 275 g active ingredient MCPA per hectare. Refer to the table at end of book for correct volume of MCPA.

Note: MCPA ester is preferred although other formulations may be used. Add MCPA to the spray tank first, agitate, then add IPCO BROTEX 240 Liquid Herbicide.

SECTION 18 - FORAGE SORGHUM AND FORAGE MILLET

IPCO BROTEX 240 Liquid Herbicide used ALONE	
Weeds Controlled	See Section 2.
Spray volume and pressure	Apply in 200–300 L water per hectare at a pressure of 275 kPa. DO NOT apply by air.
Timing	Apply when the crop is at or beyond the 4-leaf stage and less than 20 cm in height by ground application only. One application per year. Observe a PHI of 30 days.
Application rate	1.2 L/ha. (9.7 L treats 8.1 ha.)

SECTION 19 - MIXING TABLE:

For calculating the amount of product to add to the spray tank when applying 100 L spray solution per hectare.

		IPCO BROTEX 240 Emulsifiable Concentrate Herbicide (240 g/L)		MCPA AMINE 500 or MCPA ESTER 500		MCPA SODIUM SALT 300	
Rate active per hectare		280 g	340 g	275 g	550 g	275 g	550 g
Rate product per hectare		1.2 L	1.4 L	550 mL	1.1 L	920 mL	1.83 L
Size of spray tank		Litres of chemical to add to spray tank (approx.)					
Gallons	Litres						
500	2273	27.2	31.8	12.5	25.0	20.9	41.6
400	1818	21.8	25.5	10.0	20.0	16.7	33.3
350	1591	19.1	22.3	8.75	17.5	14.6	29.2
300	1363	16.3	19.0	7.5	15.0	12.5	25.0
250	1136	13.6	15.9	6.25	12.5	10.5	20.8
200	909	10.9	12.7	5.0	10.0	8.4	16.6
100	455	5.4	6.3	2.5	5.0	4.2	8.4
50	227	2.7	3.2	1.25	2.5	2.1	4.2
22	100	1.2	1.4	550 mL	1.1	920 mL	1.8
10	45	540 mL	630 mL	250 mL	500 mL	420 mL	820 mL

	IPCO BROTEX 240 Emulsifiable Concentrate		2,4-D AMINE 500		2,4-D 600 LV ESTER		2,4-D 700 LV ESTER	
	Herbicide (240 g/L)							
Rate active per hectare	280 g	340 g	275 g	275 g	275 g	275 g	275 g	275 g
Rate product per hectare	1.2 L	1.4 L	550 mL	550 mL	458 mL	458 mL	393 mL	393 mL
Size of spray tank		Litres of chemical to add to the spray tank (approx.)						
Gallons	Litres							
500	2273	27.2	31.8	12.5	10.4	8.9		
400	1818	21.8	25.5	10.0	8.3	7.1		
350	1591	19.1	22.3	8.75	7.3	6.25		
300	1363	16.3	19.0	7.5	6.25	5.4		
250	1136	13.6	15.9	6.25	5.2	4.5		
200	909	10.9	12.7	5.0	4.2	3.6		
100	455	5.4	6.3	2.5	2.1	1.8		
50	227	2.7	3.2	1.25	1.0	890 mL		
22	100	1.2	1.4	550 mL	458 mL	390 mL		
10	45	540 mL	630 mL	250 mL	210 mL	180 mL		

(Base Label)

GROUP	6	HERBICIDE
--------------	----------	------------------

IPCO[®]

BROTEX[®] 240

**EMULSIFIABLE CONCENTRATE
HERBICIDE**

CONTAINS BROMOXYNIL

For use in WHEAT (Spring and Winter), BARLEY, OATS, FLAX (including low linolenic acid varieties), CORN, FALL RYE, CANARY SEED (for seed production), TRITICALE, GARLIC, ONION (dry bulb only), SEEDLING ALFALFA, ESTABLISHED ALFALFA (for seed production only), and SEEDLING GRASSES, FORAGE SORGHUM, FORAGE MILLET and ZERO TILL

COMMERCIAL (AGRICULTURAL)

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

ACTIVE INGREDIENT: BROMOXYNIL (present as the octanoate

ester): 240 g/L REGISTRATION NO: 28519

PEST CONTROL PRODUCTS ACT

NET CONTENTS: **9.7 litres**



WARNING

POISON

EYE IRRITANT

INTERPROVINCIAL COOPERATIVE LIMITED
P.O. BOX 1050, SASKATOON, SASKATCHEWAN S7K 3M9
1-204-233-3461

[®]T.M. Reg'd.