

(Container)

Maverick™ III

Herbicide

GROUP	9	HERBICIDE
--------------	----------	------------------

Water soluble herbicide for nonselective weed control in CROPLAND SYSTEMS AND IN NON-CROPLAND AREAS.

AGRICULTURAL and INDUSTRIAL

**READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

ACTIVE INGREDIENT: Glyphosate (present as dimethylamine salt) 480 g/L
solution

REGISTRATION NO. 28977 PEST CONTROL PRODUCTS ACT

**CAUTION: EYE AND SKIN IRRITANT
POTENTIAL SKIN SENSITIZER**

Net Contents: 10L

Albaugh, LLC
1525 NE 36th Street
Ankeny, IA 50021
USA
1-800-247-8013

PRECAUTIONS

May irritate eyes and skin

Avoid contact with eyes or with skin

KEEP OUT OF REACH OF CHILDREN

DO NOT APPLY BY AIR

Wear long sleeved shirt, long pants and chemical resistant gloves during mixing, loading, application, clean up and repair. In addition, wear goggles or a face shield during mixing and loading.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind directions, temperature inversions, application equipment and sprayer settings.

PHYSICAL OR CHEMICAL HAZARDS

Spray solutions of this product should be mixed, stored and applied only in stainless steel, aluminum, fibreglass, plastic and plastic-lined steel containers. **DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.** This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

No specific antidote. Employ supportive care. Treatment should be based on judgement of the physician in response to reactions of the patient.

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL HAZARDS

Avoid direct applications to any body of water. Do not contaminate water by disposal of waste or cleaning of equipment. TOXIC to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE. To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

STORAGE

Avoid contamination of seed, feed, and foodstuffs.

Soak up small amounts of spill with absorbent clays.

DISPOSAL

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

(Booklet)

Maverick™ III

Herbicide

GROUP	9	HERBICIDE
--------------	----------	------------------

Water soluble herbicide for nonselective weed control in CROPLAND SYSTEMS AND IN NON-CROPLAND AREAS.

AGRICULTURAL and INDUSTRIAL

**READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

ACTIVE INGREDIENT: Glyphosate (present as dimethylamine salt) 480 g/L
solution

REGISTRATION NO. 28977 PEST CONTROL PRODUCTS ACT

**CAUTION: EYE AND SKIN IRRITANT
POTENTIAL SKIN SENSITIZER**

Net Contents: 7.5 L, 10 L, 115L, 450L, 960 L and Bulk

Albaugh, LLC
1525 NE 36th Street
Ankeny, IA 50021
USA
1-800-247-8013

TABLE OF CONTENTS

PRECAUTIONS

DIRECTIONS FOR USE

GENERAL INFORMATION

MIXING AND APPLICATION

Precautions

Mixing and Application Equipment Information

WEEDS CONTROLLED

Annual Weeds

Perennial Weeds

Woody Brush and Trees

CROPLAND USES

ANNUAL WEED CONTROL

Annual Weed Control with Maverick III Herbicide

Annual Weed Control with Maverick III Herbicide Tank Mixtures for Summerfallow & Minimum Tillage Systems

Addition of Surfactant

Additional Important Information For Annual Weed Control

Weed Control in Glyphosate Tolerant Canola

Maverick III Herbicide plus Lontrel™360

Weed Control in Glyphosate Tolerant Soybean

Weed Control in Glyphosate Tolerant Corn

PERENNIAL WEED CONTROL

Perennial Weed Control With Maverick III Herbicide

Special Notes for Perennial Weed Control:

Quackgrass

Surfactant Information

Canada Thistle

Toadflax

Dandelion

Alfalfa Control with 2,4-D Tank-Mix

All Perennial Weeds

CROPLAND SITUATIONS

Prior to Planting - All Crops

Post Harvest Stubble Treatment

Spot treatment In-Crop

Grazing Restrictions

Summerfallow Treatment

Minimum and Zero Tillage Cropping Systems

Minimum and Zero Tillage Tank Mixtures

Maverick III Herbicide plus Pardner

Maverick III Herbicide plus Pursuit

Forage Legumes and Grasses

Pasture Renovation

Forage Seed Production

Preharvest Control of Quackgrass, Canada Thistle, Milkweed, Toadflax and Dandelion

Timing of Preharvest Applications

Tree Plantings

Tree, Vine and Berry Crops

Selective Equipment

NON-CROPLAND USES

Weed Control In Non-Cropland Areas with Maverick III Herbicide
Application Information for Non-Cropland Uses
Ground Applications
Purple Loosestrife Control
Selective Equipment for Non-Cropland Uses
Turfgrass
Injection Applications
Cut Stump Application
Resistance Management Recommendations

PRECAUTIONS

May irritate eyes and skin
Avoid contact with eyes or with skin

KEEP OUT OF REACH OF CHILDREN

DO NOT APPLY BY AIR

Wear long sleeved shirt, long pants and chemical resistant gloves during mixing, loading, application, clean up and repair. In addition, wear goggles or a face shield during mixing and loading.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind directions, temperature inversions, application equipment and sprayer settings.

PHYSICAL OR CHEMICAL HAZARDS

Spray solutions of this product should be mixed, stored and applied only in stainless steel, aluminum, fibreglass, plastic and plastic-lined steel containers. **DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.** This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

No specific antidote. Employ supportive care. Treatment should be based on judgement of the physician in response to reactions of the patient.

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL HAZARDS

Avoid direct applications to any body of water. Do not contaminate water by disposal of waste or cleaning of equipment. **TOXIC** to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE. To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

STORAGE

Avoid contamination of seed, feed, and foodstuffs.
Soak up small amounts of spill with absorbent clays.

DISPOSAL

Recyclable Containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

Returnable Containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable Containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

DIRECTIONS FOR USE

GENERAL INFORMATION

The restricted entry interval is 12 hours after application for all agricultural uses.

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests. **DO NOT** contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Glyphosate is not to be applied using hand-wicking or hand-daubing methods.

Overspray or drift to important wildlife habitats such as bodies of water, wetlands (e.g., sloughs), shelterbelts, woodlots and other cover on the edges of fields frequented by wildlife, should be avoided. Leave a 15 meter buffer zone between the last spray swath and the edge of any of these habitats.

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. Boom height must be 60 cm or less above the crop or ground.

Airblast or mist blower application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** direct spray above plants to be treated. **DO NOT** apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side. For airblast applications, turn off outward pointing nozzles at row ends and outer rows.

Aerial application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 km/h at flying height at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. To reduce drift caused by turbulent wingtip vortices, the nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotorspan.

544

Maverick III Herbicide, a water soluble liquid, mixes readily with water for application as a foliage spray for the control or destruction of most herbaceous plants. It may be applied through most standard industrial or field type sprayers after dilution and thorough mixing with water in accordance with the booklet instructions.

This herbicide moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days but on most perennial weeds may not occur until 7 to 10 days. Extremely cool or cloudy weather at treatment time may slow down activity of this product and delay visual effects of control. Visible effects are a gradual wilting and yellowing of the plant which advances to complete browning of above ground growth and deterioration of underground plant parts.

Delay application until vegetation has emerged to the stages described for control of such vegetation under the annual and perennial weed control sections of this booklet to provide adequate leaf surface to receive the spray. Unemerged plants arising from underground rhizomes or root stocks of perennials will not be affected by the spray and will continue to grow. For this reason best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity.

Always use the higher rate of this product per hectare within the recommended range when weed growth is heavy or dense, or weeds are growing in an undisturbed (non-cultivated) area.

Do not treat weeds under poor growing conditions such as drought stress, disease or insect damage, as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust.

This product does not provide residual weed control. For subsequent residual weed control follow a label approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

Heavy rainfall immediately after application may wash the chemical off the foliage and a repeat treatment may be required. Do not apply if rainfall is forecast for the time of application.

Do not mix with any surfactant, pesticide, herbicide oils or any other material other than water unless specified in this booklet. For best results, spray coverage should be uniform and complete. Do not spray weed foliage to the point of runoff.

MIXING AND APPLICATION

PRECAUTIONS

ATTENTION: AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES.

AVOID DRIFT - EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURING DESIRABLE PLANTS AND CROPS. Even minute quantities of spray drift can cause severe damage or destruction to nearby crops, plants or other areas on which treatment is not intended, or may cause other unintended consequences.

DO NOT USE IN GREENHOUSES. REDUCED RESULTS MAY OCCUR IF WATER CONTAINING SOIL IS USED, SUCH AS WATER FROM PONDS AND UNLINED DITCHES.

Clean sprayer and parts immediately after using this product by thoroughly flushing with water. Do not contaminate water sources by disposal of wastes or cleaning of equipment.

NOTE: Use of this product in any manner not consistent with this booklet may result in injury to persons, animals or crops, or other unintended consequences. Keep container closed to prevent spills and contamination.

MIXING AND APPLICATION EQUIPMENT INFORMATION

MIXING

For ground or industrial type sprayers, fill the spray tank with one-half the required amount of water. Add the proper amount of herbicide (see "Weed Control" sections of this booklet) and mix well before adding the remaining portion of water. Placing the filling hose below the surface of the liquid solution will prevent excessive foaming. Removing hose from tank immediately will avoid back siphoning into water source. Use of mechanical agitators may cause excessive foaming. Bypass lines should terminate at the bottom of the tank.

For use in knapsack sprayers, it is suggested that the proper amount of this herbicide be mixed with water in a larger container. Fill sprayer with the mixed solution.

APPLICATION EQUIPMENT

BOOM EQUIPMENT

For control of perennial weeds and woody brush and trees listed on this booklet using conventional boom equipment--, Apply this product in 50 to 300 L of clean water per hectare as a broadcast spray using no more than 275 kPa pressure. See "**Weed Control**" sections of this booklet for rates to control specific weeds.

For control of annual weeds listed on this booklet using conventional boom equipment--Apply this product in 50 to 100 L of clean water per hectare as a broadcast spray, except as otherwise stated on this label using no more than 275 kPa pressure. See "**Weed Control**" sections of this booklet for rates to control specific weeds.

HAND HELD AND HIGH VOLUME EQUIPMENT

(use coarse sprays only)

For control of weeds and woody brush and trees listed in the "Weed Controlled" section of this label using knapsack sprayers or high volume spraying equipment utilizing handguns or other suitable nozzle arrangements. Unless otherwise specified, make a 0.75% solution of this product in water (0.75 litre of this product in 100 litres of water) and apply to foliage of vegetation to be controlled. For best results, use a 1.5% solution (1.5 litres of this product in 100 litres of water) on harder to control perennials such as field bindweed, hemp dogbane, milkweed and Canada thistle.

Applications should be made on a spray-to-wet basis. Spray coverage should be uniform and complete. Do not spray to point of runoff. Hand gun applications should be properly directed to avoid spraying desirable plants.

SELECTIVE EQUIPMENT

Selective equipment such as **WIPER** and **ROLLER** applicators can be used for weed control in soy and dry beans, orchards, vineyards, cranberries, strawberries and non-crop areas. For information regarding use of this product with selective equipment, refer to "**Selective Equipment**" section of this label.

WEEDS CONTROLLED

This product controls many annual and perennial grasses, broadleaf weeds, and woody brush and trees when applied as recommended and under conditions described. For information on how to control specific weeds including herbicide rate refer to the "**Annual Weed Control**" and "**Perennial Weed Control**" sections of this label. The following is a partial list of weeds controlled:

ANNUAL WEEDS

Annual Grasses

Barnyard Grass

Echinochloa crusgalli

Blue Grass (annual)

Poa annua

Crab Grass (large)

Digitaria sanguinalis

Crab Grass (smooth)

Digitaria ischaemum

Downy Brome

Bromus tectorum

Fall Panicum

Panicum dichotomiflorum

Giant Foxtail

Setaria faberii

Green Foxtail

Setaria viridis

Persian Darnel

Lolium persicum

Volunteer Barley

Hordeum spp.

Volunteer Corn

Zea Mays

Volunteer Wheat

Triticum spp.

Wild Oats

Avena fatua

Wild Proso Millet

Panicum miliaceum

Yellow Foxtail

Setaria glauca

Other

Dodder

Cuscuta spp.

Annual Broadleaf Weeds

Chickweed

Stellaria media

Cleavers

Galium aparine

Cocklebur

Xanthium strumarium

Corn Spurry

Spergula arvensis

Cowcockle

Saponaria vaccaria

Eastern Black Flowering Nightshade

Solanum ptycanthum

Fleabane (Canada)

Erigeron canadensis

Flixweed

Descurania sophia

Green Smartweed

Polygonum scabrum

Hempnettle

Galeopsis tetrahit

Kochia

Kochia scoparia

Lady's-Thumb

Polygonum persicaria

Lamb's-Quarters (common)

Chenopodium album

Narrow-leaved Hawk's Beard

Crepis tectorum

Narrow-leaved Vetch

Vicia angustifolia

Night-flowering Catchfly

Silene noctiflora

Pennsylvania Smartweed

Polygonum pennsylvanicum

Prickly Lettuce

Lactuca scariola

Ragweed (common)

Ambrosia artemisiifolia

Redroot Pigweed

Amaranthus retroflexus

Round-Leaved Mallow

Malva pusilla

Russian Thistle

Salsola pestifer

Shepherd's Purse

Capsella bursa-pastoris

Smooth Pigweed

Amaranthus hybridus

Sowthistle (annual)

Sonchus oleraceus

Stinkweed

Thlaspi arvense

Storksbill

Erodium cicutarium

Volunteer Canola

Brassica spp

Volunteer Flax

Linaria spp

Wild Buckwheat

Polygonum convolvulus

Wild Mustard

Sinapsis arvensis

Wild Tomato

Solanum triflorum

Velvetleaf

Abutilon theophrasti

PERENNIAL WEEDS

Perennial Grasses / Sedges

Blue Grass (Canada)

Poa compressa

Blue Grass (Kentucky)

Poa pratensis

Brome Grass (smooth)

Bromus inermis

Cattail (common)

Typha latifolia

Foxtail Barley

Hordeum jubatum

Quackgrass

Agropyron repens

Yellow Nutsedge

Cyperus esculentus

Wire-stemmed Muhly

Muhlenbergia frondosa

Perennial Broadleaved Weeds

Alfalfa

Medicago spp.

Cottontop

Eriophorum chamissionis

Curled Dock

Rumex crispus

Dandelion

Taraxacum officinale

Field Bindweed

Convolvulus arvensis

Hemp Dogbane

Apocynum cannabinum

Hoary Cress

Cardaria draba

Knotweed (Japanese)

Polygonum cuspidatum

Milkweed (common)

Asclepias syriaca

Poison Ivy

Rhus radicans

Purple Loosestrife

Lythrum salicaria

Sow Thistle (perennial)

Sonchus arvensis

Thistle (Canada)

Cirsium arvense

Toad Flax

Linaria vulgaris

Wormwood (Absinth)

Artemisia absinthium

WOODY BRUSH AND TREES

Alder

Alnus spp.

Birch

Betula spp.

Broadleaved meadowsweet

Spiraea latifolia

Canadian rhododendron

Rhododendron canadenses

Cedar

Thuja spp.

Cherry

Prunus spp.

Douglas Fir

Pseudotsuga spp.

Hemlock

Tsuga spp.

Maple

Acer spp.

Mountain-fly honeysuckle

Lonicera villosa

Pine

Pinus spp.

Poplar

Populus spp.

Raspberry / Salmonberry

Rubus spp.

Sheep laurel

Kalmia angustifolia

Snowberry (Western)

Symphoricarpos occidentalis

Sweet fern

Comptonia peregrina

Willow

Salix spp.

Withrod

Viburnum cassinoides

CROPLAND USES

CROPLAND USES INCLUDE:

In cropping systems before planting of all crops; in minimum tillage systems, post emergent in glyphosate tolerant soybean, canola and corn; preharvest applications in wheat, barley, oats, canola (rapeseed), flax (including low linolenic acid varieties), peas, lentils, dry beans, soybeans and forages; in pasture renovation; in forage, legume and grass establishments; in tree crops including apple, pear, cherry, plum, peach, apricot, filbert, hazelnut, walnut, chestnut; in grapes, cranberries, blueberries and strawberry; in sugar beets; in tree plantings; and grasses for seed production.

ALWAYS READ PRECAUTIONARY STATEMENTS, GENERAL INFORMATION and MIXING and APPLICATION PRECAUTIONS PRIOR TO SPECIFIC APPLICATION INFORMATION IN ANY LABEL SECTION.

ANNUAL WEED CONTROL

The following tables provide rates and specific application instructions for control of the annual weeds listed.

ANNUAL WEED CONTROL WITH MAVERICK III HERBICIDE

RATE L/HA	GROWTH STAGE	WEEDS CONTROLLED	COMMENTS (Apply in 50 - 100 L/ha water)
0.56	weeds up to 8 cm in height	wild oats, green foxtail, volunteer barley, volunteer wheat volunteer canola (rapeseed), wild mustard, lady's-thumb, stinkweed	.for wild oats apply at 1 - 3 leaf stage .add 350 mL of the surfactant Agral 90, or Ag Surf, or Companion™ . .for heavy wild oat infestations use 0.75 L/ha rate.
0.75	weeds 8 cm to 15 cm in height	all annual grasses listed above all annual broadleaved weeds listed above plus flixweed and Kochia	.add 350 mL of surfactant registered for use as listed above. † suppression only. Refer to higher rates of this table or tank-mix table for control options.
0.94 - 1.4	weeds up to 15 cm in height	all annual grasses listed above plus downey brome, giant foxtail, and Persian dandelion all annual broadleaved weeds listed above plus lamb's quarters, redroot pigweed, hempnettle, flixweed, Russian thistle, volunteer flax, common ragweed, Canada fleabane, wild buckwheat†, narrow-leaved hawk's beard†	.no surfactant required for tank-mix weed control options see annual weed control with tank mixture section † DO NOT use these rates on plants greater than 8 cm in height †† for 3 - 4 leaf stage use 1.4 L/ha rate ††† for weeds 8 cm to 15 cm in height use 1.4 L/ha rate
1.69	weeds up to 15 cm in height	all annual grasses listed above plus crab grass and annual blue grass. all annual broadleaved	.for additional annual broadleaved weed control options, refer to tank-mix table

		weeds listed above plus kochia, prickly lettuce, shepherd's purse, annual sow thistle, and narrow-leaved vetch	
2.63	weeds over 15 cm in height	all annual grasses and broadleaved weeds listed above	.for additional annual broadleaved weed control options refer to tank-mix table

NOTE: For spot treatment, 0.56 – 2.63 L/ha is approximately equivalent to 6-26 mL/100 m², respectively.

ANNUAL WEED CONTROL WITH MAVERICK III HERBICIDE TANK MIXTURES FOR SUMMERFALLOW & MINIMUM TILLAGE SYSTEMS

TANK MIXTURES	RATE L/ha	WEEDS CONTROLLED	COMMENTS (Apply in 50 - 100 L/ha water; add 350mL/ha of surfactant)
Maverick III Herbicide + Banvel II	0.56– 0.75 + 0.29	Volunteer cereal, wild oats, green foxtail, Volunteer canola (rapeseed), wild mustard, flixweed, lamb's quarters, lady's thumb, stinkweed, kochia, Russian thistle, cow cockle, redroot pigweed†, wild buckwheat†.	This tank-mix is registered for summerfallow use only . Weeds should be less than 15 cm tall and actively growing for best results. Use higher rate if weeds are beyond 8 cm in height. † Maverick III Herbicide applied at 0.75 L/ha rate only. †† Suppression only. See other tank mixtures for control options.
Maverick III Herbicide + Pardner	0.56– 0.75 + 1.25	Volunteer cereals, green foxtail, volunteer canola (rapeseed), wild mustard, lady's thumb, stinkweed, wild buckwheat Redroot pigweed†, kochia†, wild oats†	This tank-mix is registered only for use in summerfallow, and prior to wheat, oats and barley in minimum tillage systems . Weeds should be less than 15 cm tall and actively growing for best results. Use higher rate if weeds are beyond 8 cm in height. † use Maverick III Herbicide at 0.75 L/ha rate only for wild buckwheat control. †† 0.75 L rate, suppression only. See other tank mixtures for control options.
Maverick III Herbicide + 2,4-D#	0.56– 0.75 + 1.2	Volunteer cereals, wild oats† and green foxtail, volunteer canola (rapeseed), wild mustard, flixweed, redroot pigweed, lady's thumb, stinkweed, kochia. Lamb's quarters††, Russian thistle†.	This tank-mix is registered for summerfallow use only . Weeds should be less than 15 cm tall and actively growing for best results. Use higher rate if weeds are beyond 8 cm in height † use Maverick III Herbicide at 0.75 L/ha rate only for wild oat and green foxtail control. †† suppression only. See other tank mixtures

<p>Maverick III Herbicide + MCPA### 500 g/L formulation, if another formulation is used, adjust rate accordingly</p>	<p>0.94-1.4 + 0.5– 0.7¹ OR 0.5– 1.0²</p>	<p>Volunteer cereals, wild oats, green foxtail, downy brome, giant foxtail, and Persian darnel.</p> <p>Volunteer canola (rapeseed) (non-glyphosate tolerant), wild mustard, flixweed, redroot pigweed, lady's thumb, stinkweed, kochia, lamb's quarters, hempnettle, Russian thistle, volunteer flax, common ragweed, Canada fleabane, wild buckwheat[†], narrowleaved hawk's beard^{†††}</p> <p>Volunteer glyphosate tolerant canola (1-4 leaf stage)^{1,2}, bluebur³, burdock³ (before 4 leaf stage), false flax³, flixweed³, lamb's quarters³, mustards³ (except dog and tansy), prickly lettuce³, ragweeds³, redroot pigweed³, Russian pigweed³, shepherd's purse³, stinkweed (field pennycress)³, vetch³, wild radish³, wild sunflower³</p>	<p>for control options.</p> <p>Weeds should be less than 15 cm tall and actively growing for best results.</p> <p>Use higher rate if weeds are beyond 8 cm in height.</p> <p>No surfactant required.</p> <p>† DO NOT use these rates on plants greater than 8 cm in height.</p> <p>†† For 3-4 leaf stage use 1.4 L/ha rate.</p> <p>††† For weeds 8 cm to 15 cm in height use 1.4 L/ha rate.</p> <p>¹MCPA amine at 0.5– 0.7 L/ha (250-350 g ai/ha) prior to peas.</p> <p>² MCPA at 0.5– 1.0 L/ha (250-500 g ai/ha) prior to wheat, barley, oats, corn (field and sweet)###, rye and flax.</p> <p>³ MCPA at 0.7– 1.0 L/ha (350– 500 g ai/ha) only.</p> <p>Use this tank mix prior to seeding in wheat, barley, rye, oats, corn (field and sweet)###, flax, and field peas###.</p>
<p>Maverick III Herbicide + Buctril M herbicides</p>	<p>0.94-1.4 + 0.5– 1.0¹</p>	<p>Volunteer cereals, wild oats, green foxtail, downy brome, giant foxtail, and Persian darnel.</p> <p>Volunteer canola (rapeseed) (non-glyphosate tolerant), wild mustard, flixweed, redroot pigweed, lady's thumb, stinkweed, kochia, lamb's quarters, hempnettle, Russian thistle, volunteer flax, common ragweed, Canada fleabane, wild buckwheat[†], narrowleaved hawk's beard^{†††}</p> <p>Volunteer glyphosate tolerant Canola (1-4 leaf stage)^{1,2}</p> <p>Seedlings up to the 4-leaf stage²: green smartweed,</p>	<p>Weeds should be less than 15 cm tall and actively growing for best results.</p> <p>Use higher rate if weeds are beyond 8 cm in height.</p> <p>No surfactant required.</p> <p>† DO NOT use these rates on plants greater than 8 cm in height.</p> <p>†† For 3-4 leaf stage use 1.4 L/ha rate.</p> <p>††† For weeds 8 cm to 15 cm in height use 1.4 L/ha rate.</p> <p>¹ Buctril M at 0.5– 1.0 L/ha (280- 560 g ai/ha) for all crops listed.</p> <p>² Buctril M at 1.0 L/ha (560 g ai/ha only).</p> <p>³ Spray before plants are 5 cm high.</p> <p>⁴ Spring annuals only.</p>

		<p>pale smartweed , lady's thumb, cow cockle, redroot pigweed, flixweed, bluebur, shepherd's purse, kochia, Russian thistle³, scentless chamomile⁴, volunteer sunflower, night flowering catchfly, cocklebur, velvetleaf⁵, ball mustard, American nightshade</p> <p>Seedlings up to the 6-leaf stage²: wild tomato Seedlings up to the 8-leaf stage²: wild buckwheat, tartary buckwheat, common buckwheat, stinkweed, wild mustard, wormseed mustard, lamb's quarters, common ragweed, common groundsel</p> <p>Perennials (top growth)²: Canada thistle, perennial sow thistle</p>	<p>⁵ Spray before plants are 8 cm high.</p> <p>Use this tank mix prior to seeding in wheat, barley, rye, oats, corn, flax, canary seed and seedling grasses (including brome grass, crested wheatgrass, intermediate wheat grass, slender wheatgrass, tall wheatgrass, Russian wild rye, Timothy, orchard grass, creeping red fescue, meadow fescue, meadow foxtail, seedling tall fescue, seedling meadow bromegrass, seedling streambank wheatgrass and reed canary grass.</p>
<p>Maverick III Herbicide + MCPA Amine (500 g/L formulation, if another formulation is used, adjust rate accordingly)</p>	<p>0.94-1.4 + 0.5_ 0.7</p>	<p>Volunteer cereals, wild oats, green foxtail, downy brome, giant foxtail, and Persian darnel.</p> <p>Volunteer canola (rapeseed) (non-glyphosate tolerant), wild mustard, flixweed, redroot pigweed, lady's thumb, stinkweed, kochia, lamb's quarters, hempnettle, Russian thistle, volunteer flax, common ragweed , Canada fleabane, wild buckwheat[†], narrowleaved hawk's beard^{†††}</p> <p>Volunteer glyphosate tolerant canola (1-4 leaf stage)³, bluebur⁴, burdock⁴ (before 4 leaf stage), false flax⁴, flixweed⁴, lamb's quarters⁴, mustards⁴ (except dog and tansy), prickly lettuce⁴, ragweeds⁴, redroot pigweed⁴, Russian pigweed⁴, shepherd's purse⁴, stinkweed⁴ (field pennycress), vetch⁴, wild radish⁴, wild sunflower⁴</p>	<p>Weeds should be less than 15 cm tall and actively growing for best results.</p> <p>Use higher rate if weeds are beyond 8 cm in height.</p> <p>No surfactant required.</p> <p>† DO NOT use these rates on plants greater than 8 cm in height.</p> <p>†† For 3-4 leaf stage use 1.4 L/ha rate.</p> <p>††† For weeds 8 cm to 15 cm in height use 1.4 L/ha rate.</p> <p>³ MCPA amine at 0.5_ 0.7 L/ha (250_ 350 g ai/ha) prior to lentils and chickpeas.</p> <p>⁴ MCPA amine at 0.7 L/ha (350 g ai/ha) only.</p> <p>• Use this tank mix prior to seeding in lentil and chickpea.</p>

For foxtail barley suppression, refer to “**Annual Weed Control**” table

#0.56 kg ai/ha of 2,4-D.

#, ## Adjust rates accordingly for other 2,4-D formulations. Use only low volatile ester or amine formulations of 2,4-D.

Use only amine formulations of MCPA prior to seeding in corn and field peas.

ADDITION OF SURFACTANT

All Maverick III Herbicide tank mixtures for annual weed control require the addition of the surfactant Agral 90, or Ag Surf, or Companion. Surfactant should be added at a rate of 350 mL per hectare, in 50-100 L of clean water.

Additional Important Information for Annual Weed Control

Allow at least 1 day after treatment before tillage

Annual weeds generally will continue to germinate from seed throughout the growing season. Repeat treatments may be necessary to control later germinating weeds, in some situations.

For additional information and precautions, refer to the "General Information" and "Mixing and Application" sections of this label.

WEED CONTROL IN GLYPHOSATE TOLERANT CANOLA

WARNING: APPLY MAVERICK III HERBICIDE ON GLYPHOSATE TOLERANT CANOLA VARIETIES ONLY.

NOTE: ALWAYS USE PEDIGREED (I.E. CERTIFIED) GLYPHOSATE TOLERANT CANOLA SEED. CANOLA WHICH IS NOT DESIGNATED AS GLYPHOSATE TOLERANT WILL BE DAMAGED OR DESTROYED BY THIS TREATMENT.

- For additional information and precautions refer to the “General Information” and “Mixing and Application” sections of the Maverick III Herbicide label.
- Apply Maverick III Herbicide in glyphosate tolerant canola only as directed in the following weed control table.
- Some short-term, visual yellowing may occur when Maverick III Herbicide is applied at the late application 4 to 6 leaf stage of the crop. This effect is temporary and will not influence crop growth, maturity or yield.

DO NOT APPLY BY AIR

The following table describes the rate and specific application instructions for control of annual and perennial weeds in glyphosate tolerant canola varieties.

WEED CONTROL IN GLYPHOSATE TOLERANT CANOLA

Rate (L/ha)	Growth Stage Of Crop	Weeds Controlled	Comments (Apply in 50 - 100 L/ha water)
0.60 - 1.4	0 to 6 leaf	Annual Grasses wild oats, green foxtail, volunteer barley, volunteer wheat, barnyard grass Annual Broadleaves stinkweed, redroot pigweed, wild mustard, Russian thistle, lamb's-quarters, non-glyphosate tolerant volunteer canola (rapeseed), hempnettle, lady's thumb, kochia, chickweed, corn spurry, wild	<ul style="list-style-type: none"> • No additional surfactant is required • Repeat applications may be required if a second flush of weeds germinates prior to canopy closure. <p>Ensure the crop has not advanced beyond the recommended growth stage.</p>

		<p>tomato, cleavers , wild buckwheat , shepherd's purse , cow cockle , night-flowering catchfly , smartweed , storksbill , flixweed , narrow- leaved hawk's beard , round-leaved mallow††</p> <p>Perennials (suppression)</p> <p>Canada thistle, Perennial sow thistle, Dandelion</p> <p>Perennials (season long control)</p> <p>Quackgrass , foxtail barley Canada thistle, Perennial sow thistle</p>	<p>† Use the 0.94 L/ha rate for control of these weeds at all crop growth stages. The lower rate can be used for control of shepherd's purse, cow cockle and night-flowering catchfly at the 1-3 leaf stage of the crop, or for control of smartweed at the 4-6 leaf stage.</p> <p>†† A single application at the 0.94 L/ha rate is required</p> <p>††† Sequential applications at the 0.94 L/ha rate are required.</p> <p>†††† Sequential applications at the 0.94 L/ha rate are required or a single application of 1.4 L/ha.</p> <p>For sequential applications, ensure the crop has not advanced beyond the recommended growth stage. Maximum 1.88 L/ha is allowed for the postemergence use.</p>
--	--	--	---

Maverick III Herbicide plus Lontrel™ 360 Herbicide Tank Mixture

For hard-to-control weeds (see list below) in glyphosate tolerant canola apply a tank mixture of 0.28 L/ha of Lontrel 360 with 0.94 L/ha of Maverick III Herbicide in 100L of water per hectare. Apply when canola is in the 2 - 6 leaf stage. Refer to the Lontrel 360 and the Maverick III Herbicide labels for lists of other weeds controlled, timing of application, water volumes and use precautions. **Apply this tank-mixture in glyphosate tolerant canola only.**

Weeds Controlled

Canada thistle (season-long top growth)
dandelions <15cm diameter (season-long top growth)
dandelions >15cm diameter (suppression)
perennial sowthistle (season-long top growth)
wild buckwheat

WEED CONTROL IN GLYPHOSATE TOLERANT SOYBEAN

WARNING: APPLY MAVERICK III HERBICIDE ON GLYPHOSATE TOLERANT SOYBEAN VARIETIES ONLY.

NOTE: ALWAYS USE PEDIGREED (CERTIFIED) SOYBEAN SEED DESIGNATED AS GLYPHOSATE TOLERANT. SOYBEANS WHICH ARE NOT DESIGNATED AS GLYPHOSATE TOLERANT WILL BE DAMAGED OR DESTROYED BY THIS TREATMENT.

DO NOT APPLY BY AIR

WEED CONTROL IN GLYPHOSATE TOLERANT SOYBEAN

Rate (L/ha)	Growth Stage of Crop	Weeds Controlled	Comments (use 100-200 L/ha water volumes)
1.88	First trifoliate leaf stage through to flowering.	velvetleaf, common ragweed, common lambsquarters, redroot pigweed, smooth pigweed, cocklebur, green smartweed, ladythumb, Pennsylvania smartweed, eastern black flowering nightshade, wild mustard, wild buckwheat, foxtail (green, yellow, giant), barnyard grass, crabgrass (smooth, large), quackgrass, milkweed , yellow nutsedge† , fall panicum, wild proso millet	<ul style="list-style-type: none"> • A second application may be used for late weed flushes emerging after the initial treatment † suppression only. • This second application must be made no later than the flowering stage of the soybean.
1.88-3.75	First trifoliate leaf stage through to flowering.	Perennial sow thistle, Canada thistle, wire- stemmed muhly	<ul style="list-style-type: none"> • A single application at the higher rate or a second (sequential) application of 1.88 L/ha will improve control in heavy weed infestations. • If sequential applications of 1.88 L/ha are used they should be at least 2 weeks apart for best results on perennial weeds. • This second application must be made no later than the flowering stage of the soybean. • Perennial sow thistle and Canada thistle should be from the rosette stage to 50 cm in height and actively growing. • Wire -stemmed muhly should be 10-20 cm in height and actively growing. • Plants not fully emerged at the time of application will escape the treatment.
3.75	First trifoliate leaf stage through to flowering.	All weeds listed above, plus milkweed† , yellow nutsedge† , field bindweed†	<ul style="list-style-type: none"> • Only one application season at 3.75 L/ha. †† Will also be controlled

			<p>by sequential applications of 1.88 L/ha. Applications should be at least 2 weeks apart for optimum control.</p> <ul style="list-style-type: none">• This second application must be made no later than the flowering stage of the soybean.• Milkweed should be 15 - 60 cm in height and actively growing; nutsedge should be 5-15 cm in height and actively growing. Plants not fully emerged at the time of application will not be controlled
--	--	--	---

† Weeds will be more easily controlled and early crop competition avoided with applications made when the weeds are small. Control of weeds greater than 25 cm in height will be inconsistent, although some weeds may be controlled.

Maverick III Herbicide plus Pursuit Herbicide Tank Mixture

For added residual control of late germinating eastern black nightshade, common lamb's quarters, redroot pigweed, velvetleaf, fall panicum and wild proso millet, Pursuit herbicide may be tank mixed with Maverick III Herbicide at a rate of 1.88 litres per hectare. Use 0.16 to 0.21 litres per hectare of Pursuit Herbicide and apply up to and including the 3rd trifoliolate leaf stage of the glyphosate tolerant soybeans in 100-200 litres per hectare of clean water. The higher rate is recommended for heavier infestations. This tank mix is recommended primarily for soybean systems with row spacings of 50 centimetres (20 inches) or more where a single application timing is desired.

Mixing: Add and mix Pursuit Herbicide as per instructions on the Pursuit Herbicide label and then add Maverick III Herbicide as per instructions on this label.

A PHI of 100 days is required for the tank mix of Maverick III Herbicide and Pursuit herbicide on glyphosate tolerant soybeans.

Only one application per season of Maverick III Herbicide at 1.88 litres per hectare tank mixed with Pursuit herbicide at 0.16 to 0.21 litres per hectare is permitted.

Refer to the Pursuit herbicide label for further safety precautions and handling instructions.

Maverick III Herbicide plus Assure II Tank Mixture

For control of volunteer glyphosate tolerant corn, Assure II Herbicide may be tank mixed with Maverick III Herbicide. Use 1.88 – 3.75 litres per hectare Maverick III Herbicide and 0.38 litres per hectare of Assure II Herbicide.

Apply in 100 – 300 litres per hectare of clean water.

Mixing: Add and mix Assure II Herbicide as per instructions on the Assure II Herbicide label and then add Maverick III Herbicide as per instructions on this label.

This tank mix is to be applied when the crop is from the first trifoliolate leaf stage through flowering and when the volunteer glyphosate tolerant corn is at the 2-6 leaf stage.

A PHI (preharvest interval) of 80 days is required for the tank mix of Maverick III Herbicide and Assure II Herbicide on glyphosate tolerant soybeans.

Refer to the Assure II Herbicide label for further safety precautions and handling

WEED CONTROL IN GLYPHOSATE TOLERANT CORN

WARNING: APPLY MAVERICK III HERBICIDE ON GLYPHOSATE TOLERANT CORN VARIETIES ONLY.

NOTE: ALWAYS USE PEDIGREED (CERTIFIED) CORN SEED DESIGNATED AS GLYPHOSATE TOLERANT. CORN WHICH IS NOT DESIGNATED AS GLYPHOSATE TOLERANT WILL BE DAMAGED OR DESTROYED BY THIS TREATMENT.

DO NOT APPLY BY AIR.

Rate (L/ha)	Growth Stage of Crop	Weeds Controlled†	Comments (use 100-200 L/ha water volumes)
1.88	Up to and including 8 leaf stage.	<p>Velvetleaf, common ragweed, common lambsquarters, redroot pigweed, smooth pigweed, cocklebur, green smartweed, ladysthumb, Pennsylvania smartweed, eastern black flowering nightshade, wild mustard, wild buckwheat, foxtail (green, yellow, giant), barnyard grass, crabgrass (smooth, large), quackgrass, fall panicum, wild proso millet</p> <p>Wild oats, volunteer barley, volunteer wheat, stinkweed, wild mustard, Russian thistle, nonglyphosate tolerant canola (rapeseed), hempnettle, kochia, chickweed, corn spurry, wild tomato, cleavers, shepherd's purse, cow cockle, nightflowering catchfly, stork's-bill, flixweed, narrow-leaved hawk's beard</p>	<p>A second application may be used for late weed flushes emerging after the initial treatment.</p> <p>This second application must be made no later than the 8 leaf stage of the corn.</p>

1.88		Common milkweed, yellow nutsedge, roundleaved mallow, field bindweed	For control of common milkweed, yellow nutsedge, roundleaved mallow and field bindweed use two applications of 1.88 L/ha. This second application must be made no later than the 8 leaf stage of the corn. Milkweed should be 15-60 cm in height and actively growing. Yellow nutsedge should be 5-15 cm in height and actively growing.
1.88		Perennial sow thistle, Canada thistle, wire-stemmed muhly	A second (sequential) application of 1.88 L/ha will improve control in heavy weed infestations. If sequential applications are used they should be at least 2 weeks apart for best results on perennial weeds. This second application must be made no later than the 8 leaf stage of the corn. Perennial sow thistle and Canada thistle should be from the rosette stage to 50 cm in height and actively growing. Wire-stemmed muhly should be 10-20 cm in height and actively growing. Plants not fully emerged at the time of application will escape treatment.
1.88 Maverick III Herbicide + 0.75–1.0 kg ai/ha atrazine†	Up to and including 5th leaf stage.	Residual control of lamb's-quarters, redroot pigweed, common ragweed	Tank mix should be used when only a single application timing is desired. Use higher rate of atrazine for heavier weed infestations.
1.88 Maverick III Herbicide + 2.5-3.7 Marksman Herbicide	Up to and including 5th leaf stage.	Residual control of lamb's-quarters, redroot pigweed, common ragweed, velvetleaf	Tank mix should be used when only a single application timing is desired. Use higher rate Marksman Herbicide for heavier weed infestations.

† 0.75-1.0 kg ai atrazine/ha is equivalent to 1.56-2.08 L/ha of Aatrex Liquid 480

†Weeds will be more easily controlled and early crop competition avoided with applications made when the weeds are small. Control of weeds greater than 25 cm in height will be inconsistent, although some weeds may be controlled.

PERENNIAL WEED CONTROL

ALWAYS READ PRECAUTIONARY STATEMENTS, GENERAL INFORMATION and MIXING and APPLICATION SECTIONS PRIOR TO SPECIFIC APPLICATION INFORMATION IN ANY LABEL SECTION.

When applied as recommended under the conditions described, this product will control the perennial weeds listed in the following table:

PERENNIAL WEED CONTROL WITH MAVERICK III HERBICIDE

WEED	APPLICATION			COMMENTS
	GROWTH STAGE	RATE (L/ha)	WATER VOLUME (L/ha)	
Quackgrass (control, light to moderate infestations)	3 to 4 green leaves or more	1.88	50 - 300	.Apply in clean water using flat fan nozzles. .Allow 3 or more days after treatment before tillage. .Refer to " Quackgrass " notes for more information. .For higher water volumes (i.e. 150 - 300 L/ha) an approved surfactant must be added at 0.5 litres per 100 litres of clean water (0.5% v/v). Refer to list of surfactants. See also below.
Quackgrass (long term control, heavy infestations, high water volumes)	3 to 4 green leaves or more	1.88– 5.25	50 - 300	.Allow 3 or more days after treatment before tillage. .Rates higher than 1.88 L/ha will provide more consistent, longer term control, especially with heavier infestations and/or higher water volumes (i.e. 150-300 L/ha) .Refer to " Quackgrass " notes for more information.
Canada Thistle	rosette stage (summer-fallow)	1.88	50 - 100	.Apply in clean water using flat fan nozzles. .Allow 10 or more days after treatment before tillage. .Refer to notes in "Canada Thistle" section for more information.
Canada Thistle	bud stage or beyond	3.56– 5.25	100 - 300	.Allow 5 or more days after treatment before tillage.
Field Bindweed	full bloom or beyond	5.25– 9	100 - 300	.Allow 7 or more days after treatment before tillage.

Common Milkweed	bud to full bloom (preharvest)	1.88	50 - 100	.See preharvest application section .Allow 7 or more days after treatment before tillage. .Reduced control may occur after full bloom.
	bud to full bloom	9	100 - 300	.Milkweed may not all be in the correct stage, therefore, repeat treatments may be required.
Toadflax	Vegetative Stage (summerfallow)	1.88	50-100	.Apply in clean water using flat fan nozzles .Allow 7 or more days after treatment before tillage in summerfallow .For more information, see summerfallow control, or preharvest control
	Bud to Full Bloom (preharvest)			
Alfalfa	Early bud to full bloom stage. Fall applications only	2.8– 3.75	50 - 300	.Allow 5 or more days after treatment before tillage. Use the higher rates when alfalfa populations are high or when heavy grass infestations are also present. .For spring applications and control in minimum tillage systems using a 2,4-D tank-mix
Dandelion	< 15 cm	1.88	50 - 100	.Allow 3 or more days after treatment before tillage for all rates. .Use the higher rate when infestations are heavy. .Refer to notes in Dandelion Section for more information. .Allow 7 or more days after treatment before tillage. For more information, see preharvest control section.
	> 15 cm	2.78– 3.75	50 - 300	
	Rosette to full bloom (preharvest)	1.88	50 - 100	
Foxtail barley	Seedling to heading	1.88– 3.75	50 -100	-Allow a minimum of 1 day after treatment before tillage or seeding. -Use higher rates for larger, more established plants, heavy infestations or if plants are stressed
Other Perennials (see perennial weeds listing)	early heading or early bud stage	5.25– 9	100 - 300	.Allow 7 or more days after application before tillage.

†NOTE: For spot treatment, mix 90 mL of product in 5L clean water per 100 m². (1.88 9 L/ha is approximately equivalent to 19– 90 L/100 m², respectively).

SPECIAL NOTES FOR PERENNIAL WEED CONTROL

QUACKGRASS

For **season-long control on fall tilled ground**: Apply 1.88L/ha of this product in spring prior to seeding. Apply in 50 to 100 L/ha of clean water as described in the preceding table. Delay application until the majority of quackgrass plants have 4-5 green leaves. This stage usually occurs 1 to 4 weeks later on fall tilled ground than on undisturbed ground. Reduced control may result on ground tilled deeper than 15 cm.

NOTE:

This treatment will provide season-long control of quackgrass on fall tilled ground. Reduced control will be experienced versus this product on non-fall tilled ground. Repeat treatments may be necessary.

Applications on forages should be followed by tillage 3 days or later and should be made when good growing conditions exist.

If a frost has occurred, wait several days to determine if the quackgrass has recovered. Quackgrass can be treated after a mild frost provided there are 3 to 4 green leaves actively growing at the time of application. Do not apply after the first damaging frost in the fall.

Surfactant Information:

The following is a list of approved surfactants for use with Maverick III Herbicide for control of quackgrass:

Agral 90	Companion
Ag Surf	Frigate

Always refer to surfactant label for specific instructions regarding use of that product.

CANADA THISTLE

Control of Canada thistle at the rosette stage: To ensure the proper timing of application the following steps must be followed:

1. Conduct summerfallow tillage as usual and perform the last tillage operation between July 15th and August 1.
2. Allow the thistles to regrow for a minimum of 5 weeks until they are a minimum of 15cm in diameter and in the rosette stage of growth.

NOTE: Canada thistle can be treated after a mild frost provided the leaves are still green and actively growing at the time of application. Do not apply after the first damaging frost in the fall.

Maverick III Herbicide plus Banvel II Tank Mixtures

For control of Canada thistle (and perennial sow thistle) in summerfallow or in post-harvest stubble, apply 1.28 L/ha Maverick III Herbicide plus 1.25 L/ha Banvel II in 100 – 200 L/ha of clean water. In addition, add 350 mL/ha of a non-ionic surfactant registered for use with this product, such as Agral 90, Ag Surf, or Companion.

For best results in summerfallow, cultivate in the spring and apply when the majority of thistles are 15 cm to 25 cm tall and before the bud stage. Cultivate 3 weeks after application.

In post harvest stubble, apply this tank mixture to actively growing thistles at least 2 weeks prior to a killing frost.

NOTE:

Grow only cereals, canola (including rapeseed), soybeans, field corn, sweet corn, or white beans after application of this tank mixture.

If application is made after September 1st, or if soil moisture levels are extremely low after application, crop injury may occur in the spring following application.

TOADFLAX

Control of Toadflax in a Summerfallow Vegetative Stage

To ensure the proper timing of application, the following steps must be followed:

1. Conduct summerfallow tillage as usual and perform the last tillage operation between July 10-21.
2. Allow toadflax to regrow for a minimum of 4-5 weeks until they are minimum of 15 cm tall and at a lush green vegetative stage.

NOTE: Toadflax can be treated after a mild frost provided the leaves are still green and actively growing at the time of application. Do not apply after the first damaging frost.

DANDELION

Applications should be made up to and including bloom for best results. Follow-up control measures should be used to manage new dandelions germinating from seed to maintain control throughout the season.

ALFALFA CONTROL WITH 2,4-D TANK-MIX:

The addition of 2,4-D may improve alfalfa control in situations where control may be more difficult to obtain, such as in minimum tillage systems where populations are heavy, and with spring applications.

For fall control of established stands of alfalfa, apply 1.88 – 3.75 L/ha Maverick III Herbicide – and 1.2 – 2.4 L/ha of any 500 g/L 2,4-D amine or low volatile ester formulation in 100 – 200 L water/ha. (Adjust product rates accordingly for other 2,4-D formulations).

For spring applications, use only the low rate of 2,4-D (i.e. 1.2 L/ha) and 1.88 – 3.75 L/ha Maverick III Herbicide. Only cereal crops not underseeded to legumes may be planted following spring applications of this tank-mix, and a 14 day interval between application and planting is required.

Use the higher Maverick III Herbicide rates when perennial grasses are prevalent.

ALL PERENNIAL WEEDS

Weed Stages: Weeds must be at the proper stage for effective control. Refer to "**Perennial Weed Control with Maverick III Herbicide** "

Nozzle Type: For best results with conventional boom equipment apply this product with 50 to 300 L/ha of clean water using flat fan nozzles and no more pressure than 275 kPa.

Rhizome Dormancy: Reduced control may result if rhizomes have become dormant. Dormancy may occur if soil fertility is low and/or the land has not been tilled for several years.

Mowing Effects: Mowing prior to application will reduce effectiveness unless weeds are allowed to regrow to the proper stage before application.

Tillage Effects: Fall or spring tillage prior to spring applications and tillage between harvesting and fall applications will reduce the effectiveness on perennial weeds. Follow-up tillage after application should be delayed 5-7 days for best results (see Weed Control Table for specific tillage interval for each weed).

Rainfall Effects: Heavy rainfall immediately after application may wash the chemical off the foliage and a repeat treatment may be required. Do not apply if rainfall is forecast for the time of application.

Regrowth from Germinating Seeds: This product only controls emerged plants. Repeat treatments or other weed control measures may be required to control weeds regenerating from seeds or other underground parts.

Frost Effects: Heavy frosts prior to application may reduce control. Do not apply after the first damaging frost in the fall.

CROPLAND SITUATIONS

ALWAYS READ PRECAUTIONARY STATEMENTS, GENERAL INFORMATION and MIXING and APPLICATION SECTIONS PRIOR TO SPECIFIC APPLICATION INFORMATION IN ANY LABEL SECTION.

This product can be applied as a broadcast spray or spot treatment prior to planting all crops, post harvest to annual crops, preharvest in wheat, barley, oats, canola (rapeseed), flax (including low linolenic acid varieties), lentils, peas, soybeans, dry beans and forages, and in summerfallow. It may also be applied as a broadcast spray in glyphosate tolerant canola, soybean or corn (refer to sections on Weed Control in Glyphosate Tolerant Canola, Soybean or Corn). It can also be applied as a directed spray in orchards, vineyards, blueberries and strawberry, and using selective equipment in soy and dry beans, orchards, vineyards, cranberries and strawberry (refer to specific sections below for more information). **For specific instructions on weed control in the following cropping situations, always refer to the Annual and Perennial Weed Control sections for more information.**

Prior to Planting - All Crops

This product may be applied prior to planting all crops for control of emerged weeds listed on this label. Ensure weeds are at the desired stage at the time of application. This product does not provide pre-emergent weed control and newly germinating weeds may be a problem in the crop. **APPLY BEFORE SEEDING OR TRANSPLANTING.**

Post Harvest Stubble Treatment

This product may be applied in the fall as a postharvest stubble treatment for control of perennial weeds such as quackgrass and Canada thistle. Allow weeds to regrow to the desired stage (20-25 cm tall for quackgrass and Canada thistle) before application and ensure they have a high proportion of green coloration. Straw should be removed or evenly spread to allow for proper regrowth and spray coverage. Heavy frosts prior to application may decrease control.

Spot Treatment (In-Crop)

This product can be applied as an in-crop spot treatment in barley, corn, oats, soybeans, wheat, strawberry, blueberry, forage grasses and legumes including seed production. Applications should be made using the same rates and at the same growth stages as listed in the weed control tables or use a 0.75% solution for annual weeds and quackgrass and a 1.5% solution for other perennial weeds (a 0.75% solution equals 0.75 litre Maverick III Herbicide in 100 litres of spray solution). The 0.75 or 1.5 per cent solutions should be applied to wet, but not run-off. Applications can be made using a boom sprayer, hose and handgun, or hand sprayer in accordance with instructions in the "Application Equipment" section.

Grazing Restrictions

Applications can be made up to heading of small grains, initial pod set on soy and dry beans, silking of corn and emergence of seed heads. The crop in the treated area will be killed. Take care to avoid drift for the same reason. **DO NOT APPLY IF CROP GROWTH HAS ADVANCED BEYOND SEED SET. ALLOW 3 TO 5 DAYS FOR MAVERICK III HERBICIDE TO TRANSLOCATE INTO ALL PLANT PARTS BEFORE GRAZING OR HARVESTING TREATED AREAS IN FORAGES.**

Summerfallow Treatment

This product, or labelled tank mixtures, may be applied in summerfallow to control weeds listed on this label. Ensure weeds are at the desired growth stage and actively growing at application for best results. Reduced control may result if weeds are drought stressed. Weeds will continue to germinate from seed throughout the growing season. Repeat treatments may be necessary to control later germinating weeds.

Minimum and Zero Tillage Cropping Systems (All Field Crops, including cereals, oilseeds, pulses, forages, corn and potatoes)

This product may be applied prior to seeding or after seeding, but before crop emergence for control of emerged weeds in minimum and zero tillage cropping systems for all field crops. Applications made too far in advance of seeding may allow weeds to emerge between application and crop emergence, as this product does not provide residual weed control.

Minimum and Zero Tillage Tank Mixtures

Maverick III Herbicide plus bromoxynil (Pardner) can be applied prior to seeding or after seeding, but before crop emergence in **wheat, barley and oats**. Refer to **Annual Weed Control with Maverick III Herbicide Tank Mixtures** table for information.

Maverick III Herbicide plus Pursuit can be applied prior to, or after, seeding, but before crop emergence in soybeans. Maverick III Herbicide will control emerged weeds listed on this label when applied as directed (refer to Annual and Perennial Weed Control sections). Pursuit will control weeds germinating from seed. Add the recommended rates of both products in 100 L of water/ha, following the instructions on the Pursuit herbicide label.

ALWAYS REFER TO THE PURSUIT LABEL FOR FURTHER INFORMATION ON WEEDS CONTROLLED, APPLICATION DIRECTIONS, AND USE PRECAUTIONS. ONLY SOYBEANS, FIELD CORN, SPRING BARLEY, SPRING WHEAT AND WINTER WHEAT MAY BE PLANTED THE SEASON FOLLOWING A PURSUIT APPLICATION. WINTER WHEAT MAY BE PLANTED THE SAME YEAR AS A PURSUIT APPLICATION TO SOYBEANS, BUT NOT EARLIER THAN 120 DAYS AFTER THE APPLICATION.

DO NOT APPLY AFTER CROP EMERGENCE

Maverick III Herbicide plus MCPA can be applied prior to seeding in wheat, barley, rye, oats, corn (field and sweet; MCPA amine only), flax and field peas (MCPA amine only). Refer to **Annual Weed Control with Maverick III Herbicide Tank Mixtures** table for information.

Maverick III Herbicide plus Buctril M can be applied prior to seeding in **wheat, rye, corn, barley, oats, flax, canary seed and seedling grasses (including brome grass, crested wheatgrass, intermediate wheat grass, slender wheatgrass, tall wheatgrass, Russian wild rye, Timothy, Orchard grass, creeping red fescue, meadow fescue, meadow foxtail, seedling tall fescue, seedling meadow brome grass, seedling streambank wheatgrass and reed canary grass** Refer to **Annual Weed Control with Maverick III Herbicide Tank Mixtures** table for information.

Maverick III Herbicide plus MCPA amine can be applied prior to seeding in **lentil and chickpea** Refer to **Annual Weed Control with Maverick III Herbicide Tank Mixtures** table for information.

Forage Legumes and Grasses

This product may be applied for control of emerged weeds prior to emergence of forage legumes and grasses. If the forages are to be under-seeded with a cover crop, this product must be applied prior to planting the cover crop.

Pasture Renovation

Use this product to control or suppress existing vegetation for zero-tillage seeding of legumes into established sod for pasture renovation. Delay spraying until weed growth is at least 20 cm in height and a maximum number of seedlings or shoots have emerged. Application can be made immediately before, during or after seeding, but before crop emergence.

Forage Seed Production

For spot treatment control of perennial weed problems such as quackgrass and Canada thistle in seed fields, apply as directed to vegetation that is at least 20 to 25 cm in height but before emergence of seed head. The crop in the treated areas will be killed. Take care to avoid drift outside target area for the same reason.

PRE-HARVEST CONTROL OF QUACKGRASS, CANADA THISTLE, MILKWEED, TOADFLAX and DANDELION; SEASON-LONG CONTROL OF PERENNIAL SOW THISTLE, AND HARVEST MANAGEMENT

For control of quackgrass, Canada thistle, common milkweed, toadflax and dandelion; and season-long control of perennial sow thistle, Maverick III Herbicide can be applied prior to harvest of wheat, barley (including malting barley), oats, canola (rapeseed), flax (including low linolenic acid varieties), lentils, peas, dry beans, soybeans and forages. DO NOT apply to crops if grown for seed production.

This treatment may also provide harvest management benefits, by drying down crop and weed vegetative growth, for example, where late flushes of annual weeds, green vegetative crop growth, or late tillering may interfere with harvest operations. EXTREMELY COOL, WET AND/OR CLOUDY WEATHER CONDITIONS BETWEEN THE TIME OF APPLICATION AND THE ANTICIPATED HARVEST DATE MAY SLOW DOWN ACTIVITY OF THIS PRODUCT, THEREBY DELAYING CROP DRYDOWN AND HARVEST DATE.

Maverick III Herbicide should be applied pre-harvest at 1.88 L/ha in 50 to 100 L/ha of clean water, by ground application only. Apply only when the crop has 30% or less grain moisture content. This stage typically occurs 7 to 14 days before harvest. For forage crops, apply this product at 1.88-3.75 L/ha 3-7 days prior to the last cut before rotation or forage renovation. Consult the table "Guidelines for Timing of Preharvest Applications" for visual indicators of this stage in each crop. For the best weed control results quackgrass should be actively growing and have at least 4 to 5 green leaves. Canada thistle and perennial sow thistle should be actively growing and at or beyond the bud stage for best results. Common milkweed should be at the bud to bloom stage and actively growing for best results. Applications for weed control (not for harvest management) must be made at the correct stage of both weed and crop growth.

Apply only during the period 7-14 days (or 3-7 days for forage applications) before harvest to ensure best weed control and to maximize harvest management benefits. Earlier application may reduce crop yield and/or quality, and may lead to excess glyphosate residues in the crop.

Do not expose or contaminate any body of water or non-target vegetation by direct application, spray drift, or when cleaning and rinsing spray equipment.

DO NOT APPLY BY AIRCRAFT

GUIDELINES FOR TIMING OF PREHARVEST APPLICATIONS

CROP(S)	PERCENT GRAIN MOISTURE	VISUAL SYMPTOMS
WHEAT/BARLEY/ OATS	Less than 30	Hard dough stage; a thumbnail impression remains on seed.
CANOLA	Less than 30	Pods are green to yellow; most seeds are yellow to brown.
FLAX (INCLUDING LOW LINOLENIC ACID VARIETIES)	Less than 30	Majority (75%-80%) of bolls are brown.
PEAS	Less than 30	Majority (75%-80%) of pods are brown.
LENTILS	Less than 30	Lowermost pods (bottom 15%) are brown and seeds rattle.
DRY BEANS	Less than 30	Stems are green to brown in colour; pods are mature (yellow to brown in colour); 80%-90% leaf drop (original leaves).

SOYBEANS	Less than 30	Stems are green to brown in colour; pod tissue is dry and brown in appearance; 80%-90% leaf drop.
FORAGES	Not applicable	Normal stage for forage harvesting.

Tree Plantings

Shelterbelts and Nursery Stock (Woody Ornamentals)

This product may be used to control listed annual or perennial weeds prior to planting, or as a post directed spray in established nurseries or shelterbelts of the following species:

Deciduous

Ash - Fraxinus spp.
Caragana - Caragan spp.
Cherry - Prunus spp.
Elm - Ulmus spp.
Lilac - Syringa spp.
Maple - Acer spp.
Mountain Ash - Sorbus spp.
Poplar - Populus spp.
Russian Olive - Elaeagnus spp.
Willow - Salix spp.

Coniferous

Fir – Abies spp.
Juniper - Juniperus spp.
Pine - Pinus spp.
Spruce - Picea spp.
Yew - Taxus spp.

NOTE: This product is not recommended for use as an over-the-top broadcast spray in forest tree nurseries or in Christmas tree plantations. Application in such sites should be limited to directed sprays. DO NOT treat Christmas tree plantations in the year of anticipated harvest.

TREE, VINE and BERRY CROPS

This product is recommended for annual and perennial weed control in established vineyards or orchards, in blueberry, cranberry and strawberry, or for site preparation prior to transplanting tree and vine crops. Applications may be made with boom equipment, shielded sprayers, hand-held and high volume orchards guns, or with wiper applicator equipment (orchards, vineyards, cranberry and strawberry only). See the "Mixing and Application Equipment Information" section of this label and the following table for specific information on the use of equipment.

Repeat treatments may be necessary to control weeds originating from underground parts of untreated weeds or from seeds. This product does not provide residual or pre-emergent weed control. For subsequent weed control, follow a program using residual herbicides or use repeated applications of this product. Do not apply more than 26 litres of this product per hectare per year.

EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE SOLUTION, SPRAY, DRIFT, OR MIST WITH FOLIAGE OR GREEN BARK OF TRUNK, BRANCHES, SUCKERS, FRUIT, CANES OF BLUEBERRY BUSHES, OR OTHER PARTS OF TREES OR VINES. CONTACT OF THIS PRODUCT WITH OTHER THAN MATURED BROWN BARK CAN RESULT IN SERIOUS CROP DAMAGE.

Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed or cut and have not been allowed to regrow to the recommended stage for treatment.