



****CONTAINER****

GROUP	3	FUNGICIDE
-------	----------	-----------

JAU6476 100FS

SEED TREATMENT FUNGICIDE SUSPENSION

A seed treatment product for control or suppression of certain seed-borne and soil-borne diseases of Cereals, Chickpea, Corn, Dry Beans, Dry Peas (including field peas), Lentil and Soybean.

COMMERCIAL

ACTIVE INGREDIENT: prothioconazole..... 100 g/L

Contains 1,2-benzisothiazolin-3-one at 0.027%, and 5-chloro-2-methyl-4-isothiazolin-3-one at 0.0009%, and 2-methyl-4-isothiazolin-3-one at 0.0003% as preservatives.

REGISTRATION NO. 30101 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND BOOKLET BEFORE USING

POTENTIAL SKIN SENSITIZER

Bayer CropScience Inc.
Suite200, Quarry Park Blvd. SE
Calgary, Alberta
T2C 3G3

NET CONTENTS: 0.5 – 1000 L

PROTECT FROM FREEZING

For product information, call: 1-888-283-6847

In case of spills, poisoning, fire or other emergencies,
call: 1-800-334-7577 (24 hours a day)

GENERAL INFORMATION

JAU6476 100FS Seed Treatment Fungicide is a concentrate flowable seed treatment formulation containing the systemic, broad-spectrum fungicide prothioconazole. JAU6476 100FS Seed Treatment Fungicide provides protection from listed seed, seedling and soil-borne diseases of cereals, chickpea, corn, dry beans, dry peas (including field peas), lentil and soybean.

JAU6476 100FS Seed Treatment Fungicide may be applied in commercial or on-farm conventional treating equipment which can accurately control application rates and provide good distribution of the chemical onto the seed in the mixing chamber.

PRECAUTIONS:

- KEEP OUT OF REACH OF CHILDREN.
- Harmful if swallowed, inhaled, or absorbed through skin. Potential skin sensitizer. Avoid contact with skin or clothing. Prolonged or frequent repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.
- Workers must wear long pants, a long-sleeved shirt and chemical-resistant gloves during mixing, loading, treating, clean-up, maintenance of seed treatment equipment, bagging, sewing or stacking of bagged treated seed. In addition, workers must wear a suitable dust mask when bagging or sewing bags of treated seed or when transferring seed to a storage bin.
- Do not apply in a way that this product will contact workers or other persons. Only handlers (mixers, loaders and applicators) wearing personal protective equipment may be in the area being treated during application.
- Wear long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seed.
- All bags containing treated seed for sale or use in Canada must be labeled or tagged as follows: "This seed has been treated with JAU6476 100FS Seed Treatment Fungicide containing prothioconazole. Use chemical-resistant gloves when handling treated seed. DO NOT use for feed, food or oil processing. Store away from feeds and other foodstuffs. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil."
- If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact Bayer CropScience Canada Inc. at 1-888-283-6847 or www.cropscience.bayer.ca.

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

There is no specific antidote if this product is ingested. Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS:

- TOXIC to aquatic organisms.

STORAGE:

Store product in original container only, away from other pesticides, fertilizer, food or feed. Store in a cool, dry place and avoid excessive heat. Keep container closed. To prevent contamination, store this product away from food or feed.

DISPOSAL:

FOR NON-RETURNABLE CONTAINERS:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.

FOR RECYCLABLE CONTAINERS:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the mixture in the tank.
2. Make the empty container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

FOR RETURNABLE CONTAINERS:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

FOR REFILLABLE CONTAINERS:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not use this container for any other purpose.

TRIPLE RINSE INSTRUCTIONS (FOR RECYCLABLE AND DISPOSABLE CONTAINERS ONLY): This container should be triple-rinsed prior to its disposal. The rinsate can be added to the seed treatment. Follow these steps for rinsing of containers:

1. Transfer JAU6476 100FS Seed Treatment Fungicide from the container to the seed treating equipment applicator tank.
2. Add 1/3 of the rinse water to the container and agitate the contents.
3. Transfer the rinsate to the applicator tank.
4. Rinse as above 2 more times.
5. Thoroughly mix the JAU6476 100FS Seed Treatment Fungicide and rinse water in the applicator tank prior to use.

CAUTION: Be sure to adjust the undiluted application rate by the same volume of water added to compensate for the dilution of the product caused by the addition of rinsate.

DISPOSAL OF UNUSED, UNWANTED PRODUCT:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

DISPOSAL OF TREATED SEED:

Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. Dispose of seed packaging in accordance with local requirements. **DO NOT** re-use bags from treated seed to handle food or feed products.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.



****BOOKLET****

GROUP	3	FUNGICIDE
-------	----------	-----------

JAU6476 100FS

SEED TREATMENT FUNGICIDE

SUSPENSION

A seed treatment product for control or suppression of certain seed-borne and soil-borne diseases of Cereals, Chickpea, Corn, Dry Beans, Dry Peas (including field peas), Lentil and Soybean.

COMMERCIAL

ACTIVE INGREDIENT: prothioconazole..... 100 g/L

Contains 1,2-benzisothiazolin-3-one at 0.027%, and 5-chloro-2-methyl-4-isothiazolin-3-one at 0.0009%, and 2-methyl-4-isothiazolin-3-one at 0.0003% as preservatives.

REGISTRATION NO. 30101 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND BOOKLET BEFORE USING

POTENTIAL SKIN SENSITIZER

Bayer CropScience Inc.
Suite 200, 160 Quarry Park Blvd. SE
Calgary, Alberta
T2C 3G3

NET CONTENTS: 0.5 – 1000 L

PROTECT FROM FREEZING

For product information, call: 1-888-283-6847

In case of spills, poisoning, fire or other emergencies,
call: 1-800-334-7577 (24 hours a day)

Table of Contents:	Section	Page Number
GENERAL INFORMATION	1. The Product	3
SAFETY AND HANDLING	2. Precautions, Protective Clothing and Equipment ..	3
	3. First Aid and Toxicological Information	4
	4. Environmental Precautions	4
	5. Storage	4
	6. Disposal	5
	7. Notices	6
DIRECTIONS FOR USE	8. Pests Controlled	6
	9. Recommended Uses	7
	10. Application Cautions	9
	11. Use Limitations	10
	12. Resistance Management Recommendations	10

GENERAL INFORMATION

Section 1: The Product

JAU6476 100FS Seed Treatment Fungicide is a concentrate flowable seed treatment formulation containing the systemic, broad-spectrum fungicide prothioconazole. **JAU6476 100FS** Seed Treatment Fungicide provides protection from listed seed, seedling and soil-borne diseases of cereals, chickpea, corn, dry beans, dry peas (including field peas), lentil and soybean.

JAU6476 100FS Seed Treatment Fungicide may be applied in commercial or on-farm conventional treating equipment which can accurately control application rates and provide good distribution of the chemical onto the seed in the mixing chamber.

SAFETY AND HANDLING

Section 2: Precautions, Protective Clothing and Equipment

PRECAUTIONS:

- KEEP OUT OF REACH OF CHILDREN.
- Harmful if swallowed, inhaled, or absorbed through skin. Potential skin sensitizer. Avoid contact with skin or clothing. Prolonged or frequent repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.
- Workers must wear long pants, a long-sleeved shirt and chemical-resistant gloves during mixing, loading, treating, clean-up, maintenance of seed treatment equipment, bagging, sewing or stacking of bagged treated seed. In addition, workers must wear a suitable dust mask when bagging or sewing bags of treated seed or when transferring seed to a storage bin.
- Do not apply in a way that this product will contact workers or other persons. Only handlers (mixers, loaders and applicators) wearing personal protective equipment may be in the area being treated during application.
- Wear long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seed.
- All bags containing treated seed for sale or use in Canada must be labeled or tagged as follows: "This seed has been treated with **JAU6476 100FS** Seed Treatment Fungicide containing prothioconazole. Use chemical-resistant gloves when handling treated seed. DO NOT use for feed, food or oil processing. Store away from feeds and other foodstuffs. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil."
- If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please

contact Bayer CropScience Canada Inc. at 1-888-283-6847 or www.cropscience.bayer.ca.

Section 3: First Aid and Toxicological Information

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

There is no specific antidote if this product is ingested. Treat symptomatically.

Section 4: Environmental Precautions

TOXIC to aquatic organisms.

Section 5: Storage

Store product in original container only, away from other pesticides, fertilizer, food or feed. Store in a cool, dry place and avoid excessive heat. Keep container closed. To prevent contamination, store this product away from food or feed.

Section 6: Disposal

FOR NON-RETURNABLE CONTAINERS:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.

FOR RECYCLABLE CONTAINERS:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the mixture in the tank.
2. Make the empty container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

FOR RETURNABLE CONTAINERS:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

FOR REFILLABLE CONTAINERS:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not use this container for any other purpose.

TRIPLE RINSE INSTRUCTIONS (FOR RECYCLABLE AND DISPOSABLE CONTAINERS ONLY): This container should be triple-rinsed prior to its disposal. The rinsate can be added to the seed treatment. Follow these steps for rinsing of containers:

1. Transfer **JAU6476 100FS** Seed Treatment Fungicide from the container to the seed treating equipment applicator tank.
2. Add 1/3 of the rinse water to the container and agitate the contents.
3. Transfer the rinsate to the applicator tank.
4. Rinse as above 2 more times.
5. Thoroughly mix the **JAU6476 100FS** Seed Treatment Fungicide and rinse water in the applicator tank prior to use.

CAUTION: Be sure to adjust the undiluted application rate by the same volume of water added to compensate for the dilution of the product caused by the addition of rinsate.

DISPOSAL OF UNUSED, UNWANTED PRODUCT:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in

case of a spill, and for clean-up of spills.

DISPOSAL OF TREATED SEED:

Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. Dispose of seed packaging in accordance with local requirements. DO NOT re-use bags from treated seed to handle food or feed products.

Section 7: Notices

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

DIRECTIONS FOR USE

JAU6476 100FS Seed Treatment Fungicide is a seed treatment formulation for use in commercial seed treatment operations and for on-farm treatment with conventional seed treating equipment which can accurately meter, mix and apply flowable seed treatment formulations. Allow seeds to dry before bagging, storing or seeding.

This product is recommended to be diluted with water or another suitable liquid just prior to application to ensure uniform coverage to the seed during the application process. Do not store diluted mixture. Additional or continuous agitation or mixing of a diluted mixture may be necessary to prevent **JAU6476 100FS** Seed Treatment Fungicide from settling out. Uniform application to seed is necessary to ensure best disease protection and seed safety.

JAU6476 100 FS Seed Treatment Fungicide-treated seed may reduce seed flow in the seed drill. Recalibration of the seed drill may be required to obtain correct seeding rate before planting.

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Section 8: Pests Controlled

JAU6476 100 FS Seed Treatment Fungicide is a seed treatment containing the fungicide prothionconazole. **JAU6476 100 FS** Seed Treatment Fungicide provides protection from the following seed, seedling and soil-borne diseases on labeled crops.

Section 9: Recommended Uses

CROP	PEST	APPLICATION RATE	DIRECTIONS
<p>Cereal grains (wheat, barley, oat, rye, triticale, millet, pearl millet, proso millet)</p>	<p>Diseases Controlled:</p>		<p>Use only in treating equipment which can accurately control application rates and provide good distribution of the chemical onto the seed in the mixing chamber. Not for use in hopper box, slurry-box or similar seed treatment applications.</p> <p>Uniform application to seed is necessary to ensure best disease protection and seed safety.</p>
	<p>Seed rot / pre-emergence damping-off caused by seed- and soil-borne <i>Fusarium</i> spp. and <i>Cochliobolus sativus</i></p> <p>Seedling blight caused by seed-borne <i>Fusarium</i> spp. and <i>Cochliobolus sativus</i></p> <p>Post-emergence damping-off caused by soil-borne <i>Fusarium</i> spp.</p> <p>Common bunt (<i>Tilletia foetida</i>)</p> <p>False loose smut and covered smut of barley (<i>Ustilago nigra</i> and <i>U. hordei</i>);</p> <p>Barley leaf stripe (<i>Pyrenophora graminea</i>)</p> <p>Loose smut and covered smut of oat (<i>Ustilago avenae</i> and <i>U. kolleri</i>)</p>	<p>Apply 50 mL per 100 kg of seed</p>	
	<p>Diseases Suppressed:</p>		
	<p>Loose smut of wheat (<i>Ustilago tritici</i>)</p> <p>Root rot caused by seed-borne and soil-borne <i>Fusarium</i> spp.</p> <p>Root rot caused by soil-borne <i>Cochliobolus sativus</i></p> <p>Crown rot of small-grain cereals caused by seed-borne <i>Fusarium</i> spp.</p> <p>Seedling blight caused by seed-borne <i>Penicillium</i> spp.</p>	<p>Apply 50 mL per 100 kg of seed</p>	

CROP	PEST	APPLICATION RATE	DIRECTIONS
Corn (field, sweet, popcorn)	Diseases Controlled:		Use only in treating equipment which can accurately control application rates and provide good distribution of the chemical onto the seed in the mixing chamber. Not for use in hopper box, slurry-box or similar seed treatment applications. Uniform application to seed is necessary to ensure best disease protection and seed safety.
	Seed rot and pre-emergence damping-off caused by seed-borne and soil-borne <i>Fusarium</i> spp. Post-emergence damping-off caused by soil-borne <i>Fusarium</i> spp. Seed rot and pre-emergence damping-off caused by seed-borne <i>Cladosporium</i> spp.	Apply 50 mL per 100 kg of seed	
	Diseases Suppressed:		
	Seed rot and pre-emergence damping-off caused by <i>Penicillium</i> spp.	Apply 50 mL per 100 kg of seed	

CROP	PEST	APPLICATION RATE	DIRECTIONS
Dry bean, dried and field pea, chickpea, lentil and soybean, immature soybean seed	Diseases Controlled:		Use only in treating equipment which can accurately control application rates and provide good distribution of the chemical onto the seed in the mixing chamber. Not for use in hopper box, slurry-box or similar seed treatment applications. Uniform application to seed is necessary to ensure best disease protection and seed safety.
	Seed rot / pre-emergence damping-off and post-emergence damping-off caused by soil-borne <i>Fusarium</i> spp. Seed rot and pre-emergence damping-off of soybean caused by seed-borne <i>Phomopsis longicolla</i> Seed rot and pre-emergence damping-off of chickpea caused by <i>Ascochyta rabiei</i>	Apply 50 mL per 100 kg of seed	
	Diseases Suppressed:		
	Seed infection on soybean caused by <i>Penicillium</i> spp.	Apply 50 mL per 100 Kg of seed	

TANK-MIXTURES

For control of certain insect pests and/or additional diseases not on the JAU6476 100FS Seed Treatment Fungicide product label, JAU6476 100FS Seed Treatment Fungicide may be combined with certain seed treatment insecticides and fungicides.

For recommended tank-mixes, read the label directions for each product and follow the most restrictive label precautions and limitations.

CROP	TANK-MIX PARTNER
Dry Bean Pea (dried and field) Chickpea Lentil Soybean	Trilex AL
Corn (field, sweet, popcorn)	Allegiance FL (except popcorn) Poncho 600 Poncho 600 + Allegiance FL
Wheat, barley, oats & rye for export only	Allegiance FL

Section 10: Application Cautions

SEED QUALITY: Lab and field studies have shown that **JAU6476 100 FS** Seed Treatment Fungicide-treated seed can be stored without loss in germination or disease performance. Many seed types are prone to mechanical damage during seed handling, and a reduction in germination within a seed lot can occur should handling cause seed damage irrespective of seed treatment application. **JAU6476 100 FS** Seed Treatment Fungicide-treated or untreated seed may drop in germination with age and germination drops are accelerated under adverse storage conditions or in seed with a high level of mechanical damage. Use handling equipment designed to minimize mechanical damage when moving seed. Due to seed quality and seed storage conditions beyond the control of Bayer CropScience, no claims are made to guarantee the germination of carry-over seed or propagating materials for all crop seed. Treatment of highly mechanically scarred or damaged seed, or seed known to be of low vigour and poor quality, may result in reduced germination and/or reduction of seed and seedling vigour. Treat a small quantity of seed using equipment similar to that planned for treating the total seed lot. Conduct germination tests on a

small portion of seed before committing the total seed lot to a selected seed treatment.

COLOURATION OF TREATED SEED: This product contains a pigment which will colour the treated seed. However, users are responsible for ensuring that the treated seed, when dried and ready for bagging, storage or seeding, has an unnatural colour. If the pigment contained in the formulation does not colour the seed adequately, additional colourant must be added to the mixture while treating the seed. Regulations pertaining to colouration of treated seed enforced under the *Seeds Act* must be strictly adhered to when using this product.

SEED TREATMENT AND INOCULANTS: **JAU6476 100 FS** Seed Treatment Fungicide is compatible with *Rhizobium*-based inoculants. Please check with inoculant manufacturers for further details prior to use. Mixing with inoculants may increase drying time while treating, extending the processing time.

Section 11: Use Limitations

USE RESTRICTIONS:

- Treated seed must not be used for food, feed or oil processing.

LABELLING TREATED SEED:

All bags containing treated seed must be labelled or tagged as follows: “This seed has been treated with **JAU6476 100 FS** Seed Treatment Fungicide, which contains prothioconazole. Use chemical-resistant gloves when handling treated seed. **DO NOT** use for feed, food or oil processing. Store away from feeds and other foodstuffs. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil.”

Section 12: Resistance Management Recommendations

For resistance management, please note that **JAU6476 100 FS** Seed Treatment Fungicide contains a Group 3 fungicide. Any fungal population may contain individuals naturally resistant to **JAU6476 100 FS** Seed Treatment Fungicide and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

- Where possible, rotate the use of **JAU6476 100 FS** Seed Treatment Fungicide or other Group 3 fungicides with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group when such use is permitted.

- Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications.
- Monitor fungal populations for resistance development. Notify Bayer CropScience Inc. if reduced sensitivity of the pathogen to **JAU6476 100 FS** Seed Treatment Fungicide is suspected.
- If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different site of action, if available.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens. For further information and to report suspected resistance, contact Bayer CropScience Inc. at 1-888-283-6847 or at www.cropscience.bayer.ca .