



**** CARTON****

GROUP	3	4	FUNGICIDE
GROUP		4	INSECTICIDE

NipsIt™ SUITE Canola Seed Protectant

Flowable Insecticide and Fungicide Seed Protectant

SUSPENSION

NipsIt SUITE Canola Seed Protectant is a systemic insecticide and fungicide seed protection product for use on canola, carinata and rapeseed for early season control of flea beetles, seed and seedling diseases caused by *Pythium* spp., *Fusarium* spp. and *Rhizoctonia solani* and suppression of seed-borne blackleg (*Leptosphaeria maculans*).

For use in commercial seed treatment facilities with closed transfer including closed mixing, loading, calibrating, and closed treatment equipment only. No open transfer of NipsIt SUITE Canola Seed Protectant.

COMMERCIAL

GUARANTEE:

Clothianidin 279 g/L
Metalaxyl.....5.23 g/L
Metconazole.....1.04 g/L

Contains 5-chloro-2-methyl-4-isothiazolin-3-one at 0.00113% and 2-methyl-4-isothiazolin-3-one at 0.00037% as preservatives.

REGISTRATION NUMBER 31355 PEST CONTROL PRODUCTS ACT

**READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

CAUTION

<<Skull and Cross Bones in Diamond>>

POISON

NET CONTENTS: 72 L - 14400 L (bulk tanker)

This product contains blue colourant for application to canola/rapeseed, according to regulations pertaining to the "Seeds Act".

2019-1665
2019-05-17

HEALTH EMERGENCY OR SPILL (24 hr):
(800) 682-5368
TRANSPORTATION (24 hr.): CHEMTREC
(800) 424-9300 or (202) 483-7616

GROWER INFORMATION:

Valent Canada, Inc.

201-230 Hanlon Creek Blvd., Guelph, Ontario, N1C 0A1 Contact Phone Number: (519) 767-9262

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NipsIt is a trademark of Valent U.S.A. Corporation.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is

2019-1665
2019-05-17

inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

FIRST AID:

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

No specific antidote is available. Treat the patient symptomatically.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Harmful or fatal if swallowed. Harmful if absorbed through skin. Avoid contact with skin or clothing. Wash hands thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Seed Treatment Workers: Workers mixing, loading, treating, clean-up, or maintaining and repairing seed treatment equipment, bagging, sewing, or stacking treated seed must wear coveralls over long-sleeved shirt, long pants, chemical-resistant gloves, head gear, and respiratory protection (i.e., NIOSH-approved dust-filtering respirator or fresh air hood).

Commercial planters must use a closed cab planter when planting treated seeds.

User Safety Recommendations:

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove personal protective equipment (PPE) immediately after handling this product.
- Wash the outside of gloves before removing.
- As soon as possible, wash thoroughly and change into clean clothing.
- Follow manufacturer's instructions for cleaning/maintaining. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

2019-1665
2019-05-17

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's website at: www.croplife.ca.

ENVIRONMENTAL HAZARDS:

Toxic to aquatic organisms.

Treated seed is toxic to birds and small wild animals. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface.

Toxic to bees. Bees can be exposed to product residues in flowers, leaves, pollen and/or nectar resulting from seed treatments.

This product demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of Nipsit Suite Canola Seed Protectant in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

USE RESTRICTIONS:

- Do not use treated seed for food, feed, or oil processing.
- Due to seed quality conditions beyond the control of Valent Canada, Inc., no claims are made to guarantee germination of carry-over seed.
- Rape greens and seed grown or harvested from NipsIt SUITE Canola Seed Protectant-treated seed must not be used for feed or human consumption.
- Rapeseed and carinata grown and harvested from NipsIt SUITE Canola Seed Protectant-treated seed is only for industrial uses and can not be used for edible oil or any other human/feed consumption.
- Do not graze or feed livestock on treated areas for four weeks after planting.

LABELLING OF TREATED SEED:

All bags containing treated seed must be labeled or tagged as follows:

- This seed was treated with a product containing the active ingredients metconazole, clothianidin, and metalaxyl.
- Do not use for feed, food or oil processing.
- Rape greens and seed grown or harvested from NipsIt SUITE Canola Seed Protectant-treated seed must not be used for feed or human consumption.
- Store away from feeds and other foodstuffs.
- Wear long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seed.
- This product is toxic to aquatic invertebrates.
- Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements.
- Dispose of seed packaging in accordance with local requirements.
- Cover or incorporate spilled treated seeds.

2019-1665
2019-05-17

STORAGE AND DISPOSAL:

Do not contaminate water, food or feed by storage, disposal, or by cleaning of equipment.

PESTICIDE STORAGE: Store in a cool place. Do not store in direct sunlight. Protect from freezing temperatures.

PESTICIDE DISPOSAL: For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

CONTAINER DISPOSAL:

RECYCLABLE CONTAINER:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

RETURNABLE CONTAINER:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

REFILLABLE CONTAINER:

1. After use, return the container to the point of purchase or designated locations.
2. This container must only be refilled with NipsIt SUITE Canola Seed Protectant at designated locations.
3. Do not reuse container for other purposes.
4. Prior to refilling inspect thoroughly for damage.
5. Check for leaks after refilling and before transportation.
6. Do not refill or transport damaged or leaking containers.
7. If the container is not being refilled, return to the point of purchase.

NON-RECYCLABLE/RETURNABLE/REFILLABLE CONTAINER:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.



**** CONTAINER****

GROUP	3	4	FUNGICIDE
GROUP		4	INSECTICIDE

NipsIt™ SUITE Canola Seed Protectant

Flowable Insecticide and Fungicide Seed Protectant

SUSPENSION

NipsIt SUITE Canola Seed Protectant is a systemic insecticide and fungicide seed protection product for use on canola, carinata and rapeseed for early season control of flea beetles, seed and seedling diseases caused by *Pythium* spp., *Fusarium* spp. and *Rhizoctonia solani* and suppression of seed-borne blackleg (*Leptosphaeria maculans*).

For use in commercial seed treatment facilities with closed transfer including closed mixing, loading, calibrating, and closed treatment equipment only. No open transfer of NipsIt SUITE Canola Seed Protectant.

COMMERCIAL

GUARANTEE:

Clothianidin 279 g/L
Metalaxyl.....5.23 g/L
Metconazole.....1.04 g/L

Contains 5-chloro-2-methyl-4-isothiazolin-3-one at 0.00113% and 2-methyl-4-isothiazolin-3-one at 0.00037% as preservatives.

REGISTRATION NUMBER 31355 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

CAUTION

<<Skull and Cross Bones in Diamond>>

POISON

NET CONTENTS: 72 L - 14400 L (bulk tanker)

This product contains blue colourant for application to canola/rapeseed, according to regulations pertaining to the “Seeds Act

HEALTH EMERGENCY OR SPILL (24 hr):

2019-1665
2019-05-17

(800) 682-5368
TRANSPORTATION (24 hr.): CHEMTREC
(800) 424-9300 or (202) 483-7616

GROWER INFORMATION:

Valent Canada, Inc.

201-230 Hanlon Creek Blvd., Guelph, Ontario, N1C 0A1 Contact Phone Number: (519) 767-9262

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

FIRST AID:

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

No specific antidote is available. Treat the patient symptomatically.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Harmful or fatal if swallowed. Harmful if absorbed through skin. Avoid contact with skin or clothing. Wash hands thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Seed Treatment Workers: Workers mixing, loading, treating, clean-up, or maintaining and repairing seed treatment equipment, bagging, sewing, or stacking treated seed must wear coveralls over long-sleeved shirt, long pants, chemical-resistant gloves, head gear, and respiratory protection (i.e., NIOSH -approved dust-filtering respirator or fresh air hood).

Commercial planters must use a closed cab planter when planting treated seeds.

2019-1665
2019-05-17

User Safety Recommendations:

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove personal protective equipment (PPE) immediately after handling this product.
- Wash the outside of gloves before removing.
- As soon as possible, wash thoroughly and change into clean clothing.
- Follow manufacturer's instructions for cleaning/maintaining. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL HAZARDS:

Toxic to aquatic organisms.

Treated seed is toxic to birds and small wild animals. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface.

Toxic to bees. Bees can be exposed to product residues in flowers, leaves, pollen and/or nectar resulting from seed treatments.

This product demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of Nipsit Suite Canola Seed Protectant in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the US visit CropLife Canada's Website at www.Croplife.ca.

STORAGE AND DISPOSAL:

Do not contaminate water, food or feed by storage, disposal, or by cleaning of equipment.

PESTICIDE STORAGE: Store in a cool place. Do not store in direct sunlight. Protect from freezing temperatures.

PESTICIDE DISPOSAL: For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

CONTAINER DISPOSAL:

RECYCLABLE CONTAINER:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

2019-1665
2019-05-17

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

RETURNABLE CONTAINER:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

REFILLABLE CONTAINER:

1. After use, return the container to the point of purchase or designated locations.
2. This container must only be refilled with NipsIt SUITE Canola Seed Protectant at designated locations.
3. Do not reuse container for other purposes.
4. Prior to refilling inspect thoroughly for damage.
5. Check for leaks after refilling and before transportation.
6. Do not refill or transport damaged or leaking containers.
7. If the container is not being refilled, return to the point of purchase.

NON-RECYCLABLE/RETURNABLE/REFILLABLE CONTAINER:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.

NipsIt is a trademark of Valent U.S.A. Corporation



** BOOKLET**

GROUP	3	4	FUNGICIDE
GROUP	4		INSECTICIDE

NipsIt™ SUITE Canola Seed Protectant

Flowable Insecticide and Fungicide Seed Protectant

SUSPENSION

NipsIt SUITE Canola Seed Protectant is a systemic insecticide and fungicide seed protection product for use on canola, carinata and rapeseed for early season control of flea beetles, seed and seedling diseases caused by *Pythium* spp., *Fusarium* spp. and *Rhizoctonia solani* and suppression of seed-borne blackleg (*Leptosphaeria maculans*).

For use in commercial seed treatment facilities with closed transfer including closed mixing, loading, calibrating, and closed treatment equipment only. No open transfer of NipsIt SUITE Canola Seed Protectant.

AGRICULTURAL

GUARANTEE:

Clothianidin	279 g/L
Metalaxyl.....	5.23 g/L
Metconazole.....	1.04 g/L

Contains 5-chloro-2-methyl-4-isothiazolin-3-one at 0.00113% and 2-methyl-4-isothiazolin-3-one at 0.00037% as preservatives.

REGISTRATION NUMBER 313155 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

CAUTION

<<Skull and Cross Bones in Diamond>>

POISON

NET CONTENTS: 72 L - 14400 L (bulk tanker)

This product contains blue colourant for application to canola/rapeseed, according to regulations pertaining to the “Seeds Act”.

2019-1665
2019-05-17

HEALTH EMERGENCY OR SPILL (24 hr):
(800) 682-5368
TRANSPORTATION (24 hr.): CHEMTREC
(800) 424-9300 or (202) 483-7616

GROWER INFORMATION:
Valent Canada, Inc.
201-230 Hanlon Creek Blvd., Guelph, Ontario, N1C 0A1 Contact Phone Number: (519) 767-9262

2019-1665
2019-05-17

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

FIRST AID:

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

No specific antidote is available. Treat the patient symptomatically.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Harmful or fatal if swallowed. Harmful if absorbed through skin. Avoid contact with skin or clothing. Wash hands thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Seed Treatment Workers: Workers mixing, loading, treating, clean-up, or maintaining and repairing seed treatment equipment, bagging, sewing, or stacking treated seed must wear coveralls over long-sleeved shirt, long pants, chemical-resistant gloves, head gear, and respiratory protection (i.e., NIOSH -approved dust-filtering respirator or fresh air hood).

Commercial planters must use a closed cab planter when planting treated seeds.

User Safety Recommendations:

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove personal protective equipment (PPE) immediately after handling this product.
- Wash the outside of gloves before removing.

2019-1665
2019-05-17

- As soon as possible, wash thoroughly and change into clean clothing.
- Follow manufacturer's instructions for cleaning/maintaining. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's website at: www.croplife.ca.

ENVIRONMENTAL HAZARDS:

Toxic to aquatic organisms.

Treated seed is toxic to birds and small wild animals. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface.

Toxic to bees. Bees can be exposed to product residues in flowers, leaves, pollen and/or nectar resulting from seed treatments.

This product demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of Nipsit Suite Canola Seed Protectant in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

USE RESTRICTIONS:

- Do not use treated seed for food, feed, or oil processing.
- Due to seed quality conditions beyond the control of Valent Canada, Inc., no claims are made to guarantee germination of carry-over seed.
- Rape greens and seed grown or harvested from NipsIt SUITE Canola Seed Protectant-treated seed must not be used for feed or human consumption.
- Rapeseed and carinata grown and harvested from NipsIt SUITE Canola Seed Protectant-treated seed is only for industrial uses and can not be used for edible oil or any other human/feed consumption.
- Do not graze or feed livestock on treated areas for four weeks after planting.

LABELLING OF TREATED SEED:

All bags containing treated seed must be labeled or tagged as follows:

- This seed was treated with a product containing the active ingredients metconazole, clothianidin, and metalaxyl.
- Do not use for feed, food or oil processing.
- Rape greens and seed grown or harvested from NipsIt SUITE Canola Seed Protectant-treated seed must not be used for feed or human consumption.
- Store away from feeds and other foodstuffs.
- Wear long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seed.
- This product is toxic to aquatic invertebrates.
- Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements.
- Dispose of seed packaging in accordance with local requirements.

2019-1665
2019-05-17

- Cover or incorporate spilled treated seeds.
- Commercial planters must use a closed cab planter when planting treated seeds.

STORAGE AND DISPOSAL:

Do not contaminate water, food or feed by storage, disposal, or by cleaning of equipment.

PESTICIDE STORAGE: Store in a cool place. Do not store in direct sunlight. Protect from freezing temperatures.

PESTICIDE DISPOSAL: For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

CONTAINER DISPOSAL:

RECYCLABLE CONTAINER:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

RETURNABLE CONTAINER:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

REFILLABLE CONTAINER:

1. After use, return the container to the point of purchase or designated locations.
2. This container must only be refilled with NipsIt SUITE Canola Seed Protectant at designated locations.
3. Do not reuse container for other purposes.
4. Prior to refilling inspect thoroughly for damage.
5. Check for leaks after refilling and before transportation.
6. Do not refill or transport damaged or leaking containers.
7. If the container is not being refilled, return to the point of purchase.

NON-RECYCLABLE/RETURNABLE/REFILLABLE CONTAINER:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.

GENERAL INFORMATION:

NipsIt SUITE Canola Seed Protectant is a systemic insecticide and fungicide seed protectant product for use on canola, carinata and rapeseed for early season control of flea beetle, seed and seedling diseases caused by *Pythium* spp., *Fusarium* spp. and *Rhizoctonia solani* and suppression of seed-borne blackleg (*Leptosphaeria maculans*). Clothianidin, the insecticide component within NipsIt SUITE Canola Seed Protectant, provides seedling protection against early season flea beetle feeding damage. The fungicide components of NipsIt SUITE Canola Seed Protectant are metalaxyl and metconazole. Metalaxyl protects seed and seedlings against seed rot, damping-off, seedling blight and root rot associated with *Pythium* spp. Metconazole fungicide gives protection against seed rot and damping off associated with *Fusarium* spp., post-emergence damping-off associated with *Rhizoctonia solani* and seed-borne blackleg (*Leptosphaeria maculans*).

Clothianidin is a Group 4 Insecticide. Do not make any subsequent application of a Group 4 Insecticide (i.e., in-furrow or foliar application) following treatment with NipsIt SUITE Canola Seed Protectant.

For use in commercially available equipment designed for seed treatment only. Not for use in hopper-box, slurry-box, or similar seed treatment applications used at planting. This product is to be used in liquid or slurry treaters. Mix thoroughly before use or use entire container at one time.

DIRECTIONS FOR USE:

NipsIt SUITE Canola Seed Protectant is a ready to use product, which includes blue colourant (according to regulations pertaining to the “Seeds Act”). No additional colourant, dyes, binders, polymers, or water are required.

Calibrate seed treatment equipment prior to application of NipsIt SUITE Canola Seed Protectant to assure proper rate of application is achieved. Consult with the seed treatment equipment manufacturer for guidance on the operation and proper calibration of the treating equipment, as non-uniform, inaccurate seed coverage may result suboptimal insect or disease protection to canola, carinata and rapeseed seedlings.

Apply 1.43 L/100 KG seed of NipsIt SUITE Canola Seed Protectant using standard commercial seed treatment equipment to secure uniform seed coverage. Allow treated seed to dry prior to bagging.

RECOMMENDED APPLICATION

Crop	Insect Claims	Disease Claims	Application Rate of NipsIt SUITE Canola Seed Protectant	Remarks

Canola, Carinata, Rapeseed	Flea Beetles	<p>Suppression of seed-borne blackleg (<i>Leptosphaeria maculans</i>)</p> <p>Seed rot/pre-emergence damping-off and post-emergence damping-off, caused by <i>Fusarium</i> spp.</p> <p>Post-emergence damping-off caused by <i>Rhizoctonia solani</i></p> <p>Seed rot/pre-emergence damping-off, post-emergence damping-off, seedling blight and seedling root rot caused by <i>Pythium</i> spp.</p>	1.43 L/100 kg seed	Do not make any subsequent application of a Group 4 Insecticide (i.e., in-furrow or foliar application) following treatment with NipsIt SUITE Canola Seed Protectant.
----------------------------------	--------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------

SEED QUALITY AND STORAGE OF TREATED SEED

- Laboratory and field studies have shown that NipsIt SUITE Canola Seed Protectant applied to canola seed will not adversely affect germination.
- Due to seed quality conditions beyond the control of Valent Canada, Inc., no claims are made to guarantee germination of carry-over seed. Treatment of mechanically scarred or damaged seed, or of seed known to be of low germ, low vigour and poor quality is not recommended.
- Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Valent Canada, Inc., no claims are made to guarantee germination of carry-over seed.

SEEDING EQUIPMENT:

NipsIt SUITE Canola Seed Protectant-treated seeds may not flow through seeding equipment at the same rate as other canola seed treatment products or untreated seed. Re-calibration of seeding equipment is recommended before planting seed treated with NipsIt SUITE Canola Seed Protectant to insure proper planting rate in the field.

ROTATIONAL CROP RESTRICTIONS:

Canola, carinata and rapeseed may be replanted at any time. A one year plantback interval is required for leafy, root and tuber vegetables. A 30-day plantback interval on cereal grains, grasses, non-grass animal feeds and soybeans and dried beans is required.

RESISTANCE MANAGEMENT RECOMMENDATIONS:

2019-1665
2019-05-17

For resistance management, please note that NipsIt SUITE Canola Seed Protectant contains a Group 4 insecticide and Group 3 and 4 fungicides. Any insect or fungal population may contain individuals naturally resistant to NipsIt INSIDE Canola Seed Protectant and other Group 4 insecticides or Group 3 and 4 fungicides. A gradual or total loss of pest control may occur over time if these insecticides or fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance:

- Where possible, rotate the use of NipsIt SUITE Canola Seed Protectant or other Group 4 insecticides and Groups 3 and 4 fungicides with different groups that control the same insects or pathogens.
- Insecticide and fungicide use should be based on an IPM program that includes scouting, record keeping, and considers cultural, biological and other chemical control practices.
- Monitor treated insect and fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.

For further information or to report suspected resistance contact Valent Canada, Inc. at 1-800-682-5368.

NipsIt is a trademark of Valent U.S.A. Corporation