

2023-2406
2024-11-07

(Container Label)

GROUP	2	HERBICIDE
-------	---	-----------

GP Herbicide

**FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND
INTERIOR OF BRITISH COLUMBIA (INCLUDING THE PEACE RIVER REGION) ONLY**

COMMERCIAL - AGRICULTURAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: pyroxsulam 21.5%

Wettable granules

REGISTRATION NO. 32068 PEST CONTROL PRODUCTS ACT

CAUTION: EYE IRRITANT

NET CONTENTS: 0.5 kg – bulk

FMC of Canada Limited
6755 Mississauga Road, Suite 204
Mississauga, ON L5N 7Y2
1-833-362-7722

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN CAUTION EYE IRRITANT

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

At all times: Wear clean clothing with long sleeved shirt and long pants. Wear coveralls over long sleeved shirt and long pants, chemical-resistant gloves, socks and chemical resistant footwear during mixing, loading, application, clean-up and repair. Wear goggles or face shield during mixing/loading. For closed cab or aerial application, coveralls and gloves are not necessary.

Apply to agricultural crops only when the potential for drift to areas of human habitation and areas of human activity, such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

At completion of spraying or end of the day: Take a shower immediately. Wash thoroughly with soap and water before eating, drinking or smoking. Remove contaminated clothing at the end of the work session and store and wash separately from household laundry using detergents and hot water before reuse.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IN CASE OF EMERGENCY, CALL TOLL FREE, DAY OR NIGHT: 1-800-331-3148

TOXICOLOGICAL INFORMATION

No specific antidote. Employ supportive care. Treatment should be based on judgment of the physician in response to reactions of the patient.

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE. To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

2023-2406
2024-11-07

STORAGE

Keep in original container during storage. Store product in cool, dry, well ventilated place away from seed, feed, fertilizer, or other pesticides. Keep away from fire or open flame, or other source of heat. Store this product away from food or feed.

DISPOSAL

Recyclable Containers

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

Refillable Containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

Returnable Containers

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

2023-2406
2024-11-07

(Directions for Use)

GROUP	2	HERBICIDE
-------	---	-----------

GP Herbicide

**FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND
INTERIOR OF BRITISH COLUMBIA (INCLUDING THE PEACE RIVER REGION) ONLY**

COMMERCIAL - AGRICULTURAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: pyroxsulam 21.5%

Wettable granules

REGISTRATION NO. 32068 PEST CONTROL PRODUCTS ACT

CAUTION: EYE IRRITANT

FMC of Canada Limited
6755 Mississauga Road, Suite 204
Mississauga, ON L5N 7Y2
1-833-362-7722

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN CAUTION EYE IRRITANT

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

At all times: Wear clean clothing with long sleeved shirt and long pants. Wear coveralls over long sleeved shirt and long pants, chemical-resistant gloves, socks and chemical resistant footwear during mixing, loading, application, clean-up and repair. Wear goggles or face shield during mixing/loading. For closed cab or aerial application, coveralls and gloves are not necessary.

Apply to agricultural crops only when the potential for drift to areas of human habitation and areas of human activity, such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

At completion of spraying or end of the day: Take a shower immediately. Wash thoroughly with soap and water before eating, drinking or smoking. Remove contaminated clothing at the end of the work session and store and wash separately from household laundry using detergents and hot water before reuse.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IN CASE OF EMERGENCY, CALL TOLL FREE, DAY OR NIGHT: 1-800-331-3148

TOXICOLOGICAL INFORMATION

No specific antidote. Employ supportive care. Treatment should be based on judgment of the physician in response to reactions of the patient.

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE. To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

GENERAL INFORMATION

GP Herbicide is a selective postemergence herbicide for the control of wild oats and certain broadleaved weeds in spring wheat, durum wheat and winter wheat not underseeded with legumes. GP Herbicide is mixed with water and applied as a uniform broadcast spray. It is non-corrosive, non-flammable, and non-volatile.

GP Herbicide must be applied early postemergence, to the main flush of actively growing wild oats and broadleaved weeds. Conditions favouring active weed growth enhance the activity of GP Herbicide by allowing maximum foliar uptake and contact activity. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and re-growth may occur. For best results, ensure thorough spray coverage of target weeds. See "DIRECTIONS FOR USE" section of this label for complete use details.

GP Herbicide stops growth of susceptible weeds rapidly. However, typical symptoms (discolouration) of dying weeds may not be noticeable for 1 to 3 weeks after application, depending upon growing conditions and weed susceptibility. Degree of control and duration of effect are dependent on weed sensitivity, weed size, crop competition, growing conditions at and following treatment, and spray coverage.

MODE OF ACTION

GP Herbicide inhibits the production of the ALS enzyme in plants. This enzyme is essential for the production of certain amino acids which are essential for plant growth. GP Herbicide is a Group 2 mode of action herbicide.

GENERAL USE PRECAUTIONS

Non-Target Sites

Do not apply where proximity of susceptible crops (e.g. canola and legumes) or other desirable plants is likely to result in exposure to spray or spray drift. See Environmental Hazards section of the label.

Crop Rotation

Fields previously treated with GP Herbicide can be seeded 11 months following an application of GP Herbicide, with the following crops: barley, brown mustard, canola, dry bean (of the species *Phaseolus vulgaris*), flax, canola quality Brassica juncea, lentils, oats, field peas, chickpea, spring wheat, soybean and yellow mustard or fields can be summerfallowed. Fields previously treated with GP Herbicide can be seeded 10 months following an application of GP Herbicide with the following crops: field corn, sunflower and potatoes.

TANK MIXES:

This product may be tank mixed with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact FMC of Canada Limited at 1-833-362-7722 for information before applying any tank mix that is not specifically recommended on this label.

Spray Equipment Precaution

Do not apply through any type of irrigation system.

Sprayer Clean-Out Instructions

To avoid injury to desirable plants, thoroughly clean equipment used to apply this product before re-use or using it to apply other chemicals.

1. Immediately after spraying, completely drain the sprayer tank. Any contamination on the outside of the spraying equipment should be removed by washing with clean water.
2. First rinse:

- Spray the inside of tank with clean water and fill the sprayer with at least one tenth of the spray tank volume.
 - Agitate and circulate for 15 minutes, and flush through booms and hoses.
 - Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
 - Drain tank completely.
3. Second rinse:
- Fill the tank with clean water.
 - Add All Clear Spray Tank Decontaminator as per manufacturer's recommendations while filling the tank with clean water.
 - Agitate and then flush the boom and hoses with the cleaning solution. Top up with water making sure the tank is completely full. Allow to stand for 15 minutes with agitation. Flush the solution out of the spray tank through the spray booms. Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
 - If possible, let the solution stand in the sprayer tank and booms for an extended period of time, overnight if possible.
 - After flushing the boom and hoses, drain tank completely.
 - Remove nozzles and screens and clean separately with a cleaning agent or an ammonia solution (100 mL in 10 L water).
4. Third rinse:
- Rinse the tank with clean water and flush through the boom and hoses using at least one tenth of the spray tank volume.
 - Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
 - Drain tank completely.

Do not use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty odour which may cause eye, nose, throat and lung irritation. Do not clean equipment in an enclosed area.

DIRECTIONS FOR USE

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND INTERIOR OF BRITISH COLUMBIA (INCLUDING THE PEACE RIVER REGION) ONLY

READ THE ENTIRE LABEL BEFORE USE. FAILURE TO FOLLOW LABEL INSTRUCTIONS MAY RESULT IN ERRATIC WEED CONTROL OR CROP DAMAGE. DO NOT APPLY TO CROPS UNDERSEEDED WITH LEGUMES.

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. Boom height must be 60 cm or less above the crop or ground.

Aerial application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 km/h at flying height at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotorspan.

GP HERBICIDE ALONE

Application Directions - Spring Wheat, Durum Wheat

Apply GP Herbicide at a rate of 70 g per hectare for control of grass and broadleaf weeds (see below). For control of low wild oat populations (<75 plants/m²) only, apply GP Herbicide at a rate of 52 g per hectare. Always add a non-ionic surfactant (Agra*I 90, Ag-Surf* or Surf* 92) at 0.25 v/v (0.25 L per 100 L of spray solution) when applying GP Herbicide alone. Apply using aerial equipment or ground equipment that will assure uniform coverage. See table below for weeds species controlled. GP Herbicide will control only emerged weeds. Weeds that germinate after application will not be controlled.

Weeds Controlled or Suppressed with GP Herbicide

Weeds Controlled:

barnyard grass (1-5 leaf)	foxtail, yellow (1-5 leaf)
brome, Japanese ^{††} (1-6 leaf)	hemp nettle (1-8 leaf)
canola, volunteer [†] (1-6 leaf)	oats, wild (up to 4 leaf, 2 tiller)
chickweed, common (up to 10 cm)	pigweed, red-root (1-8 leaf)
cleavers (up to 6 whorl)	round-leaved mallow (up to 6 leaf stage, <10 cm in size)
corn spurry (up to 2 whorl stage, <10 cm in height)	smartweed (lady's-thumb, 1-5 leaf)
cow cockle (up to 8 leaf)	shepherd's purse (up to 30 cm tall)
flixweed (up to 10 cm)	stinkweed (up to 30 cm tall)

[†]will not control volunteer Imidazolinone-tolerant canola

^{††} For control of Japanese brome from the 1 leaf to 4 leaf, 2 tiller stage under good growing conditions, apply GP Herbicide at the 52 g/ha rate. For larger Japanese brome (up to the 6 leaf stage) and less than optimum growing conditions, use the 70 g/ha rate.

Weeds Suppressed:

brome, downy (2-6 leaf, 4 tillers)	foxtail, green (1-5 leaf)
dandelion (seedlings and over-wintered rosettes less than or equal to 20 cm)	wild buckwheat (1-4 leaf)
white cockle (up to the first flower stage, less than 20 cm in height)	Canada thistle (up to 30 cm tall, prebud)
	Russian thistle (up to 10 cm tall)
	Persian darnel (1 leaf to 4 leaf, 2 tiller)

Application Timing

Apply to actively growing wheat from the 3 leaf stage to before the flag leaf stage. Apply when weeds are actively growing: grass weeds up to the 4 leaf, 2 tiller stage and broadleaved weeds as indicated in the table above. Only weeds emerged at the time of treatment will be controlled. Best results are obtained from applications made to seedling weeds. Extreme growing conditions such as drought or near freezing temperature prior to, at or following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. Under conditions of low crop and high weed density, control may be reduced.

Application Directions - Winter Wheat (Fall or Spring Application) for Downy Brome

GP Herbicide applied in the fall will provide control of downy brome while a spring application will provide suppression. Apply GP Herbicide at a rate of 70 g per hectare. Add BASF Assist Oil Concentrate at 0.8% v/v (0.8 L/100 L of spray mixture) or Merge* Adjuvant at 0.5% v/v (0.5 L/100 L of spray mixture; spring

application only). Apply using aerial equipment or ground equipment that will assure uniform coverage. In addition to suppression of downy brome when applied in the spring GP Herbicide will also control/suppress weeds listed. GP Herbicide will control only emerged weeds. Weeds that germinate after application will not be controlled.

Application Timing

Apply in the fall when winter wheat is in the 1-3 leaf stage or in the spring when winter wheat is in the 2-7 leaf, 4 tiller stage. For fall applications, apply when downy brome is actively growing at the 1 to 3 leaf stage; for spring applications, apply when weeds are actively growing at the stages indicated. Only weeds emerged at the time of treatment will be controlled. Best results are obtained from applications made to seedling weeds. Extreme growing conditions such as drought or near freezing temperature prior to, at or following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. Under conditions of low crop and high weed density, control may be reduced.

Occasionally slight yellowing or height reduction may be observed in the treated crop. These transient symptoms disappear within 14 days with no reduction to yield. Do not apply to crops suffering from drought, nutrient deficiency or exposed to frost or other agronomic factors affecting plant growth.

If foliage is wet at the time of application control may be decreased. Applications of GP Herbicide are rainfast within 2 hours after application.

Mixing Instructions (GP Herbicide used alone)

1. Fill sprayer tank 1/2 full of water.
2. Start sprayer tank agitation.
3. Add the required amount of GP Herbicide
4. Fill the sprayer tank with sufficient water to spray 30-100 L of spray mixture per hectare.
5. Add the required surfactant.
6. Follow sprayer directions and precautions as outlined above, especially when applying next to sensitive crops (e.g. canola and legumes).
7. Follow sprayer clean-up directions.

Pre-harvest/Grazing Intervals

Livestock may be grazed on treated crops 7 days following application. Do not harvest the treated crop within 60 days after application.

TANK MIX COMBINATIONS – FOR USE IN SPRING WHEAT AND DURUM WHEAT ONLY

GP Herbicide can be tank mixed with herbicides and/or fungicides as indicated in the table below.

For application of GP Herbicide in tank mixtures, only add surfactant where the label of the tank mix partner product indicates its use is required, in accordance with that label. See table below for surfactant recommendations for specific tank mixtures.

Tank Mix Combinations with GP Herbicide at 70 g/ha in Spring Wheat and Durum Wheat

Herbicide Tank Mix Partner	Fungicide Tank-Mix Partner	Rate of herbicide TM partner (Product /ha)	Additional Pests Controlled
GP Herbicide alone	Tilt* 250 E or Stratego* 250 EC	70 g/ha Consult fungicide tank-mix partner label for fungicide rates.	Consult fungicide tank-mix partner label for diseases controlled
Attain*XC A Herbicide + Attain* XC B Herbicide Tank-Mix	Tilt* 250 E or Stratego* 250 EC	0.23 L/ha Attain* XC A Herbicide + 0.64 L/ha Attain* XC B Herbicide	stork's-bill (1-8 leaf) wild buckwheat + labelled weeds for Low Rate of Attain* XC A Herbicide + Attain* XC B Herbicide tank-mix

		Consult fungicide tank-mix partner label for fungicide rates	Consult fungicide tank-mix partner label for diseases controlled
OcTTain* XL Herbicide		1.1 L/ha	Consult label
Curtail* M Herbicide		1.5 L/ha	annual sowthistle (1-5 leaf) consult label
Frontline*XL Herbicide		1.25 L/ha	annual sowthistle (1-5 leaf) stork's-bill (1-8 leaf) consult label
Frontline* 2,4-D Herbicide Tank-Mix		66.7 mL/ha Frontline 2,4-D A + 667 mL/ha Frontline 2,4-D B	annual sowthistle (1-5 leaf) stork's-bill (1-8 leaf) wild buckwheat (1-7 leaf)+ weeds controlled by 2,4-D ester at 350 g a.e./ha ^{††}
Spectrum* Herbicide Tank-Mix		100 mL/ha Spectrum* A + 1.5 L/ha Spectrum* B	consult label
MCPA Ester 600	Tilt* 250 E or Stratego* 250 EC	0.58 – 0.92 L/ha Consult fungicide tank-mix partner label for fungicide rates	stork's-bill (1-8 leaf) ^{†††} wild buckwheat ^{†+} weeds labelled for this rate range of MCPA Consult fungicide tank-mix partner label for diseases controlled
Buctril* M		1.0 L/ha	consult label
Refine® SG Herbicide		30 g/ha A registered adjuvant is required for this mixture	consult label
PP-3317 Herbicide		30 g/ha A registered adjuvant is required for this mixture	consult label
PP-2525 Herbicide		30 g/ha A registered adjuvant is required for this mixture	consult label
Thumper* Emulsifiable Selective Weedkiller		1.0 L/ha	consult label
Prestige* XC Herbicide Tank-Mix		0.31 L/ha Prestige XC A + 1.5 L/ha Prestige XC B	annual sowthistle (1-5 leaf) wild buckwheat (1-9 leaf) Russian thistle (2-10 cm high)
2,4-D ester		280 g a.e./ha (e.g. 425 mL/ha of 2,4-D Ester 700)	volunteer canola (1-6 leaf stage)* lamb's quarters (1-6 leaf stage) *Including imidazolinone tolerant volunteer canola

MCPA ester		280 g a.e./ha (e.g. 465 mL/ha of MCPA Ester 600)	volunteer canola (1-6 leaf stage)* lamb's quarters (1-6 leaf stage) *Including imidazolinone tolerant volunteer canola
-------------------	--	--	--

†For larger weeds or heavy infestations use MCPA at 1.05 L/ha

††Seedlings at the 2-4 leaf stage; actively growing mustard (except dog and green tansy), flixweed, bluebur, burdock, cocklebur, field horsetail (top growth control only), stinkweed, goats beard, prickly lettuce, hoary cress (top growth control only), kochia, lamb's-quarters, plantain, ragweeds, Russian knapweed, Russian thistle, shepherd's-purse, annual sunflower, sweet clover, vetch, wild radish

†††Use the 1.05 L/ha rate for control of stork's-bill

Tank Mix Combinations with GP Herbicide at 52 g/ha in Spring Wheat and Durum Wheat Under Low Wild Oat Populations (<75 plants/m²)

Herbicide Tank Mix Partner	Fungicide Tank Mix	Rate (Product /ha)	Additional Weeds Controlled
Attain* XC A Herbicide + Attain* XC B Herbicide Tank-Mix	Tilt* 250 E or Stratego* 250 EC	0.23 L/ha Attain* XC A Herbicide + 0.64 L/ha Attain* XC B Herbicide Consult fungicide tank-mix partner label for fungicide rates.	cow cockle (up to 8 leaf), wild oats, wild buckwheat (1-4 leaf) + labelled weeds for Low Rate of Attain XC A Herbicide + Attain XC B Herbicide tank-mix Consult fungicide tank-mix partner label for diseases controlled
OcTTain* XL Herbicide		1.1 L/ha	Consult label

Mixing Instructions (GP Herbicide tank mixed with other products)

1. Fill sprayer tank 1/2 full of water.
2. Start sprayer tank agitation.
3. Add the required amount of GP Herbicide
4. Add the correct amount of tank-mix partner and continue to agitate.
5. Fill the sprayer tank with the remaining amount of water required.

Notes: Follow tank-mix partner label for order of mixing.

APPLICATION INSTRUCTIONS (GROUND AND AERIAL APPLICATIONS)

(1) Ground Application (GP Herbicide applied alone and with all tank mix partners)

Using ground equipment, apply GP Herbicide alone or with tank mix partners as a broadcast treatment. Apply GP Herbicide alone at the recommended rates in a spray volume of 30-100 L water / ha.

(2) Aerial Application

Directions for Use

Do not use human flaggers.

2023-2406
2024-11-07

Use GP Herbicide alone or in tank mix with either MCPA ester or Buctril* M as a broadcast treatment by air. Apply the recommended rates of GP Herbicide and either MCPA ester or Buctril* M as indicated above. Apply in a minimum spray volume of 30 L water / ha.

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate swath marking devices.

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing should be laundered regularly.

Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-833-362-7722 or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

- The interaction of many equipment-and-weather-related factors determine the potential for spray drift. Users are responsible for considering all these factors when making decisions.
- The following drift management requirement must be followed to avoid off-target drift movement from aerial applications:
 - Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

SPRAY BUFFER ZONES

A spray buffer zone is NOT required for:

- uses with hand-held application equipment permitted on this label

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), and sensitive freshwater habitats

(such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

Method of application	Crop		Spray Buffer Zones (metres) Required for the Protection of:	
			Freshwater Habitat of Depths	Terrestrial Habitat
Field sprayer	Spring wheat, durum wheat, winter wheat		1	2
Aerial	Spring wheat, durum wheat, winter wheat	Fixed wing	1	70
		Rotary wing	1	60

When a tank mixture is used, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture. Apply using the coarsest spray (ASAE) category indicated on the label for those tank-mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, GP Herbicide is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to GP Herbicide and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of GP Herbicide or other Group 2 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact FMC of Canada Limited at 1-833-362-7722.

STORAGE

Keep in original container during storage. Store product in cool, dry, well ventilated place away from seed, feed, fertilizer, or other pesticides. Keep away from fire or open flame, or other source of heat. Store this product away from food or feed.

DISPOSAL

Recyclable Containers

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

Refillable Containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

Returnable Containers

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

REFINE is a trademark of FMC Corporation or an affiliate.

* All other products mentioned are trademarks of their respective companies.