

LUNA PRIVILEGE GREENHOUSE

FUNGICIDE

SUSPENSION

For control of botrytis grey mould, botrytis leaf blight, botrytis stem canker and powdery mildew diseases in greenhouse cucumbers, peppers/eggplants (crop subgroup 8-09B) and tomatoes, early blight in greenhouse tomatoes, botrytis grey mould and powdery mildew in greenhouse strawberries and botrytis rot in greenhouse lettuce.

COMMERCIAL

ACTIVE INGREDIENT: Fluopyram500 g/L

Contains 1,2-benzisothiazolin-3-one at 0.029% and 5-chloro-2-methyl-4-isothiazolin-3-one at 0.0007% and 2-methyl-4-isothiazolin-3-one at 0.00025% as preservatives.

REGISTRATION NUMBER: 32208 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND BROCHURE BEFORE USING

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS: 0.5 L -Bulk

Distributed by:
Envu Canada
137 Glasgow Street, Suite 210
Office 111
Kitchener, ON N2G 4X8
Canada

Product Information: 1-800-331-2867
www.ca.envu.com

Registrant: Bayer CropScience Inc.

CAUTION: Do not apply under low light conditions as crop injury may occur

In case of spills, poisoning or fire, call telephone emergency response number 1-800-334-7577 (24 hours a day).

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

GENERAL INFORMATION:

THE PRODUCT:

LUNA PRIVILEGE GREENHOUSE is a broad-spectrum fungicide with preventative, systemic, and curative properties recommended for control of certain disease on vegetable and fruit plants in greenhouses. LUNA PRIVILEGE GREENHOUSE is best suited for use in a preventative treatment program.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Avoid contact with the skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. If product comes in contact with clothing, remove all contaminated clothing, wash skin with soap and water and dress in clean clothing. Launder applicator clothing separate from other laundry.

PROTECTIVE CLOTHING AND EQUIPMENT:

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair.

For Drip/Drench applications, when applied as a tank mix combination, read and observe all the label directions, including rates, personal protective equipment, restrictions and precautions for each product used in the tank-mix. Always use in accordance with the most restrictive label restrictions and precautions. When applied as a foliar application, DO NOT apply LUNA PRIVILEGE GREENHOUSE in tank mixture with other pesticides, fertilizers or adjuvants.

Do not apply this product using fogging equipment (handheld or automated) or using handheld mist blowers/airblast equipment.

RE-ENTRY RESTRICTION:

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when

seeking medical attention.

TOXICOLOGICAL INFORMATION:

Treat symptomatically. Medical Personnel should call emergency response number 1-800-334-7577 (24 hours a day).

ENVIRONMENTAL PRECAUTIONS:

Toxic to aquatic organisms and non-target terrestrial plants.

STORAGE:

Store this product away from food or feed. Do not contaminate water, food, or feed by storage or disposal. Do not store below freezing. If stored for 1 year or longer, shake well before using. Store the tightly closed container away from seeds, fertilizer, and plants and. Do not use or store in or around the home. Keep the product in the original container during storage.

DISPOSAL:

Recyclable Container Disposal:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

Disposal of Unused, Unwanted Product:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills.

Always read and follow label directions.

Luna Privilege™ is a trademark of the Bayer Group.

Bayer CropScience Inc. is a member of CropLife Canada.

230130

LUNA PRIVILEGE GREENHOUSE

FUNGICIDE

SUSPENSION

For control of botrytis grey mould, botrytis leaf blight, botrytis stem canker and powdery mildew diseases in greenhouse cucumbers, peppers/eggplants (crop subgroup 8-09B) and tomatoes, early blight in greenhouse tomatoes, botrytis grey mould and powdery mildew in greenhouse strawberries and botrytis rot in greenhouse lettuce.

COMMERCIAL

ACTIVE INGREDIENT: Fluopyram500 g/L

Contains 1,2-benzisothiazolin-3-one at 0.029% and 5-chloro-2-methyl-4-isothiazolin-3-one at 0.0007% and 2-methyl-4-isothiazolin-3-one at 0.00025% as preservatives.

REGISTRATION NUMBER: 32208 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND BROCHURE BEFORE USING

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS : 0.5 L – Bulk

Distributed by:
Envu Canada
137 Glasgow Street, Suite 210
Office 111
Kitchener, ON N2G 4X8
Canada

Product Information: 1-800-331-2867
www.ca.envu.com

Registrant: Bayer CropScience Inc.

In case of spills, poisoning or fire, call telephone emergency response number
1-800-334-7577 (24 hours a day)

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

GENERAL INFORMATION**THE PRODUCT:**

LUNA PRIVILEGE GREENHOUSE is a broad-spectrum fungicide with preventative, systemic, and curative properties recommended for control of certain disease on vegetable and fruit plants in greenhouses. LUNA PRIVILEGE GREENHOUSE is best suited for use in a preventative treatment program.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Avoid contact with the skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. If product comes in contact with clothing, remove all contaminated clothing, wash skin with soap and water and dress in clean clothing. Launder applicator clothing separate from other laundry.

PROTECTIVE CLOTHING AND EQUIPMENT:

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair.

For Drip/Drench applications, when applied as a tank mix combination, read and observe all the label directions, including rates, personal protective equipment, restrictions and precautions for each product used in the tank-mix. Always use in accordance with the most restrictive label restrictions and precautions. When applied as a foliar application, DO NOT apply LUNA PRIVILEGE GREENHOUSE in tank mixture with other pesticides, fertilizers or adjuvants.

Do not apply this product using fogging equipment (handheld or automated) or using handheld mist blowers/airblast equipment.

RE-ENTRY RESTRICTION:

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when

seeking medical attention.

TOXICOLOGICAL INFORMATION:

Treat symptomatically. Medical Personnel should call emergency response number 1-800-334-7577 (24 hours a day).

ENVIRONMENTAL PRECAUTIONS:

Toxic to aquatic organisms and non-target terrestrial plants.

STORAGE:

Store this product away from food or feed. Do not contaminate water, food, or feed by storage or disposal. Do not store below freezing. If stored for 1 year or longer, shake well before using. Store the tightly closed container away from seeds, fertilizer, and plants. Do not use or store in or around the home. Keep the product in the original container during storage.

DISPOSAL:

Recyclable Container Disposal:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

Disposal of Unused, Unwanted Product:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills.

DIRECTIONS FOR USE

Thorough coverage of all plant parts to be protected is essential for good disease control. Use sufficient water volume and spray pressure to provide thorough and uniform coverage for optimum disease control. Use a minimum water volume for thorough coverage of plant foliage.

DO NOT apply under low light conditions as crop injury may occur.

DO NOT allow releases, effluent or runoff from greenhouses containing this product to enter lakes, streams, ponds or other waters.

GREENHOUSE CUCUMBERS: Foliar Application		
Disease Control	Rate mL/hectare	Use Information

<p>Grey mould (fruit) Botrytis leaf blight Botrytis stem canker (<i>Botrytis cinerea</i>)</p>	<p>500</p>	<p>Use a minimum water volume for thorough coverage of plant foliage. Suggested spray volumes by crop height: 1.1 m - 1000 L/Ha spray volume 2.1 m - 2500 L/Ha spray volume</p> <p>Begin fungicide applications preventatively. Maximum of two applications per cropping cycle with a minimum re-application interval of 6 weeks. Rotate with non-Group 7 fungicides that control <i>Botrytis cinerea</i> 7 to 10-days later.</p> <p>Do not make a second application of LUNA PRIVILEGE GREENHOUSE within 6 weeks of the initial application.</p>
<p>Powdery Mildew (<i>Sphaerotheca fuliginea</i>) (<i>Erysiphe cichoracearum</i>)</p>	<p>100</p>	<p>Use a minimum water volume for thorough coverage of plant foliage. Suggested spray volumes by crop height: 1.1 m - 1000 L/Ha spray volume 2.1 m - 2500 L/Ha spray volume</p> <p>Begin fungicide applications preventatively. Maximum of three applications per cropping cycle with a minimum re-application interval of 6 weeks. Rotate with non-Group 7 fungicides that control Powdery mildew 7 to 14-days later.</p> <p>Do not make a second application of LUNA PRIVILEGE GREENHOUSE within 6 weeks of the initial application.</p>
<p>CAUTION: Not all varieties have been tested for phytotoxicity under all conditions. The user should test a small number of plants for sensitivity before considering application to the entire crop. When testing a small number of plants, wait at least one week after application to determine sensitivity. Do not apply to the crop if any negative or adverse crop response is observed, or if a test for sensitivity as not conducted.</p> <p>Restrictions: Do not apply more than 500 g of fluopyram per hectare per season. To limit the potential for development of disease resistance to this fungicide class do not make sequential applications of LUNA PRIVILEGE GREENHOUSE or any Group 7 containing fungicide before rotating with a fungicide from a different Group. DO NOT apply LUNA PRIVILEGE GREENHOUSE in tank mixture with other pesticides, fertilizers or adjuvants. Refer to RESISTANCE MANAGEMENT RECOMMENDATIONS section for details on multi-spray crops.</p> <p>LUNA PRIVILEGE GREENHOUSE may be applied up to the day of harvest.</p>		

<p>CROP SUBGROUP 8-09B: GREENHOUSE PEPPER/EGGPLANTS: Foliar Application [African Eggplant, Eggplant, Martynia, Okra, Pea Eggplant, Pepino, Pepper (bell and nonbell), Roselle, Scarlet Eggplant, and cultivars, varieties and hybrids of these commodities]</p>		
Disease Control	Rate mL/hectare	Use Information
<p>Grey mould (fruit) Botrytis leaf blight Botrytis stem canker (<i>Botrytis cinerea</i>)</p>	<p>500</p>	<p>Use a minimum water volume for thorough coverage of plant foliage. Suggested spray volumes by crop height: 1.7 m - 1000 L/Ha spray volume 3.4 m - 1500 - 2000 L/Ha spray volume</p> <p>Begin fungicide applications preventatively.</p>

		<p>Maximum of two applications per cropping cycle with a minimum re-application interval of 6 weeks. Rotate with non-Group 7 fungicides that control <i>Botrytis cinerea</i> 7 to 12-days later.</p> <p>Do not make a second application of LUNA PRIVILEGE GREENHOUSE within 6 weeks of the initial application.</p>
Powdery mildew (<i>Leveillula taurica</i>)	100	<p>Use a minimum water volume for thorough coverage of plant foliage. Suggested spray volumes by crop height: 1.7 m - 1000 L/Ha spray volume 3.4 m - 1500 - 2000 L/Ha spray volume</p> <p>Begin fungicide applications preventatively. Maximum of three applications per cropping cycle with a minimum re-application interval of 6 weeks. Rotate with non-Group 7 fungicides that control Powdery mildew 7 to 10-days later.</p> <p>Do not make a subsequent application of LUNA PRIVILEGE GREENHOUSE within 6 weeks of the preceding application.</p>
<p>CAUTION: Not all varieties have been tested for phytotoxicity under all conditions. The user should test a small number of plants for sensitivity before considering application to the entire crop. When testing a small number of plants, wait at least one week after application to determine sensitivity. Do not apply to the crop if any negative or adverse crop response is observed, or if a test for sensitivity was not conducted.</p> <p>Restrictions: Do not apply more than 500 g of fluopyram per hectare per season. To limit the potential for development of disease resistance to this fungicide class do not make sequential applications of LUNA PRIVILEGE GREENHOUSE or any Group 7 containing fungicide before rotating with a fungicide from a different Group. DO NOT apply LUNA PRIVILEGE GREENHOUSE in tank mixture with other pesticides, fertilizers or adjuvants. Refer to RESISTANCE MANAGEMENT RECOMMENDATIONS section for details on multi-spray crops.</p> <p>LUNA PRIVILEGE GREENHOUSE may be applied up to the day of harvest.</p>		

GREENHOUSE TOMATO: Foliar Broadcast Application		
Disease Control	Rate mL/hectare	Use Information
Grey mould (fruit) Botrytis leaf blight Botrytis stem canker (<i>Botrytis cinerea</i>)	500	<p>Use a minimum water volume for thorough coverage of plant foliage.</p> <p>Suggested spray volumes by crop height:</p> <p>1.2 m - 1000 L/Ha spray volume 2.7 m - 1000 - 1500 L/Ha spray volume</p> <p>Begin fungicide applications preventatively. Maximum of two applications per cropping cycle with a minimum re-application interval of 6 weeks. Rotate with non-Group 7 fungicides that control <i>Botrytis cinerea</i> 7 to 12-days later.</p> <p>Do not make a subsequent application of LUNA PRIVILEGE GREENHOUSE within 6 weeks of the preceding application.</p>
Early blight (<i>Alternaria solani</i>)	150-300	<p>Use a minimum water volume for thorough coverage of plant foliage.</p> <p>Suggested spray volumes by crop height:</p> <p>1.2 m - 1000 L/Ha spray volume 2.7 m - 1000 - 1500 L/Ha spray volume</p> <p>Begin fungicide applications preventatively. Maximum of three applications per cropping cycle with a minimum re-application interval of 6 weeks. Rotate with non-Group 7 fungicides that control Early blight 7 to 12-days later.</p> <p>Utilize the high rate when disease pressure is high or when agronomic or weather conditions are conducive to severe disease development.</p> <p>Do not make a subsequent application of LUNA PRIVILEGE GREENHOUSE within 6 weeks of the preceding application.</p>
Powdery mildew (<i>Leveillula taurica</i>)	100	<p>Use a minimum water volume for thorough coverage of plant foliage.</p> <p>Suggested spray volumes by crop height:</p> <p>1.2 m - 1000 L/Ha spray volume 2.7 m - 1000 - 1500 L/Ha spray volume</p> <p>Begin fungicide applications preventatively. Maximum of three applications per cropping cycle with a minimum re-application interval of 6 weeks. Rotate with non-Group 7 fungicides that control Powdery mildew 7 to 14-days later.</p>

		Do not make a second application of LUNA PRIVILEGE GREENHOUSE within 6 weeks of the initial application.
<p>CAUTION: Not all varieties have been tested for phytotoxicity under all conditions. The user should test a small number of plants for sensitivity before considering application to the entire crop. When testing a small number of plants, wait at least one week after application to determine sensitivity. Do not apply to the crop if any negative or adverse crop response is observed, or if a test for sensitivity was not conducted.</p> <p>Restrictions: Do not apply more than 500 g of fluopyram per hectare per season. To limit the potential for development of disease resistance to this fungicide class do not make sequential applications of LUNA PRIVILEGE GREENHOUSE or any Group 7 containing fungicide before rotating with a fungicide from a different Group. DO NOT apply LUNA PRIVILEGE GREENHOUSE in tank mixture with other pesticides, fertilizers or adjuvants. Refer to RESISTANCE MANAGEMENT RECOMMENDATIONS section for details on multi-spray crops.</p> <p>LUNA PRIVILEGE GREENHOUSE may be applied up to the day of harvest.</p>		

GREENHOUSE TOMATO: Drip/Drench Application		
Disease Control	Rate mL/hectare	Use Information
Grey mould (fruit) Botrytis leaf blight Botrytis stem canker (<i>Botrytis cinerea</i>)	40 ml product per 1000 plants	<p>A maximum of two applications made preventatively with a minimum re-application interval of 6 weeks from occurrence of flowering on the 9th truss provided the maximum rate per cropping cycle of 1.0 L/ha (500 g of fluopyram per hectare) is not exceeded.</p> <p>Apply the required amount in 500 L/ha to 1500 L/ha</p> <p>Do not apply prior to flowering on the 9th truss or before March 1st.</p> <p>Apply as part of a normal irrigation cycle, late in the day on last watering.</p> <p>Avoid application when conditions favour rapid growth.</p> <p>Do not make a second application of LUNA PRIVILEGE GREENHOUSE within 6 weeks of the initial application.</p> <p>Do not apply this product through any other type of irrigation system.</p>
<p>CAUTION: Not all varieties have been tested for phytotoxicity under all conditions, and ungrafted small fruited varieties (with small and thinner leaves) in particular have shown sensitivity to LUNA PRIVILEGE GREENHOUSE. The user should test a small number of plants for sensitivity before considering application to the entire crop. When testing a small number of plants, wait at least one week after application to determine sensitivity. Do not apply to the crop if any negative or adverse crop response is observed, or if a test for sensitivity was not conducted.</p> <p>Restrictions: Do not apply more than 500 g of fluopyram per hectare per season. To limit the potential for development of disease resistance to this fungicide class do not make sequential applications of LUNA PRIVILEGE GREENHOUSE or any Group 7 containing fungicide before rotating with a fungicide from a different</p>		

Group. A maximum of 2 sequential applications are permitted when applied in mixture with a non-Group 7 fungicide. Refer to RESISTANCE MANAGEMENT RECOMMENDATIONS section for details on multi-spray crops.

LUNA PRIVILEGE GREENHOUSE may be applied up to the day of harvest.

GREENHOUSE LETTUCE: Drip/Drench Application		
Disease Control	Rate mL/hectare	Use Information
Botrytis rot (<i>Botrytis cinerea</i>)	500	<p>First application can be made one week after transplanting at the earliest crop stage BBCH 12 (2nd leaf developed) and can be repeated once with a minimum interval of 6 weeks until the latest crop stage BBCH 49 (final head size for harvest is reached).</p> <p>Apply the required amount in 500 L/ha to 1500 L/ha.</p> <p>Do not make a second application of LUNA PRIVILEGE GREENHOUSE within 6 weeks of the initial application.</p> <p>Do not apply this product through any other type of irrigation system.</p>
<p>CAUTION: Not all varieties have been tested for phytotoxicity under all conditions. The user should test a small number of plants for sensitivity before considering application to the entire crop. When testing a small number of plants, wait at least one week after application to determine sensitivity. Do not apply to the crop if any negative or adverse crop response is observed, or if a test for sensitivity was not conducted.</p> <p>Restrictions: Do not apply more than 500 g of fluopyram per hectare per season. Do not apply LUNA PRIVILEGE GREENHOUSE within 7 days of harvest. To limit the potential for development of disease resistance to this fungicide class do not make sequential applications of LUNA PRIVILEGE GREENHOUSE or any Group 7 containing fungicide before rotating with a fungicide from a different Group. A maximum of 2 sequential applications are permitted when applied in mixture with a non-Group 7 fungicide. Refer to RESISTANCE MANAGEMENT RECOMMENDATIONS section for details on multi-spray crops.</p>		

MINOR USES:

NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS (BELOW): The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Bayer CropScience Inc. under the User Requested Minor Use Label Expansion program. For these uses, Bayer CropScience Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

GREENHOUSE STRAWBERRIES: Foliar Application

Disease Control	Rate mL/hectare	Use Information
Grey mould (<i>Botrytis</i> spp.)	500	Use a water volume for thorough coverage of plant foliage. Use a minimum water volume of 650 L/ha. Begin fungicide applications preventatively. Maximum of two applications per year with a minimum re-application interval of 6 weeks. Rotate with non-Group 7 fungicides that control listed diseases 7 to 10-days later. Do not make a second application of LUNA PRIVILEGE GREENHOUSE within 6 weeks of the initial application.
Powdery mildew (<i>Podosphaera aphanis</i> formerly <i>Sphaerotheca macularis</i>)		

CAUTION: Not all varieties have been tested for phytotoxicity under all conditions. The user should test a small number of plants for sensitivity before considering application to the entire crop. When testing a small number of plants, wait at least one week after application to determine sensitivity. Do not apply to the crop if any negative or adverse crop response is observed, or if a test for sensitivity as not conducted.

Restrictions: Do not apply more than 500 g of fluopyram per hectare per year. To limit the potential for development of disease resistance to this fungicide class do not make sequential applications of LUNA PRIVILEGE GREENHOUSE or any Group 7 containing fungicide before rotating with a fungicide from a different Group. DO NOT apply LUNA PRIVILEGE GREENHOUSE in tank mixture with other pesticides, fertilizers or adjuvants. Refer to RESISTANCE MANAGEMENT RECOMMENDATIONS section for details on multi-spray crops.

Preharvest Interval: The harvest of strawberries is permitted on the day of application.

MIXING INSTRUCTIONS:

LUNA PRIVILEGE GREENHOUSE must be applied with properly calibrated, clean equipment.

Prior to adding LUNA PRIVILEGE GREENHOUSE to the spray tank, ensure that the spray tank is thoroughly clean:

- Add one-half of the required amount of water to the spray or mixing tank and start agitation.
- Add the required quantity of LUNA PRIVILEGE GREENHOUSE to the water and complete filling with water to the required total volume.
- Maintain agitation throughout mixing and spraying.

RESISTANCE MANAGEMENT RECOMMENDATIONS:

For resistance management, LUNA PRIVILEGE GREENHOUSE contains a Group 7 fungicide. Any fungal population may contain individuals naturally resistant to LUNA PRIVILEGE GREENHOUSE and other Group 7 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

- Where possible, rotate the use of LUNA PRIVILEGE GREENHOUSE or other Group 7 fungicides with different groups that control the same pathogens.
- For Drip/Drench applications, use tank mixtures with fungicides from a different group that is effective on the target pathogen when such use is permitted.
- Avoid consecutive applications of LUNA PRIVILEGE GREENHOUSE or other Group 7 fungicides in a season. Alternate applications with fungicides having a different mode of action.
- For multi-spray crops: When applied alone, the total number of Group 7 fungicide applications should not exceed one third of the total number of fungicide applications in a season. When used in a mixture for Drip/Drench application, the total number of sprays containing Group 7 fungicides should not exceed half of the total number of fungicide applications in a season. When both solo and mixture applications are used in a program, no more than half of the total number of fungicide applications should contain Group 7 fungicides.
- Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications.
- Monitor treated fungal populations for resistance development. Notify Envu Canada if reduced sensitivity of the pathogen to LUNA PRIVILEGE GREENHOUSE is suspected.
- If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and disease problems in your area.
- For further information and to report suspected resistance, contact an Envu Canada representative at 1-800-331-2867 or at www.ca.envu.com.

Always read and follow label directions.
Luna Privilege™ is a trademark of the Bayer Group.
Bayer CropScience Inc. is a member of CropLife Canada.

230130