

2020-1550  
2020-08-04

**GROUP**

**9**

**HERBICIDE**

# **POWERMAX Liquid Herbicide**

**SOLUTION**

**AGRICULTURAL and INDUSTRIAL**

**CAUTION**



**POISON**

**WARNING - EYE AND SKIN IRRITANT**

**REGISTRATION NO. 32209 PEST CONTROL PRODUCTS ACT**

**ACTIVE INGREDIENT:** Glyphosate, 540 grams acid equivalent per litre, present as potassium salt.

Water Soluble Herbicide for non-selective weed control

**READ THE LABEL AND ATTACHED BROCHURE BEFORE USING.**

**NET CONTENTS: LITRES**

**BAYER CROPSCIENCE INC  
Suite 200, 160 Quarry Park Blvd SE  
Calgary, Alberta T2C 3G3  
1-888-283-6847  
[www.cropscience.bayer.ca](http://www.cropscience.bayer.ca)**

2020-1550  
2020-08-04

### **PRECAUTIONS**

KEEP OUT OF REACH OF CHILDREN.

HARMFUL IF SWALLOWED.

HARMFUL IF INHALED.

CAUSES EYE AND SKIN IRRITATION.

Avoid contact with eyes, skin or clothing.

Avoid inhaling spray mist.

Wear a long-sleeved shirt and long pants during mixing, loading, application, clean-up and repair. In addition, wear goggles or a face shield and chemical-resistant gloves during mixing and loading, clean-up and repair.

The restricted entry interval is 12 hours after application for all agricultural uses.

### **FIRST AID**

**If swallowed:** call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

### **TOXICOLOGICAL INFORMATION**

Treat symptomatically. This product contains a petroleum distillate. Vomiting may cause aspiration pneumonia.

### **ENVIRONMENTAL HAZARDS**

2020-1550  
2020-08-04

- **TOXIC** to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.
- To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil or clay.
- Avoid application when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

### **PHYSICAL OR CHEMICAL HAZARDS**

Spray solutions of this product should be mixed, stored and applied only in stainless steel, aluminum, fiberglass, plastic and plastic-lined steel containers. **DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.** This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

**In case of an emergency involving this product**, call Bayer CropScience collect, day or night:

Accident/Spills/Medical Emergency ..... 1-800-334-7577

**Read NOTICE before buying or using. If NOTICE terms are not acceptable, return at once unopened.**

For additional information on this or other Bayer CropScience agricultural products, call the Product Support Line at: 1-888-283-6847.

### **STORAGE**

Avoid contamination of seed, feed, and foodstuffs.  
Soak up small amounts of spill with absorbent clays.

### **DISPOSAL**

#### **RECYCLABLE CONTAINERS:**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or

2020-1550  
2020-08-04

municipality for the location of the nearest collection site. Before taking the container to the collection site:

- 1) Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
- 2) Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

**RETURNABLE CONTAINERS:**

Do not reuse container for any other purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

2020-1550  
2020-08-04

**REFILLABLE CONTAINERS:**

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on the disposal of unused, unwanted product, contact the manufacturer and the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for the clean-up of spills.

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

POWERMAX and Roundup Ready are registered trademarks of Bayer Group. Used under license. ©2020 Bayer Group. All rights reserved.

2020-1550  
2020-08-04

**GROUP**

**9**

**HERBICIDE**

# **POWERMAX HERBICIDE**

**SOLUTION**

**AGRICULTURAL and INDUSTRIAL**

**CAUTION**



**POISON**

**WARNING - EYE AND SKIN IRRITANT**

**REGISTRATION NO. 32209 PEST CONTROL PRODUCTS ACT**

**ACTIVE INGREDIENT:** Glyphosate, 540 grams acid equivalent per litre, present as potassium salt.

Water Soluble Herbicide for non-selective weed control

**READ THE LABEL BEFORE USING.**

**BAYER CROPSCIENCE INC**  
**Suite 200, 160 Quarry Park Blvd SE**  
**Calgary, Alberta T2C 3G3**  
**1-888-283-6847**  
**[www.cropscience.bayer.ca](http://www.cropscience.bayer.ca)**

(FRANÇAIS AU VERSO)

## CONTENTS

Page(s)

### **1.0 PRODUCT DESCRIPTION**

### **2.0 EMERGENCY NUMBERS**

#### **2.1 Information**

### **3.0 PRECAUTIONS**

#### **3.1 First Aid**

#### **3.2 Toxicological Information**

#### **3.3 Environmental Hazards**

#### **3.4 Physical or Chemical Hazards**

#### **3.5 Storage**

#### **3.6 Disposal and Decontamination**

## **DIRECTIONS FOR USE**

### **4.0 GENERAL INFORMATION**

### **5.0 MIXING AND APPLICATION**

#### **5.1 Precautions**

#### **5.2 Mixing and Application Equipment**

#### **5.3 Buffer Zones**

### **6.0 WEEDS CONTROLLED**

#### **6.1 Annual Weeds**

#### **6.2 Perennial Weeds**

#### **6.3 Woody Brush and Trees**

## **CROPLAND USES**

### **7.0 ANNUAL WEED CONTROL**

#### **7.1 Annual Weed Control with POWERMAX HERBICIDE**

#### **7.2 Annual Weed Control with POWERMAX HERBICIDE Tank Mixtures**

#### **7.3 Surfactant Information**

#### **7.4 Additional Important Information for Annual Weed Control**

#### **7.5 Weed Control in TruFlex™ Roundup Ready® Canola Varieties**

##### **7.5.1 TruFlex Roundup Ready Hybrid Canola Seed Production**

#### **7.6 Weed Control in Roundup Ready® Canola Varieties**

##### **7.6.1 Tank Mixtures**

##### **7.6.2 Roundup Ready® Hybrid Canola Seed Production**

#### **7.7 Weed Control in Roundup Ready2 Yield® or Roundup Ready Soybean Varieties**

##### **7.7.1 Weed Control in Roundup Ready2 Yield Soybean Varieties**



2020-1550

2020-08-04

- 9.8 Forage Seed Production
- 9.9 Preharvest Control of Quackgrass, Canada Thistle, Milkweed, Toadflax and Dandelion; Season-Long Control of Perennial Sow Thistle, and Harvest Management
  - 9.9.1 Guidelines for Timing of Preharvest Applications
  - 9.9.2 Preharvest Aerial Application
- 9.10 Tree Plantings
- 9.11 Tree, Vine, Berry and Other Crops
- 9.12 Selective Equipment
  
- 10.0 **NON-CROPLAND USES: INDUSTRIAL, RIGHTS-OF-WAY, RECREATIONAL, AND PUBLIC AREAS**
- 10.1 Weed Control in Non-Cropland Areas with POWERMAX HERBICIDE
- 10.2 Application Information for Non-Cropland Uses
  - 10.2.1 Ground Applications for all Non-Cropland Uses
  - 10.2.2 Purple Loosestrife Control
- 10.3 Selective Equipment for Non-Cropland Uses
- 10.4 Turf Grass
- 10.5 Injection Applications – for all Non-Cropland Uses
- 10.6 Cut Stump Application

# **POWERMAX HERBICIDE**

## **1.0 PRODUCT DESCRIPTION**

**Water soluble herbicide for non-selective weed control in CROPLAND SYSTEMS AND IN NON-CROPLAND AREAS.**

### **CROPLAND USES INCLUDE:**

In cropping systems before planting of all crops; in minimum tillage systems; postemergent in Roundup Ready 2™ Xtend soybeans, Roundup Ready® canola, soybeans and corn; preharvest applications in wheat, barley, oats, canola (rapeseed), flax (including low linolenic acid varieties), peas, lentils, dry beans, soybeans, chickpeas, dried lupins, dried fava beans and forages; in pasture renovation; in forage, legume and grass establishments; in tree crops including apple, pear, cherry, plum, peach, apricot, filbert, hazelnut, walnut, chestnut, Japanese heartnut; in grapes, cranberries, blueberries and strawberry; in sugar beets; in asparagus; in North American ginseng; in tree plantings; and grasses for seed production.

### **NON-CROPLAND USES INCLUDE:**

Industrial; recreational, rights-of-way, and public areas; turf grass renovation.

Not for relabelling or repackaging.

POWERMAX HERBICIDE and Roundup Ready are registered trademarks of Bayer Group. Used under license. ©2020 Bayer Group. All rights reserved.

## **2.0 EMERGENCY NUMBERS**

**In case of an emergency involving this product, call Bayer CropScience collect, day or night:**

Accident/Spills/Medical Emergency ..... 1-800-334-7577

**Read NOTICE before buying or using. If NOTICE terms are not acceptable, return at once unopened.**

2020-1550  
2020-08-04

## 2.1 INFORMATION

For additional information on this or other Bayer CropScience agricultural products, call the Product Support Line at: 1-888-283-6847.

## 3.0 PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.  
HARMFUL IF SWALLOWED.  
HARMFUL IF INHALED.  
CAUSES EYE AND SKIN IRRITATION.  
Avoid contact with eyes, skin or clothing.  
Avoid inhaling spray mist.

Wear a long-sleeved shirt and long pants during mixing, loading, application, clean-up and repair. In addition, wear goggles or a face shield and chemical-resistant gloves during mixing and loading, clean-up and repair.

The restricted entry interval is 12 hours after application for all agricultural uses.

## 3.1 FIRST AID

**If swallowed:** call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

## 3.2 TOXICOLOGICAL INFORMATION

2020-1550  
2020-08-04

Treat symptomatically. This product contains a petroleum distillate. Vomiting may cause aspiration pneumonia.

### **3.3 ENVIRONMENTAL HAZARDS**

- **TOXIC** to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.
- To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil or clay.
- Avoid application when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

### **3.4 PHYSICAL OR CHEMICAL HAZARDS**

Spray solutions of this product should be mixed, stored and applied only in stainless steel, aluminum, fiberglass, plastic and plastic-lined steel containers. **DO NOT MIX, STORE OR APPLY THIS PRODUCT OR SPRAY SOLUTIONS OF THIS PRODUCT IN GALVANIZED STEEL OR UNLINED STEEL (EXCEPT STAINLESS STEEL) CONTAINERS OR SPRAY TANKS.** This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

### **3.5 STORAGE**

Avoid contamination of seed, feed, and foodstuffs.  
Soak up small amounts of spill with absorbent clays.

### **3.6 DISPOSAL AND DECONTAMINATION**

#### **RECYCLABLE CONTAINERS:**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or

2020-1550  
2020-08-04

municipality for the location of the nearest collection site. Before taking the container to the collection site:

- 1) Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
- 2) Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

#### RETURNABLE CONTAINERS:

Do not reuse container for any other purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

#### REFILLABLE CONTAINERS:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on the disposal of unused, unwanted product, contact the manufacturer and the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for the clean-up of spills.

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

## **DIRECTIONS FOR USE**

### **4.0 GENERAL INFORMATION**

Do not apply this product using aerial spray equipment except under conditions as specified within this booklet.

Glyphosate is not to be applied using hand-wicking or hand-daubing methods.

Observe buffer zones specified in section 5.3.

POWERMAX HERBICIDE, a water soluble liquid, mixes readily with water for application as a foliage spray for the control or destruction of most herbaceous plants. It may be applied through most standard industrial or field type sprayers after dilution and thorough mixing with water in accordance with the booklet instructions.

2020-1550

2020-08-04

This herbicide moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days but on most perennial weeds may not occur until 7 to 10 days. Extremely cool or cloudy weather at treatment time may slow down activity of this product and delay visual effects of control. Visible effects are a gradual wilting and yellowing of the plant which advances to complete browning of above ground growth and deterioration of underground plant parts.

Delay application until vegetation has emerged to the stages described for control of such vegetation under the “**Annual and Perennial Weed Control**” (section 7.0 and 8.0) to provide adequate leaf surface to receive the spray. Unemerged plants arising from underground rhizomes or root stocks of perennials will not be affected by the spray and will continue to grow. For this reason best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity.

Always use the higher rate of this product per hectare within the recommended range when weed growth is heavy or dense, or weeds are growing in an undisturbed (non-cultivated) area.

Do not treat weeds under poor growing conditions such as drought stress, disease or insect damage, as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust.

This product does not provide residual weed control. For subsequent residual weed control follow a label approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

Rainfall occurring within 60 minutes of treatment may result in reduced weed control. Heavy rainfall immediately after application may wash the chemical off the foliage and a repeat treatment may be required. Do not apply if rainfall is forecast for the time of application.

Do not mix with any surfactant, pesticide, herbicide oils or any other material other than water unless specified in this booklet. For best results, spray coverage should be uniform and complete. Do not spray weed foliage to the point of run-off.

**As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests**

**DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.**

**RESISTANCE-MANAGEMENT RECOMMENDATIONS**

For resistance management, POWERMAX HERBICIDE is a Group 9 herbicide. Any weed population may contain or develop plants naturally resistant to POWERMAX HERBICIDE and other Group 9 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of POWERMAX HERBICIDE or other Group 9 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Bayer CropScience at 1-888-283-6847 or at [www.cropscience.bayer.ca](http://www.cropscience.bayer.ca)

## **5.0 MIXING AND APPLICATION**

### **5.1 PRECAUTIONS**

**ATTENTION: AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.**

2020-1550  
2020-08-04

**APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES.**

**DO NOT USE IN GREENHOUSES. REDUCED RESULTS MAY OCCUR IF WATER CONTAINING SOIL IS USED, SUCH AS WATER FROM PONDS AND UNLINED DITCHES.**

Clean sprayers and parts immediately after using this product by thoroughly flushing with water. Do not contaminate water sources by disposal of wastes or cleaning of equipment.

DO NOT use human flaggers.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature, application equipment and sprayer settings.

**NOTE:** Use of this product in any manner not consistent with this booklet may result in injury to persons, animals or crops, or other unintended consequences. Keep container closed to prevent spills and contamination.

2020-1550

2020-08-04

## 5.2 MIXING AND APPLICATION EQUIPMENT

### MIXING WITH WATER

For ground or industrial type sprayers, fill the spray tank with one-half the required amount of water. Add the proper amount of herbicide, see “**Weed Control**” (sections 7.1 and 8.1) and mix well before adding the remaining portion of water. Placing the filling hose below the surface of the liquid solution will prevent excessive foaming. Removing hose from tank immediately will avoid back siphoning into water source. Use of mechanical agitators may cause excessive foaming. Bypass lines should terminate at the bottom of the tank.

For use in knapsack sprayers, it is suggested that the proper amount of this herbicide be mixed with water in a larger container. Fill sprayer with the mixed solution.

### TANK MIXING PROCEDURE

**The following steps should be followed when adding tank mix partners, using an herbicide loading system or adding product directly into the tank:**

1. Fill spray tank 3/4 full of water.
2. Start agitation and run for entire mixing and spraying operation.
3. Add required amount of the tank mix partner.
4. Flush herbicide loading tank and herbicide containers with water.
5. If using an herbicide loading system - ensure that the loading tank and lines to the pump are empty and flushed out with water before adding tank mix partner.
6. Add required amount of POWERMAX HERBICIDE.
7. Flush herbicide loading tank and herbicide containers with water.
8. If using an herbicide loading system - ensure that the loading tank and lines to the pump are flushed with water and empty before starting spray operation.

Always start and end the mixing and spraying operation with a clean system.

### APPLICATION EQUIPMENT

#### BOOM EQUIPMENT

**For control of perennial weeds and woody brush and trees listed on this booklet using conventional boom equipment** – apply this product in 50 to 300 litres of clean water per hectare as a broadcast spray using no more than 275 kPa pressure. See “**Weed Control**” (sections 7.1 and 8.1) for rates to control specific weeds.

**For control of annual weeds listed on this booklet using conventional boom equipment** – Apply this product in 50 to 100 litres of clean water per hectare as a

2020-1550

2020-08-04

broadcast spray, except as otherwise stated on this label using no more than 275 kPa pressure. See “**Weed Control**” (sections 7.1 and 8.1) for rates to control specific weeds.

### **HAND HELD AND HIGH VOLUME EQUIPMENT (use coarse sprays only)**

**For control of weeds and woody brush and trees listed in the “Weed Control” section (6.0) of this label using knapsack sprayers or high volume spraying equipment utilizing handguns or other suitable nozzle arrangements** – Unless otherwise specified, make a 0.67 percent solution of this product in water (0.67 litres of this product in 100 litres of water) and apply to foliage of vegetation to be controlled. For best results, use a 1.34 percent solution (1.34 litres of this product in 100 litres of water) on harder to control perennials such as field bindweed, hemp dogbane, milkweed and Canada thistle.

Applications should be made on a spray-to-wet basis. Spray coverage should be uniform and complete. Do not spray to point of run-off. Handgun applications should be properly directed to avoid spraying desirable plants.

### **SELECTIVE EQUIPMENT**

Selective equipment such as **WIPER** and **ROLLER** applicators can be used for weed control in soy and dry beans, orchards, vineyards, cranberries, strawberries and non-crop areas. For information regarding use of this product with selective equipment, refer to “**Selective Equipment**” (section 9.12).

### **AERIAL EQUIPMENT**

Aerial application can only be used for weed control in preharvest situations. Refer to sections 5.3 and 9.9.2 for application information.

#### **Directions for use**

Apply only by fixed-wing or rotary aircraft which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label. Ensure that the maximum boom width does not exceed 65% of the wing span. Nozzle type, size and orientation must be configured to deliver a droplet size VMD in the coarse (400-600 microns) or very coarse (600-1000) range.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate(s) recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

2020-1550

2020-08-04

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices, or equivalent electronic positioning systems (GPS). The use of spotter planes is recommended.

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. **PROLONGED EXPOSURE OF THIS PRODUCT TO UNCOATED STEEL SURFACES MAY RESULT IN CORROSION AND POSSIBLE FAILURE OF THE PART. LANDING GEAR IS MOST SUSCEPTIBLE.** The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38412 may prevent corrosion.

### **Use Precautions**

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the *National Aerial Pesticide Application Manual*, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Do not apply to any body of water. Avoid drifting of spray onto any body of water or other non-target areas. Specified buffer zones should be observed.

Coarse sprays are less likely to drift, therefore, avoid combinations of pressure and nozzle type that will result in fine particles (mist). Do not apply during periods of dead calm or when wind velocity and direction pose a risk of spray drift. Do not spray when the wind is blowing towards a nearby sensitive crop, garden, terrestrial habitat (such as shelter-belt) or aquatic habitat.

Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

### **Operator Precautions**

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more

2020-1550  
2020-08-04

stringent label precautions in cases where the operator precautions exceed generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

### **Product Specific Precautions**

Read and understand the entire label before opening this product. If you have questions, call the Bayer CropScience Custom Care Line at 1-888-283-6847 or obtain technical advice from the distributor or your provincial agricultural representative.

Application of this product must meet and/or conform to the following:

Volume: Apply the recommended rate in a minimum spray volume of 30-100 litres per hectare.

## **5.3 BUFFER ZONES**

i) Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. Boom height must be 60 cm or less above the crop or ground.

Airblast or mist blower application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** direct spray above plants to be treated. **DO NOT** apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side. For airblast applications, turn off outward pointing nozzles at row ends and outer rows.

Aerial application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 km/h at flying height at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. To reduce drift caused by turbulent wingtip vortices, the nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotorspan.

ii) **Buffer zones:**

2020-1550

2020-08-04

Use of the following spray methods or equipment **DO NOT** require a buffer zone: hand-held or backpack sprayer and spot treatment, inter-row hooded sprayer, low-clearance hooded or shielded sprayers that ensure spray drift does not come in contact with orchard crop fruit or foliage, soil drench and soil incorporation.

For application to rights-of-way and for forestry uses, buffer zones for protection of sensitive terrestrial habitats are not required; however, the best available application strategies which minimize off-site drift, including meteorological conditions (for example, wind direction, low wind speed) and spray equipment (for example, coarse droplet sizes, minimizing height above canopy), should be used. Applicators must, however, observe the specified buffer zones for protection of sensitive aquatic habitats.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands) and sensitive aquatic habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, wetlands and estuarine/marine water bodies).

Agricultural and non-cropland systems	Maximum number of applications	Buffer Zones (metres) Required for the Protection of:	
		Aquatic habitats	Terrestrial habitats
<b>Agricultural crop system and ground boom application method</b>			
Pre-seeding applications for rye, cranberry, filberts, hazelnut and all other crops. Established pasture and summer fallow. Ginseng new garden.	1	1	1
Ginseng - existing established garden, Canola – Roundup Ready hybrid for seed production	2	1	1
Filberts or hazelnut, sugar beets (glyphosate tolerant varieties)	4	1	1
Corn (glyphosate non-tolerant varieties including grain, silage and ornamental types), sugar beet (glyphosate non-tolerant varieties), strawberry, blueberry highbush and lowbush, walnut, chestnut, Japanese heartnut, Turf grass (prior to establishment or renovation)	2	1	2
Wheat, barley, oats, soybean (glyphosate non-tolerant varieties), corn-sweet (glyphosate tolerant varieties), canola (glyphosate non-tolerant varieties), peas, dry beans, flax (including low linoleic acid varieties), lentils, chickpea, lupin (dried), fava bean (dried), asparagus, corn (glyphosate tolerant varieties), forage grasses and legume including seed production	3	1	2
Canola (glyphosate tolerant varieties), soybean (glyphosate tolerant varieties)	4	1	2
Apple, apricot, cherry (sweet/sour), peaches, pears, plums, grapes	3	1	3
<b>Agricultural crop system and airblast application method (including mist blower)</b>			
Pasture	1	20	30
Turfgrass (Prior to establishment or renovation)	2	25	35

Agricultural and non-cropland systems		Maximum number of applications	Buffer Zones (metres) Required for the Protection of:	
			Aquatic habitats	Terrestrial habitats
<b>Non-cropland system and ground boom application method</b>				
Non-crop land and industrial uses: Industrial and rights of way areas, Recreational and public areas		3	1	3*
<b>Non-cropland system and airblast application method (including mist blower)</b>				
Non-crop land and industrial uses: Industrial and rights of way areas, Recreational and public areas		3	1	30*
<b>Agricultural crop system and aerial application method</b>	<b>Wing type</b>			
Rye, corn (glyphosate non-tolerant varieties), corn-sweet (glyphosate tolerant varieties), chickpea, lupin (dried), fava bean (dried), sugar beet (glyphosate non-tolerant varieties), all other crops for pre-seeding treatments only	Fixed and rotary wing	1	15	20
Canola (glyphosate tolerant varieties)	Fixed and rotary wing	3	20	40
Sugar beets (glyphosate tolerant varieties)	Fixed wing	2	20	30
	Rotary wing	2	15	30
Wheat, barley, oats, soybean (glyphosate non-tolerant varieties), canola (glyphosate non-tolerant varieties), peas, dry beans, flax (including low linoleic acid varieties), lentils	Fixed wing	2	20	35
	Rotary wing	2	20	30

Agricultural and non-cropland systems		Maximum number of applications	Buffer Zones (metres) Required for the Protection of:	
			Aquatic habitats	Terrestrial habitats
Forage grasses and legume including seed production	Fixed and rotary wing	1	20	40
Soybean (glyphosate tolerant varieties)	Fixed wing	3	20	45
	Rotary wing	3	20	40
Summer fallow	Fixed wing	1	20	45
	Rotary wing	1	20	40
Corn (glyphosate tolerant varieties)	Fixed wing	2	20	50
	Rotary wing	2	20	45
Pasture	Fixed wing	1	30	70
	Rotary wing	1	30	55
<b>Non-cropland system and aerial application method</b>				
Non-crop land and industrial uses: rights-of way areas only	Fixed wing	3	100	NR
	Rotary wing	3	60	NR

\* Buffer zones for the protection of terrestrial habitats are not required for use on rights-of-way including railroad ballast, rail and hydro rights-of-way, utility easements and roads.

NR = Not Required

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

2020-1550  
2020-08-04

The buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Buffer Zone Calculator on the Pest Management Regulatory Agency web site.

## **6.0 WEEDS CONTROLLED**

This product controls many annual and perennial grasses, broadleaf weeds, and woody brush and trees when applied as recommended and under conditions described. For information on how to control specific weeds including herbicide rate, refer to “**Annual Weed Control**” and “**Perennial Weed Control**” (sections 7.1 and 8.1). The following is a partial list of weeds controlled:

2020-1550

2020-08-04

## 6.1 ANNUAL WEEDS

### ANNUAL GRASSES

#### Barnyard Grass

*Echinochloa crusgalli*

#### Blue Grass (annual)

*Poa annua*

#### Crab Grass (large)

*Digitaria sanguinalis*

#### Crab Grass (smooth)

*Digitaria ischaemum*

#### Downy Brome Grass

*Bromus tectorum*

#### Fall Panicum

*Panicum dichotomiflorum*

#### Giant Foxtail

*Setaria faberii*

#### Green Foxtail

*Setaria viridis*

#### Persian Darnel

*Lolium persicum*

#### Volunteer Barley

*Hordeum spp.*

#### Volunteer Corn

*Zea mays*

#### Volunteer Wheat

*Triticum spp.*

#### Wild Oats

*Avena fatua*

#### Wild Proso Millet

*Panicum miliaceum*

#### Yellow Foxtail

*Setaria glauca*

### OTHER

#### Dodder

*Cuscuta spp.*

### ANNUAL BROADLEAF WEEDS

#### Chickweed

*Stellaria media*

#### Cleavers

*Galium aparine*

#### Cocklebur

*Xanthium strumarium*

#### Corn Spurry

*Spergula arvensis*

#### Cow Cockle

*Saponaria vaccaria*

#### Eastern Black Nightshade

*Solanum ptycanthum*

#### Fleabane (Canada)

*Erigeron canadensis*

#### Flixweed

*Descurainia sophia*

#### Green Smartweed

*Polygonum scabrum*

#### Hempnettle

*Galeopsis tetrahit*

#### Kochia

*Kochia scoparia*

#### Lady's-Thumb

*Polygonum persicaria*

#### Lamb's-quarters (common)

*Chenopodium album*

#### Narrow-leaved Hawk's Beard

*Crepis tectorum*

#### Narrow-leaved Vetch

*Vicia angustifolia*

#### Night-flowering Catchfly

*Silene noctiflora*

2020-1550

2020-08-04

**Pennsylvania Smartweed**

*Polygonum pensylvanicum*

**Prickly Lettuce**

*Lactuca scariola*

**Ragweed (common)**

*Ambrosia artemisiifolia*

**Redroot Pigweed**

*Amaranthus retroflexus*

**Round-Leaved Mallow**

*Malva pusilla*

**Russian Thistle**

*Salsola pestifer*

**Shepherd's Purse**

*Capsella bursa-pastoris*

**Smooth Pigweed**

*Amaranthus hybridus*

**Sowthistle (annual)**

*Sonchus oleraceus*

**Stinkweed**

*Thlaspi arvense*

**Storksbill**

*Erodium cicutarium*

**Velvetleaf**

*Abutilon theophrasti*

**Volunteer Canola (rapeseed)**

*Brassica spp.*

**Volunteer Flax**

*Linum spp.*

**Wild Buckwheat**

*Polygonum convolvulus*

**Wild Mustard**

*Sinapis arvensis*

**Wild Tomato**

*Solanum triflorum*

**6.2 PERENNIAL WEEDS**

**PERENNIAL GRASSES/SEDGES**

**Blue Grass (Canada)**

*Poa compressa*

**Blue Grass (Kentucky)**

*Poa pratensis*

**Brome Grass (smooth)**

*Bromus inermis*

**Cattail (common)**

*Typha latifolia*

**Cottongrass**

*Eriophorum chamissonis*

**Foxtail Barley**

*Hordeum jubatum*

**Quackgrass**

*Elytrigia repens*

**Wire-Stemmed Muhly**

*Muhlenbergia frondosa*

**Yellow Nutsedge**

*Cyperus esculentus*

**PERENNIAL BROADLEAVED WEEDS**

**Alfalfa**

*Medicago spp.*

**Curled Dock**

*Rumex crispus*

**Dandelion**

*Taraxacum officinale*

**Field Bindweed**

*Convolvulus arvensis*

**Hemp Dogbane**

*Apocynum cannabinum*

**Hoary Cress**

*Cardaria draba*

**Knotweed (Japanese)**

*Polygonum cuspidatum*

**Milkweed (common)**

*Asclepias syriaca*

2020-1550  
2020-08-04

**Poison Ivy**

*Rhus radicans*

**Purple Loosestrife**

*Lythrum salicaria*

**Sow Thistle (perennial)**

*Sonchus arvensis*

**Thistle (Canada)**

*Cirsium arvense*

**Toad Flax**

*Linaria vulgaris*

**Wormwood (Absinth)**

*Artemisia absinthium*

2020-1550

2020-08-04

### **6.3 WOODY BRUSH AND TREES**

**Alder**

*Alnus spp.*

**Birch**

*Betula spp.*

**Broadleaved meadowsweet**

*Spiraea latifolia*

**Rhododendron (Canadian)**

*Rhododendron canadense*

**Cedar**

*Thuja spp.*

**Cherry**

*Prunus spp.*

**Douglas Fir**

*Pseudotsuga spp.*

**Hemlock**

*Tsuga spp.*

**Maple**

*Acer spp.*

**Mountain-fly honeysuckle**

*Lonicera villosa*

**Pine**

*Pinus spp.*

**Poplar**

*Populus spp.*

**Raspberry/Salmonberry**

*Rubus spp.*

**Sheep laurel**

*Kalmia angustifolia*

**Snowberry (Western)**

*Symphoricarpos occidentalis*

**Sweet fern**

*Comptonia peregrina*

**Willow**

*Salix spp.*

**Withrod**

*Viburnum cassinoides*

### **7.0 ANNUAL WEED CONTROL**

The following tables provide rates and specific application instructions for control of the annual weeds listed.

2020-1550  
2020-08-04

**7.1 ANNUAL WEED CONTROL WITH POWERMAX HERBICIDE**  
DO NOT APPLY BY AIR.

<b>RATE (L/ha)</b>	<b>GROWTH STAGE</b>	<b>WEEDS CONTROLLED</b>	<b>COMMENTS (Apply in 50-100 L/ha water)</b>
0.5	Weeds up to 8 cm in height	Wild oats, green foxtail, volunteer barley, volunteer wheat  Non-Roundup Ready® volunteer canola (rapeseed), wild mustard, lady's-thumb, stinkweed	<ul style="list-style-type: none"> <li>• For wild oats apply at 1-3 leaf stage.</li> <li>• Add 350 mL of a surfactant registered for use such as Agral® 90, Ag Surf®, or Companion™</li> <li>• For heavy wild oat infestations use 0.67 L/ha rate.</li> </ul>
0.67	Weeds 8 cm to 15 cm in height	All annual grasses listed above.  All annual broadleaved weeds listed above plus flixweed* and kochia*	<ul style="list-style-type: none"> <li>• Add 350 mL of surfactant registered for use as listed above.</li> </ul> <p>* Suppression only. Refer to higher rates of this table or tank mix table (section 7.2) for control options.</p>
0.83 – 1.27	Weeds up to 15 cm in height	All annual grasses listed above plus downy brome, giant foxtail, and Persian darnel.  All annual broadleaved weeds listed above plus cleavers, lamb's-quarters, redroot pigweed, hempnettle, flixweed, Russian thistle, volunteer flax, common ragweed*, Canada fleabane*, wild buckwheat**, narrow-leaved hawk's beard***	<ul style="list-style-type: none"> <li>• No surfactant required.</li> <li>• For tank mix weed control options see section 7.2.</li> </ul> <p>* DO NOT use these rates on plants greater than 8 cm in height.</p> <p>** For 3-4 leaf stage use 1.27 L/ha rate.</p> <p>*** For weeds 8 cm to 15 cm in height use 1.27 L/ha rate.</p>
1.5	Weeds up to 15 cm in height	All annual grasses listed above plus crab grass and annual blue grass  All annual broadleaved weeds listed above plus kochia, prickly lettuce, shepherd's purse, annual	<ul style="list-style-type: none"> <li>• For additional annual broadleaved weed control options, refer to tank mix table (section 7.2).</li> </ul>

2020-1550  
2020-08-04

<b>RATE (L/ha)</b>	<b>GROWTH STAGE</b>	<b>WEEDS CONTROLLED</b>	<b>COMMENTS (Apply in 50-100 L/ha water)</b>
		sow thistle, and narrow- leaved vetch	

2020-1550  
2020-08-04

2.33	Weeds over 15 cm in height	All annual grasses and broadleaved weeds listed above	• For additional annual broadleaved weed control options, refer to tank mix table (section 7.2).
------	----------------------------	---	--

Agral is a registered trademark of Syngenta group company.  
Ag-Surf is a registered trademark of Interprovincial Cooperative Ltd.  
Companion is a trademark of Dow AgroSciences LLC.

**NOTE:** For spot treatment, 0.5 – 2.33 litres per hectare is approximately equivalent to 5 – 23 mL/100m<sup>2</sup>, respectively.

## 7.2 ANNUAL WEED CONTROL WITH POWERMAX HERBICIDE TANK MIXTURES FOR SUMMERFALLOW & MINIMUM TILLAGE SYSTEMS

TANK MIXTURES	RATE (L/ha)	WEEDS CONTROLLED♦	COMMENTS (Apply in 50-100 L/ha water)
POWERMAX Herbicide  +  Banvel® II	0.5 – 0.67   +  0.29	Volunteer cereals, wild oats, green foxtail  Non-Roundup Ready® volunteer canola (rapeseed), wild mustard, flixweed*, lamb's-quarters, lady's-thumb, stinkweed, kochia, Russian thistle, cow cockle, redroot pigweed**, wild buckwheat**	This tank mix is registered for <b>summerfallow use only</b> . Weeds should be less than 15 cm tall and actively growing for best results.  Use higher rate if weeds are beyond 8 cm in height.  * POWERMAX Herbicide applied at 0.67 L/ha rate only.  ** Suppression only. See other tank mixtures for control options.  Add 350 mL/ha of surfactant – see list in section 7.3.
POWERMAX Herbicide  +  Banvel® II	0.61 – 1.27   +  	Volunteer cereals, wild oats, green foxtail, downy brome, Persian darnel  Non-Roundup Ready® volunteer canola (rapeseed), wild mustard, flixweed, lamb's-quarters, lady's-	Use this tank mix prior to seeding in <b>wheat, barley, rye, oats, field corn only (do not apply to sweet corn)</b> .  Certain broadleaved crops such as lentils, peas, canola and flax can be injured by a pre-seeding application and so should not be

TANK MIXTURES	RATE (L/ha)	WEEDS CONTROLLED♦	COMMENTS (Apply in 50-100 L/ha water)
	0.31	thumb, stinkweed, kochia, Russian thistle, cow cockle, redroot pigweed, wild buckwheat*, smartweed	<p>planted to a field receiving this treatment.</p> <p>Annual grasses - apply any time between emergence and heading.</p> <p>Weeds should be less than 15 cm tall and actively growing for best results.</p> <p>The higher rate should be applied when weeds are under poor growing conditions such as drought.</p> <p>*1- to 4- leaf stage.</p>
POWERMAX Herbicide + Pardner®	0.5 – 0.67  + 1.25	Volunteer cereals, green foxtail, volunteer canola (rapeseed), wild mustard, lady’s-thumb, stinkweed, wild buckwheat*  Redroot pigweed**, kochia**, wild oats**	<p>This tank mix is registered <b>only for use in summerfallow, and prior to wheat, oats and barley in minimum tillage systems.</b></p> <p>Weeds should be less than 15 cm tall and actively growing for best results.</p> <p>Use higher rate if weeds are beyond 8 cm in height.</p> <p>* Use POWERMAX Herbicide at 0.67 L/ha rate only for wild buckwheat control.</p> <p>** 0.67 L/ha rate, suppression only. See other tank mixtures for control options.</p> <p>Add 350 mL/ha of surfactant – see list in section 7.3.</p>
POWERMAX Herbicide +	0.83 – 1.27	Volunteer cereals, wild oats, green foxtail, downy brome, giant foxtail, Persian darnel	<p>Weeds should be less than 15 cm tall and actively growing for best results.</p>

TANK MIXTURES	RATE (L/ha)	WEEDS CONTROLLED♦	COMMENTS (Apply in 50-100 L/ha water)
2,4-D <sup>A</sup>	+ 0.6 – 0.9 <sup>4</sup> or 1.2 – 1.5 <sup>5</sup>	<p>Volunteer canola, (rapeseed) (non-Roundup Ready), wild mustard, flixweed, redroot pigweed, lady's-thumb, stinkweed, kochia, lamb's-quarters, hempnettle, Russian thistle, volunteer flax, common ragweed*, Canada fleabane, wild buckwheat**, narrow-leaved hawk's beard***</p> <p>Volunteer Roundup Ready canola (1-4 leaf stage)<sup>4</sup>, bluebur<sup>4</sup>, burdock<sup>4</sup>, cocklebur<sup>4</sup>, common plantain<sup>4</sup>, daisy fleabane<sup>4</sup>, false flax<sup>4</sup>, false ragweed<sup>4</sup>, goat's beard<sup>4</sup>, mustards<sup>4</sup> (except dog and tansy), prickly lettuce<sup>4</sup>, ragweeds<sup>4</sup>, Russian pigweed<sup>4</sup>, shepherd's purse<sup>4</sup>, stinging nettle<sup>4</sup>, sweet clover<sup>4</sup>, thyme-leaved spurge<sup>4</sup>, wild radish<sup>4</sup>, wild sunflower<sup>4</sup></p> <p>Volunteer Roundup Ready canola (rapeseed) (4-6 leaf stage)<sup>5</sup>, annual sowthistle<sup>5</sup>, common chickweed<sup>5</sup>, common purslane<sup>5</sup>, dog and tansy mustard<sup>5</sup>, oak-leaved goosefoot<sup>5</sup>, common groundsel<sup>5</sup>, hairy galinsoga<sup>5</sup>, hawkweed<sup>5</sup>, heal-all<sup>5</sup>, knotweed<sup>5</sup>,</p>	<p>Use higher rate if weeds are beyond 8 cm in height.</p> <p>* DO NOT use these rates on plants greater than 8 cm in height.</p> <p>** For 3- to 4-leaf stage use 1.27 L/ha rate.</p> <p>*** For weeds 8 cm to 15 cm in height use 1.27 L/ha rate.</p> <p><sup>4</sup> 2,4-D at 0.6 – 0.9 L/ha (280 – 420 g ai/ha).</p> <p><sup>5</sup> 2,4-D at 1.2 – 1.5 L/ha (560 – 700 g ai/ha). Use a minimum of 80 L/ha water when using 2,4-D amine formulations at these rates.</p> <p>Use this tank mix prior to seeding or after seeding but before crop emergence in <b>wheat, winter wheat, barley and rye.</b></p> <p>No surfactant required.</p>

TANK MIXTURES	RATE (L/ha)	WEEDS CONTROLLED♦	COMMENTS (Apply in 50-100 L/ha water)
		peppergrass <sup>5</sup> , pineapple weed <sup>5</sup> , prostrate pigweed <sup>5</sup> , purslane <sup>5</sup> , sheep sorrel <sup>5</sup> , green smartweed <sup>5</sup> , tumble pigweed <sup>5</sup> , velvetleaf <sup>5</sup> , volunteer canola (rapeseed) <sup>5</sup>	
POWERMAX Herbicide  +  2,4-D <sup>B</sup>	0.5 – 0.67          +  1.2	Volunteer cereals, wild oats*, green foxtail*  Volunteer canola (rapeseed), wild mustard, flixweed, redroot pigweed, lady’s-thumb, stinkweed, kochia  Lamb’s-quarters**, Russian thistle**	This tank mix is registered for summerfallow use only. Weeds should be less than 15 cm tall and actively growing for best results.  Use higher rate if weeds are beyond 8 cm in height.  * Use POWERMAX Herbicide at 0.67 L/ha rate only for wild oat and green foxtail control.  ** Suppression only. See other tank mixtures for control options.  Add 350 mL/ha of surfactant – see list in section 7.3.
POWERMAX Herbicide  +  MCPA <sup>C</sup> 500 g/L formulation; if another formulation is used, adjust rate accordingly.	0.83 – 1.27       +  0.5 – 0.7 <sup>1</sup>  OR 0.5 – 1.0 <sup>2</sup>	Volunteer cereals, wild oats, green foxtail, downy brome, giant foxtail, Persian darnel  Volunteer canola (rapeseed) (non-Roundup Ready), wild mustard, flixweed, redroot pigweed, lady’s thumb, stinkweed, kochia, lamb’s quarters, hempnettle, Russian thistle, volunteer flax, common ragweed*, Canada fleabane, wild	Weeds should be less than 15 cm tall and actively growing for best results.  Use higher rate if weeds are beyond 8 cm in height.  * DO NOT use these rates on plants greater than 8 cm in height.  ** For 3- to 4-leaf stage use 1.27 L/ha rate.  *** For weeds 8 cm to 15 cm in height use 1.27 L/ha rate.

TANK MIXTURES	RATE (L/ha)	WEEDS CONTROLLED♦	COMMENTS (Apply in 50-100 L/ha water)
		buckwheat**, narrow-leaved hawk's beard***  Volunteer Roundup Ready canola (1-4 leaf stage) <sup>1,2</sup> , bluebur <sup>3</sup> , burdock <sup>3</sup> (before 4 leaf stage), false flax <sup>3</sup> , flixweed <sup>3</sup> , lamb's quarters <sup>3</sup> , mustards <sup>3</sup> (except dog and tansy), prickly lettuce <sup>3</sup> , ragweeds <sup>3</sup> , redroot pigweed <sup>3</sup> , Russian pigweed <sup>3</sup> , shepherd's purse <sup>3</sup> , stinkweed (field pennycress) <sup>3</sup> , vetch <sup>3</sup> , wild radish <sup>3</sup> , wild sunflower <sup>3</sup>	<sup>1</sup> MCPA amine at 0.5 – 0.7 L/ha (250 – 350 g ai/ha) prior to peas.  <sup>2</sup> MCPA at 0.5 – 1.0 L/ha (250 – 500 g ai/ha) prior to wheat, barley, oats, corn (field and sweet) <sup>C</sup> , rye and flax.  <sup>3</sup> MCPA at 0.7 – 1.0 L/ha (350 – 500 g ai/ha) only.  Use this tank mix prior to seeding in wheat, barley, rye, oats, corn (field and sweet) <sup>C</sup> , flax and field peas <sup>C</sup> .  No surfactant required.
POWERMAX Herbicide  +  Buctril® M Herbicide	0.83 – 1.27  +  0.5 – 1.0 <sup>1</sup>	Volunteer cereals, wild oats, green foxtail, downy brome, giant foxtail, Persian darnel.  Volunteer canola (rapeseed) (non-Roundup Ready), wild mustard, flixweed, redroot pigweed, lady's thumb, stinkweed, kochia, lamb's quarters, hempnettle, Russian thistle, volunteer flax, common ragweed*, Canada fleabane, wild buckwheat**, narrow-leaved hawk's beard***  Volunteer Roundup Ready Canola (1-4 leaf stage) <sup>1,2</sup>	Weeds should be less than 15 cm tall and actively growing for best results.  Use higher rate if weeds are beyond 8 cm in height.  * DO NOT use these rates on plants greater than 8 cm in height.  ** For 3- to 4-leaf stage use 1.27 L/ha rate.  *** For weeds 8 cm to 15 cm in height use 1.27 L/ha rate.  <sup>1</sup> Buctril M at 0.5 – 1.0 L/ha (280 – 560 g ai/ha) for all crops listed.  <sup>2</sup> Buctril M at 1.0 L/ha (560 g ai/ha only).

2020-1550  
2020-08-04

TANK MIXTURES	RATE (L/ha)	WEEDS CONTROLLED♦	COMMENTS (Apply in 50-100 L/ha water)
		<p>Seedlings up to the 4-leaf stage<sup>2</sup>: green smartweed, pale smartweed, lady's thumb, cow cockle, redroot pigweed, flixweed, bluebur, shepherd's purse, kochia<sup>3</sup>, Russian thistle<sup>3</sup>, scentless chamomile<sup>4</sup>, volunteer sunflower, night flowering catchfly, cocklebur, velvetleaf<sup>5</sup>, ball mustard, American nightshade</p> <p>Seedlings up to the 6-leaf stage<sup>2</sup>: wild tomato</p> <p>Seedlings up to the 8-leaf stage<sup>2</sup>: wild buckwheat, tartary buckwheat, common buckwheat, stinkweed, wild mustard, wormseed mustard, lamb's quarters, common ragweed, common groundsel</p> <p>Perennials (top growth)<sup>2</sup>: Canada thistle, perennial sowthistle</p>	<p><sup>3</sup> Spray before plants are 5 cm high.</p> <p><sup>4</sup> Spring annuals only.</p> <p><sup>5</sup> Spray before plants are 8 cm high.</p> <p>Use this tank mix prior to seeding in <b>wheat, barley, rye, oats, corn, flax, canary seed and seedling grasses (including brome grass, crested wheatgrass, intermediate wheat grass, slender wheatgrass, tall wheatgrass, Russian wild rye, timothy, orchard grass, creeping red fescue, meadow fescue, meadow foxtail, seedling tall fescue, seedling meadow brome grass, seedling streambank wheatgrass and reed canary grass.</b></p> <p>No surfactant required.</p>
<p>POWERMAX Herbicide</p> <p>+</p> <p>MCPA amine (500 g/L formulation; if</p>	<p>0.83 – 1.27</p> <p>+</p>	<p>Volunteer cereals, wild oats, green foxtail, downy brome, giant foxtail, Persian darnel.</p> <p>Volunteer canola (rapeseed)(non Roundup Ready), wild mustard,</p>	<p>Weeds should be less than 15 cm tall and actively growing for best results.</p> <p>Use higher rate if weeds are beyond 8 cm in height.</p>

TANK MIXTURES	RATE (L/ha)	WEEDS CONTROLLED♦	COMMENTS (Apply in 50-100 L/ha water)
<p>another formulation is used, adjust rate accordingly).</p>	<p>0.5 – 0.7</p>	<p>flixweed, redroot pigweed, lady’s thumb, stinkweed, kochia, lamb’s quarters, hempnettle, Russian thistle, volunteer flax, common ragweed*, Canada fleabane, wild buckwheat**, narrow-leaved hawk’s beard***</p> <p>Volunteer Roundup Ready canola (1-4 leaf stage)<sup>3</sup>, bluebur<sup>4</sup>, burdock<sup>4</sup> (before 4 leaf stage), false flax<sup>4</sup>, flixweed<sup>4</sup>, lamb’s quarters<sup>4</sup>, mustards<sup>4</sup> (except dog and tansy), prickly lettuce<sup>4</sup>, ragweeds<sup>4</sup>, redroot pigweed<sup>4</sup>, Russian pigweed<sup>4</sup>, shepherd’s purse<sup>4</sup>, stinkweed<sup>4</sup> (field pennycress), vetch<sup>4</sup>, wild radish<sup>4</sup>, wild sunflower<sup>4</sup></p>	<p>* DO NOT use these rates on plants greater than 8 cm in height.</p> <p>** For 3- to 4-leaf stage use 1.27 L/ha rate.</p> <p>*** For weeds 8 cm to 15 cm in height use 1.27 L/ha rate.</p> <p><sup>3</sup> MCPA amine at 0.5 – 0.7 L/ha (250 – 350 g ai/ha) prior to lentils and chickpeas.</p> <p><sup>4</sup> MCPA amine at 0.7 L/ha (350 g ai/ha) only.</p> <p>Use this tank mix prior to seeding in <b>lentil and chickpea. Under drought conditions, deep seeding and/or brief rain showers after seeding may cause injury to emerging seedlings in sprayer overlaps.</b></p> <p>No surfactant required.</p>
<p>POWERMAX Herbicide + Express Toss-N-Go Herbicide Or Express Toss-N-Go Dry Flowable 75% Herbicide</p>	<p>0.83 – 1.27  +  10 g/ha (7.5 g ai/ha)</p>	<p>Volunteer cereals, Canada thistle (suppression), cow cockle, wild buckwheat, Canada fleabane common ragweed narrow-leaved hawk's beard, dandelion, downy brome, flixweed, giant foxtail, green foxtail, hempnettle, kochia, lady's thumb, lamb's quarters, persian darnel, redroot pigweed,</p>	<p>Use this tank mix in summerfallow or prior to seeding <b>wheat and barley.</b></p> <p>Refer to Express Toss-N-Go label for the appropriate weed growth stage.</p> <p>Add 350 mL/ha of surfactant – see list in section 7.3.</p>

2020-1550  
2020-08-04

<b>TANK MIXTURES</b>	<b>RATE (L/ha)</b>	<b>WEEDS CONTROLLED♦</b>	<b>COMMENTS (Apply in 50-100 L/ha water)</b>
		Russian thistle, stinkweed, volunteer canola, volunteer flax, wild mustard, wild oats	

♦ For foxtail barley, refer to “**Perennial Weed Control**” table (section 8.1).

<sup>B</sup> 0.56 kg ai/ha of 2,4-D. <sup>B</sup>, <sup>A</sup> Adjust rates accordingly for other 2,4-D formulations. Use only low volatile ester or amine formulations of 2,4-D.

<sup>C</sup> Use only amine formulations of MCPA prior to seeding in corn and field peas.

Banvel II is a registered trademark of BASF.

Pardner and Buctril® are registered trademarks of Bayer.

Express is a registered trademark of E.I.duPont de Nemours and Company.

Toss-N-Go is a registered trademark of E. I. duPont Canada Company.

2020-1550

2020-08-04

### 7.3 SURFACTANT INFORMATION

**NOTE:**

**Addition of Surfactant** – POWERMAX Herbicide tank mixtures for annual weed control may require the addition of a surfactant registered for use such as Agral 90, Ag-Surf or Companion. Refer to Section 7.2 for recommendations. Surfactant should be added at a rate of 350 millilitres per hectare, in 50 - 100 litres of clean water.

### 7.4 ADDITIONAL IMPORTANT INFORMATION FOR ANNUAL WEED CONTROL

POWERMAX Herbicide applied alone will not control volunteers from crops containing the Roundup Ready® gene.

Allow at least 1 day after treatment before tillage.

Annual weeds generally will continue to germinate from seed throughout the growing season. Repeat treatments may be necessary to control later germinating weeds, in some situations.

For additional information and precautions, refer to “**General Information**” and “**Mixing and Application**” (Sections 4.0 and 5.0, respectively).

### 7.5 WEED CONTROL IN TRUFLEX ROUNDUP READY CANOLA VARIETIES

**WARNING: APPLY POWERMAX HERBICIDE TO TRUFLEX ROUNDUP READY CANOLA VARIETIES ONLY.**

**NOTE: ALWAYS USE PEDIGREED (I.E., CERTIFIED) TRUFLEX ROUNDUP READY CANOLA SEED. CANOLA NOT DESIGNATED AS TRUFLEX ROUNDUP READY WILL BE DAMAGED OR DESTROYED BY THIS TREATMENT.**

- For additional information and precautions refer to “**General Information**” and “**Mixing and Application**” (sections 4.0 and 5.0, respectively).
- Apply to TruFlex Roundup Ready canola only as directed.

DO NOT APPLY BY AIR

The following table describes the rate and specific application instructions for weed control in TruFlex Roundup Ready canola varieties.

### WEED CONTROL IN TRUFLEX ROUNDUP READY CANOLA VARIETIES

<b>RATE (L/ha)</b>	<b>GROWTH STAGE OF CROP</b>	<b>WEEDS CONTROLLED</b>	<b>COMMENTS (Apply in 50 –100 L/ha water)</b>
0.55-0.83 Single application	Emergence to first flower*	<p><b><u>Annual Grasses</u></b> Wild oats, green foxtail, volunteer barley, volunteer wheat, barnyard grass</p> <p><b><u>Annual Broadleaves</u></b> Stinkweed, redroot pigweed, wild mustard, Russian thistle, lamb’s- quarters, non-Roundup Ready volunteer canola (rapeseed), hempnettle, lady’s- thumb, kochia, chickweed, corn spurry, wild tomato, cleavers, wild buckwheat, shepherd’s purse<sup>1</sup>, cow cockle<sup>1</sup>, night- flowering catchfly<sup>1</sup>, smartweed<sup>1</sup>, <u>stork’s-bill, flixweed, narrow-leaved hawk’s beard</u></p> <p><b><u>Perennials: (Suppression)</u></b> Canada thistle, perennial sow thistle and dandelion</p> <p><b><u>Perennials: (Season-long control)</u></b> Quackgrass,</p>	<p><sup>1</sup>The 0.55 l/ha rate can be used for control of shepherd’s purse, cow cockle and night-flowering catchfly at the 1– 3 leaf stage of the crop or for control of smartweed at the 4 –6 leaf stage.</p> <p>Repeat applications may be required if a second flush of weeds germinates prior to canopy closure.</p>
1.27 Single application	Emergence to first flower *	<p><b>All the above weeds plus:</b> <b><u>Perennials (season-long control)</u></b> Canada thistle, and perennial sow thistle</p>	
0.83 Sequential applications	Emergence to first flower *	<p><b>All the above weeds plus:</b> <b><u>Annual Broadleaves</u></b> round-leaved mallow</p> <p><b><u>Perennials (season-long control)</u></b> foxtail barley, Canada thistle, and perennial sow thistle</p>	For sequential applications, ensure the crop has not advanced beyond the recommended growth stage
1.67 Single application	Emergence to first flower *	<p><b>All the above weeds plus:</b> Foxtail barley, smooth pigweed, common ragweed, cocklebur, eastern black nightshade, pennsylvannia smartweed, foxtail (yellow and giant), fall panicum, wild proso millet, crabgrass (smooth and</p>	<p><sup>2</sup> Biennial wormwood should be at 2-8 leaf stage and actively growing.</p> <p><sup>3</sup>For control of volunteer adzuki beans (unifoliolate to the 4<sup>th</sup> trifoliolate leaf stage)</p>

		<p>large), velvet leaf, biennial wormwood<sup>2</sup> wire-stemmed muhly, volunteer adzuki beans<sup>3</sup></p> <p><b>(Suppression only)</b> Common Milkweed Yellow nutsedge</p>	<p>apply 1.67 L/ha. A second 1.67 L/ha application may be used for late flushes emerging after the initial treatment. Adzuki beans should be at unifoliate to fourth trifoliate leaf stage and actively growing.</p>
1.67 Sequential applications	Emergence to first flower *	<p><b>All the above weeds plus:</b> <b><u>Perennials (season-long control)</u></b> Dandelion Common Milkweed Field Bindweed Yellow nutsedge Horsenettle, Tall waterhemp Bur cucumber</p>	<p>A sequential application may be made at least 2 weeks after the first application.</p> <p>A second 1.67 L/ha application may be used for late weed flushes emerging after the initial treatment.</p> <p>Common milkweed should be 15-60 cm in height and actively growing.</p> <p>Yellow nutsedge should be 5-15 cm in height and actively growing.</p> <p>Horse-nettle (2-12-leaf stage)</p> <p>Tall waterhemp up to and including the 18 leaf stage)</p> <p>Bur Cucumber from the 1-18 leaf stage.</p>
3.33 Single application	Emergence to 6 leaf	All the above weeds	One application allowed in crop per season

\* First flower is when 50% of the plants in the field have no more than one flower.

Ensure the crop has not advanced beyond the recommended growth stage for all applications.

**Guidelines:**

2020-1550

2020-08-04

Repeat applications may be required if a second flush of weeds germinates prior to canopy closure.

Maximum 3.33L/ha is allowed for the postemergence use.

### **7.5.1 TRUFLEX ROUNDUP READY HYBRID CANOLA SEED PRODUCTION**

#### **For Use only in TRUFLEX ROUNDUP READY Canola Seed Production Systems**

Apply using ground boom spray equipment.

POWERMAX Herbicide may be applied for the control of non-glyphosate tolerant canola pollen parental line(s) in hybrid canola seed production fields containing both TruFlex Roundup Ready line(s) and non-TruFlex Roundup Ready line(s).

When pollination is complete or near completion, non-TruFlex Roundup Ready pollen parental line(s) may be controlled with an application of 0.83 to 1.67 litres per hectare of POWERMAX Herbicide applied in 50 to 200 litres per hectare water.

Sequential applications (**maximum 2 applications**) may be used for the control of pollen parental line(s) but the total maximum rate applied must not exceed 1.67 litres per hectare. Allow at least 5 days between sequential applications.

### **7.6 WEED CONTROL IN ROUNDUP READY® CANOLA VARIETIES**

**WARNING: APPLY POWERMAX HERBICIDE ON ROUNDUP READY® CANOLA VARIETIES ONLY.**

**NOTE: ALWAYS USE PEDIGREED (I.E., CERTIFIED) ROUNDUP READY® CANOLA SEED. CANOLA WHICH IS NOT DESIGNATED AS ROUNDUP READY® WILL BE DAMAGED OR DESTROYED BY THIS TREATMENT.**

- For additional information and precautions refer to “**General Information**” and “**Mixing and Application**” (sections 4.0 and 5.0, respectively).
- Apply POWERMAX Herbicide in Roundup Ready® canola only as directed in the following weed control table.
- Some short-term, visual yellowing may occur when POWERMAX Herbicide is applied at the late application (4 to 6 leaf stage) of the crop. This effect is temporary and will not influence crop growth, maturity or yield.

2020-1550  
2020-08-04

**DO NOT APPLY BY AIR.**

The following table describes the rate and specific application instructions for control of annual and perennial weeds in Roundup Ready® canola varieties.

**WEED CONTROL IN ROUNDUP READY® CANOLA VARIETIES**

<b>RATE (L/ha)</b>	<b>GROWTH STAGE OF CROP</b>	<b>WEEDS CONTROLLED</b>	<b>COMMENTS (Apply in 50 –100 L/ha water)</b>
0.55 – 1.27	0 to 6 leaf	<p><b><u>Annual Grasses</u></b> Wild oats, green foxtail, volunteer barley, volunteer wheat, barnyard grass</p> <p><b><u>Annual Broadleaves</u></b> Stinkweed, redroot pigweed, wild mustard, Russian thistle, lamb’s-quarters, non-Roundup Ready volunteer canola (rapeseed), hempnettle, lady’s-thumb, kochia, chickweed, corn spurry, wild tomato, cleavers*, wild buckwheat*, shepherd’s purse*, cow cockle*, night-flowering catchfly*, smartweed*, stork’s-bill*, flixweed*, narrow-leaved hawk’s beard*, round-leaved mallow***</p> <p><b><u>Perennials (suppression)**</u></b> Canada thistle, perennial sow thistle, dandelion</p> <p><b><u>Perennials (season-long control)</u></b> Quackgrass**, foxtail barley***, Canada thistle****, and perennial sow thistle****</p>	<p>Repeat applications may be required if a second flush of weeds germinates prior to canopy closure.</p> <p>Ensure the crop has not advanced beyond the recommended growth stage.</p> <p><b>* Use the 0.83 L/ha rate for control of these weeds at all crop growth stages.</b> The lower rate can be used for control of shepherd’s purse, cow cockle and night-flowering catchfly at the 1– to 3-leaf stage of the crop or for control of smartweed at the 4– to 6-leaf stage.</p> <p>** A single application of 0.83 L/ha is required.</p> <p>*** Sequential applications of 0.83 L/ha are required.</p> <p>**** Sequential applications of 0.83 L/ha are required or a single application of 1.27 L/ha.</p> <p>For sequential applications, ensure the crop has not advanced beyond the recommended growth stage.</p> <p>Maximum 1.66 L/ha is allowed for the postemergence use.</p>

2020-1550  
2020-08-04

### **7.6.1 TANK MIXTURES**

For season long control of top growth of Canada thistle and control of wild buckwheat in Roundup Ready® canola varieties, apply a tank mixture of 0.28 L/ha of Lontrel 360 with 0.83 L/ha of POWERMAX Herbicide, in 100 litres of water per hectare. Apply when canola is in the 2- to 6-leaf stage. Refer to the Lontrel 360 and to the POWERMAX Herbicide labels for a list of other weeds controlled, timing of application, water volumes and use precautions.

Lontrel® is a registered trademark of Dow AgroSciences LLC.

### **7.6.2 ROUNDUP READY® HYBRID CANOLA SEED PRODUCTION**

#### **For Use only in Roundup Ready® Hybrid Canola Seed Production Systems**

Apply using ground boom spray equipment.

POWERMAX Herbicide may be applied for the control of non-Roundup Ready® canola pollen parental line(s) in hybrid canola seed production fields containing both Roundup Ready® line(s) and non-Roundup Ready® line(s).

When pollination is complete or near completion, non-Roundup Ready® pollen parental line(s) may be controlled with an application of 0.83 to 1.67 litres per hectare of POWERMAX Herbicide applied in 50 to 200 litres per hectare water.

Sequential applications (**maximum 2 applications**) may be used for the control of pollen parental line(s) but the total maximum rate applied must not exceed 1.67 litres per hectare. Allow at least 5 days between sequential applications.

### **7.7 WEED CONTROL IN ROUNDUP READY OR ROUNDUP READY2 YIELD® SOYBEAN VARIETIES**

#### **7.7.1 WEED CONTROL IN ROUNDUP READY2 YIELD SOYBEAN VARIETIES**

**WARNING: APPLY POWERMAX HERBICIDE ON ROUNDUP READY2 YIELD SOYBEAN VARIETIES ONLY.**

**NOTE: ROUNDUP READY 2 YIELD SOYBEAN VARIETIES ARE TOLERANT OF GLYPHOSATE, THE ACTIVE INGREDIENT IN POWERMAX HERBICIDE. ALWAYS USE PEDIGREED (I.E., CERTIFIED) SOYBEAN SEED DESIGNATED AS ROUNDUP READY2 YIELD. SOYBEANS WHICH ARE NOT**

**DESIGNATED AS ROUNDUP READY2 YIELD WILL BE DAMAGED OR DESTROYED BY THIS TREATMENT.**

DO NOT APPLY BY AIR.

RATE (L/ha)	GROWTH STAGE OF CROP	WEEDS CONTROLLED◆	COMMENTS (Use 100 – 200 L/ha water volumes)
1.67	First trifoliolate leaf stage through flowering	<p>Velvetleaf, common ragweed, common lamb's quarters, redroot pigweed, smooth pigweed, cocklebur, green smartweed, lady's-thumb, Pennsylvania smartweed, Eastern black nightshade, wild mustard, wild buckwheat, foxtail (green, yellow, giant), barnyard grass, crabgrass (smooth, large), quackgrass, fall panicum, wild proso millet, wild oats, volunteer barley, volunteer wheat, stinkweed, Russian thistle, non-Roundup Ready canola (rapeseed), hemp-nettle, kochia, chickweed, corn spurry, wild tomato, cleavers, shepherd's purse, cow cockle, night flowering catchfly, stork's bill, flixweed, narrow leaved hawk's-beard</p> <p>common milkweed<sup>1,2</sup>, yellow nutsedge<sup>1,2</sup>, field bindweed<sup>2</sup>, perennial sow thistle, Canada thistle. wire-stemmed muhly.</p> <p>Bur cucumber (<i>Sicyos angulatus</i>)<sup>3</sup></p>	<p><sup>1</sup> A single application of 1.67 L/ha will provide suppression only.</p> <p><sup>2</sup> For control of common milkweed, yellow nutsedge, round-leaved mallow and field bindweed, a second sequential application may be at least 2 weeks after the first application.</p> <ul style="list-style-type: none"> <li>• A second 1.67 L/ha application may be used for late weed flushes emerging after the initial treatment.</li> <li>• Any second application made must be applied no later than the flowering stage of the soybean.</li> <li>• Common milkweed should be 15-60 cm in height and actively growing.</li> <li>• Yellow nutsedge should be 5-15 cm in height and actively growing.</li> <li>• Perennial sow thistle and Canada thistle should be from the rosette stage to 50 cm in height and actively growing.</li> </ul>

RATE (L/ha)	GROWTH STAGE OF CROP	WEEDS CONTROLLED◆	COMMENTS (Use 100 – 200 L/ha water volumes)
		Volunteer adzuki beans ( <i>Vigna angularis</i> ) <sup>4</sup>  Biennial Wormwood ( <i>Artemisia biennis</i> ) <sup>5</sup>	<ul style="list-style-type: none"> <li>• Wire-stemmed muhly should be 10-20 cm in height and actively growing.</li> <li>• Plants not fully emerged at the time of application will escape treatment.</li> <li>• <sup>3</sup>Sequential applications of 1.67 L/ha followed by 1.67 L/ha at the 1-18 leaf stage. Applications should be at least 2 weeks apart for best results.</li> <li>• <sup>4</sup>For control of volunteer adzuki beans (unifoliolate to the 4<sup>th</sup> trifoliolate leaf stage) apply 1.67 L/ha. A second 1.67 L/ha application may be used for late flushes emerging after the initial treatment. Adzuki beans should be at unifoliolate to fourth trifoliolate leaf stage and actively growing</li> <li>• <sup>5</sup> Only one application per season at 1.67L/ha. Biennial wormwood should be at 2-8 leaf stage and actively growing.</li> </ul>
3.33	First trifoliolate leaf stage through flowering	All weeds listed above plus horse-nettle <sup>6</sup> and tall waterhemp <sup>6</sup>	<ul style="list-style-type: none"> <li>• Only one application per season at 3.33 L/ha.</li> <li>• Common milkweed should be 15-60 cm in height and actively growing.</li> </ul>

RATE (L/ha)	GROWTH STAGE OF CROP	WEEDS CONTROLLED♦	COMMENTS (Use 100 – 200 L/ha water volumes)
			<ul style="list-style-type: none"> <li>• Yellow nutsedge should be 5-15 cm in height and actively growing.</li> <li>• Plants not fully emerged at the time of application will escape treatment.</li> </ul> <p><sup>6</sup>For season-long control of horse-nettle (<i>Solanum carolinense</i>) (2- to 12-leaf stage) or, for control of tall waterhemp (<i>Amaranthus tuberculatos</i>) (up to and including the 18-leaf stage) apply 3.33 L/ha. Alternatively, sequential applications of 1.67 L/ha followed by 1.67 L/ha may be applied. Applications should be at least 2 weeks apart for best results.</p> <p><sup>6</sup>For the control of Tall Waterhemp use the higher rate if weeds are beyond the 6-leaf stage.</p>
4.67	First trifoliolate leaf stage through flowering	All weeds listed above plus control of volunteer alfalfa and bromegrass	<p>Only one application per season at 4.67 L/ha.</p> <p>Alfalfa should have 9 or more leaves and be at least 10-15 cm tall.</p> <p>Bromegrass should have at least 3-5 leaves and be at least 10-15 cm tall.</p> <p>Short term yellowing may occur in sprayer overlap areas with the 4.67 L/ha application rate. This effect is temporary and will not influence crop growth or yield.</p>

2020-1550

2020-08-04

◆ Weeds will be more easily controlled and early crop competition avoided with applications made when the weeds are small. Control of annual weeds greater than 25 cm in height will be inconsistent, although some weeds may be controlled.

### **7.7.2 WEED CONTROL IN ROUNDUP READY SOYBEAN VARIETIES**

**WARNING: APPLY POWERMAX HERBICIDE ON ROUNDUP READY SOYBEAN VARIETIES ONLY.**

**NOTE: ALWAYS USE PEDIGREED (I.E., CERTIFIED) SOYBEAN SEED DESIGNATED AS ROUNDUP READY. SOYBEANS WHICH ARE NOT DESIGNATED AS ROUNDUP READY WILL BE DAMAGED OR DESTROYED BY THIS TREATMENT.**

DO NOT APPLY BY AIR.

Apply 1.67 – 3.33 L/ha of POWERMAX Herbicide to Roundup Ready soybean varieties.

See Section 7.6.1 for use directions.

Do not apply the 4.67 L/ha rate to non-Roundup Ready<sup>2</sup> Yield soybean varieties.

### **7.7.3 TANK MIXTURES**

**Tank mixtures may be applied to both Roundup Ready<sup>2</sup> Yield and Roundup Ready soybean varieties.**

#### **POWERMAX Herbicide Plus Pursuit Herbicide**

For added residual control of late germinating eastern black nightshade, common lamb's quarters, redroot pigweed, velvetleaf, fall panicum and wild proso millet, Pursuit herbicide may be tank mixed with POWERMAX Herbicide at a rate of 1.67 liters per hectare. Use 0.16 to 0.21 liters per hectare of Pursuit and apply up to and including the 3<sup>rd</sup> trifoliolate leaf stage of the Roundup Ready soybeans in 100-200 liters per hectare of clean water. The higher rate is recommended for heavier infestations. This tank mix is recommended primarily for soybean systems with row spacings of 50 centimeters (20 inches) or more where a single application timing is desired.

Mixing: Add and mix Pursuit as per instructions on the Pursuit label and then add POWERMAX Herbicide as per instructions on this label.

2020-1550

2020-08-04

A PHI of 100 days is required for the tank mix of POWERMAX Herbicide and Pursuit herbicide on Roundup Ready2 yield soybeans.

Only one application per season of POWERMAX Herbicide at 1.67 liters per hectare tank mixed with Pursuit herbicide at 0.16 to 0.21 liters per hectare is permitted.

Refer to the Pursuit herbicide label for further safety precautions and handling instructions.

**POWERMAX Herbicide Plus FirstRate™ Herbicide (For Use in Eastern Canada Only)**

For added residual control of common ragweed, velvetleaf, cocklebur, jimsonweed and giant ragweed, FirstRate Herbicide may be tank mixed with POWERMAX Herbicide at a rate of 0.83 - 1.67 liters per hectare. Use 20.8 grams per hectare of FirstRate Herbicide.

Do not harvest soybean plants for forage or hay. Do not harvest soybeans for 65 days after application.

Only one application per season of POWERMAX Herbicide tank mixed with FirstRate Herbicide is permitted.

Refer to the FirstRate Herbicide label for further safety precautions and handling instructions.

**POWERMAX Herbicide and Classic 25 DF Herbicide\***

For season-long control of dandelion, annual sow thistle, and yellow nutsedge\*, apply Classic 25 DF Herbicide at 36 grams per hectare plus POWERMAX Herbicide at 1.67 litres per hectare. Add a non-ionic surfactant such as Agral 90, Citowett Plus, or Ag-Surf at 0.2% v/v. Apply when soybeans are in the 1-3 trifoliolate stage; dandelions and annual sow thistle less than 15 cm tall and across; and up to the 8 leaf stage for yellow nutsedge. **USE THIS TANK MIXTURE ONLY ON SOYBEANS WITH THE ROUNDUP READY® TRAIT.**

Consult the Classic 25 DF Herbicide label for tank mixing instructions and use precautions including instructions on replanting to other crops.

\*Use this tank mix only in cases of heavy infestation of yellow nutsedge.

**POWERMAX Herbicide plus Sencor® 75 DF Herbicide for Control of Spreading Atriplex (Eastern Canada only)**

2020-1550  
2020-08-04

For the control of spreading atriplex, apply a preplant application of Sencor 75 DF Herbicide at 0.75 - 1.11 kg product per hectare on medium textured soils or 1.11 – 1.5 kg product per hectare on fine textured soils plus POWERMAX Herbicide at 1.67 litres per hectare. Do not apply on coarse textured soils. Apply when spreading atriplex is up to the 10-leaf stage of growth. Only one application per year is permitted.

Refer to the Sencor 75 DF Herbicide label for further use directions, safety precautions and handling instructions. Consult Table entitled "Sencor 75 DF Alone: Preemergence Application" for specific rates based on soil types and organic matter.

2020-1550  
2020-08-04

### **POWERMAX Herbicide plus Assure® II Herbicide**

<b>RATE</b>	<b>GROWTH STAGE OF CROP</b>	<b>WEEDS CONTROLLED ♦</b>	<b>COMMENTS</b>
1.67 – 3.33 L/ha POWERMAX Herbicide  +  0.25 - 0.38 L/ha Assure II Herbicide	First trifoliolate leaf stage through flowering.	Volunteer Roundup Ready corn.  Apply at the 2- to 6- leaf stage of the weed.	See additional information following this table.

\*Sure Mix may or may not be added to this tank mix

♦ Weeds will be more easily controlled and early crop competition avoided with applications made when the weeds are small. Control of annual weeds greater than 25 centimetres in height will be inconsistent, although some weeds may be controlled.

#### **Volunteer Roundup Ready Corn Control**

For control of volunteer Roundup Ready corn, Assure II herbicide may be tank mixed with POWERMAX Herbicide. Use 1.67 to 3.33 litres per hectare POWERMAX Herbicide and 0.25 - 0.38 litre per hectare of Assure II herbicide.

The higher rate of Assure II may be required when there are high populations of volunteer Roundup Ready corn, other grass weeds are present or when conditions at application are not favorable for weed growth.

Apply in 100 to 300 litres per hectare of clean water.

**Mixing:** Add and mix Assure II herbicide as per instructions on the Assure II herbicide label and then add POWERMAX Herbicide as per instructions on this label.

This tank mix is to be applied when the crop is from the first trifoliolate leaf stage through flowering and when the volunteer Roundup Ready corn is at the 2- to 6-leaf stage.

A PHI (preharvest interval) of 80 days is required for the tank-mix of POWERMAX Herbicide and Assure II herbicide on Roundup Ready2 Yield soybeans.

Refer to the Assure II Herbicide label for further safety precautions and handling instructions.

2020-1550  
2020-08-04

### POWERMAX Herbicide plus Venture® L Herbicide

RATE	GROWTH STAGE OF CROP	WEEDS CONTROLLED ♦	COMMENTS
1.67 – 3.33 L/ha POWERMAX Herbicide + 0.45 - 0.60 L/ha Venture L Herbicide**	First trifoliolate leaf stage through third trifoliolate leaf stage	Volunteer Roundup Ready corn.  Apply at the 2- to 5- leaf stage of the weed.	See additional information following this table.

\*Turbocharge may or may not be added to this tank mix

♦ Weeds will be more easily controlled and early crop competition avoided with applications made when the weeds are small. Control of annual weeds greater than 25 centimetres in height will be inconsistent, although some weeds may be controlled.

For control of volunteer Roundup Ready corn, Venture L Herbicide may be tank mixed with POWERMAX Herbicide. Use 1.67 to 3.33 litres per hectare POWERMAX Herbicide and 0.45 - 0.60 litre per hectare of Venture L Herbicide.

The higher rate of Venture L Herbicide may be required when there are high populations of volunteer Roundup Ready corn, other grass weeds are present or when conditions at application are not favorable for weed growth.

Apply in 100 to 200 litres per hectare of clean water.

**Mixing:** Add and mix Venture L Herbicide as per instructions on the Venture L Herbicide label and then add POWERMAX Herbicide as per instructions on this label.

This tank mix is to be applied when the crop is from the first trifoliolate leaf stage through third trifoliolate leaf stage and when the volunteer Roundup Ready corn is at the 2- to 5- leaf stage.

A PHI (preharvest interval) of 90 days is required for the tank-mix of POWERMAX Herbicide and Venture L Herbicide on Roundup Ready<sup>2</sup> Yield and Roundup Ready Soybeans.

Refer to the Venture L Herbicide label for further safety precautions and handling instructions.

FirstRate is a trademark of Dow AgroSciences LLC.

Pursuit is a registered trademark of BASF.

Sencor is a registered trademark of Bayer.

Assure and Classic are registered trademarks of E.I. duPont de Nemours and Company.

2020-1550  
2020-08-04

Venture is a registered trademark of a Syngenta group company.

## 7.8 WEED CONTROL IN ROUNDUP READY 2™ XTEND SOYBEANS

### Powermax Herbicide and XtendiMAX with VaporGrip Technology Herbicide Use In Roundup Ready 2 Xtend Soybeans

Apply Powermax Herbicide with XtendiMAX with VaporGrip Technology Herbicide alone or in tank mixture with in 100-220 L/ha

DO NOT apply by air.

Weeds Controlled *	Rates	Timing
<p><b>Annual Broadleaved Weeds:</b> velvetleaf, common ragweed, giant ragweed, false ragweed, common lamb's-quarters, redroot pigweed, smooth pigweed, Russian pigweed, cocklebur, green smartweed, lady's-thumb, Pennsylvania smartweed, E. black nightshade, wild mustard, hare's-ear mustard, Indian mustard, tumble mustard, wormseed mustard, wild buckwheat, tartary buckwheat, Canada fleabane, stinkweed, Russian thistle, non-glyphosate tolerant canola (rapeseed), hempnettle, kochia, chickweed, corn spurry, wild tomato, cleavers, shepherd's-purse, cow cockle, night flowering catchfly, stork's bill, narrow leaved hawk's-beard, flixweed, bur cucumber(1), volunteer adzuki beans(2), biennial wormwood(3)</p> <p><b>Annual Grass Weeds:</b> foxtail (green, yellow, giant), barnyardgrass, crabgrass (smooth, large), fall panicum</p>	<p>Powermax Herbicide at 1.67 L/ha + XtendiMAX with VaporGrip Technology Herbicide at 823 ml to 1.71 L/ha</p>	<p>Preplant or Preemergence to the crop  and/or  Postemergence to the crop once or twice up to the early flower stage of the crop (R1)</p> <p><u>Note:</u> The 1.71 L/ha rate of XtendiMAX with VaporGrip Technology Herbicide is to be used only once in a season and should be used preplant, preemergence or in-crop early postemergence</p> <p><u>Note:</u> 3.36 L/ha of XtendiMAX with VaporGrip Technology Herbicide is the maximum total to be applied to <b>Roundup Ready 2 Xtend Soybeans</b> in a single growing season (year).</p> <p>A third application of XtendiMAX with VaporGrip Technology Herbicide should only be made for the control</p>

2020-1550  
2020-08-04

<p>wild proso millet, wild oats, volunteer barley, volunteer wheat ,</p> <p><b>Perennial Weeds:</b> Dandelion (4) quackgrass(5), Canada Thistle(5), perennial sow thistle(5), wire-stemmed muhly(5), yellow nutsedge (6,7), common milkweed(6,7), field bindweed(7)</p>		<p>of glyphosate resistant weed populations.</p>
<p>All Weeds listed above plus tall water hemp (8), and horsenettle (8),</p>	<p>Powermax Herbicide at 3.33 L/ha + XtendiMAX with VaporGrip Technology Herbicide at 823 ml to 1.71 L/ha</p>	<p>See notes above for application details. - 1 application per season at 3.33 L/ha</p>
<p>All Weeds listed above plus volunteer alfalfa (9) and bromegrass (9)</p>	<p>Powermax Herbicide at 4.67 L/ha (10) + XtendiMAX with VaporGrip Technology Herbicide at 823 ml to 1.71 L/ha</p>	<p>See notes above for application details. - 1 application per season at 4.67 L/ha.</p>

\* Weeds will be more easily controlled and early crop competition avoided with applications made when the weeds are small. Control of annual weeds greater than 25 cm in height will be inconsistent, although some weeds may be controlled.

**Application Footnotes:**

(1) Two applications including Powermax Herbicide at 1.67 L/ha applied when the bur cucumber is at the 1 to 18 leaf stage. Applications should be at least 2 weeks apart for best results.

2020-1550

2020-08-04

(2) Applications including Powermax Herbicide at 1.67 L/ha applied at the unifoliolate to 4<sup>th</sup> trifoliolate leaf stage of the Adzuki beans. **A second 1.67 L/ha application may be used for late flushes emerging after the initial treatment when the Adzuki beans are in the unifoliolate to 4<sup>th</sup> trifoliolate leaf stage and actively growing.**

(3) **One application** including Powermax Herbicide at 1.67 L/ha applied at the 2-8 leaf stage of actively growing biennial wormwood.

(4) Applications including Powermax Herbicide applied preplant surface or pre-emergence at 1.67 to 3.33 L/ha. Use Powermax Herbicide rates of 2.47 to 3.33 L/ha on heavy infestations of dandelions and on dandelions greater than 15 cm in size. Apply up to and including bloom for best results.

(5) Applications including Powermax Herbicide at 1.67 L/ha applied when quackgrass has 3-4 leaves, Canada thistle and perennial sow thistle are rosette to 50 cm in height, and wire-stemmed muhly is 10-20 cm in height. Weeds should be actively growing at application.

**(6) Applications including Powermax Herbicide at 1.67 L/ha will provide suppression.**

(7) For control of common milkweed, yellow nutsedge, and field bindweed a second application **including Powermax Herbicide at 1.67 L/ha may be needed and should be applied at least 2 weeks after the first application or 3.33 L/ha should be applied once. Milkweed should be 15-60 cm in height, yellow nutsedge should be 5-15 cm in height.**

(8) Applications including Powermax Herbicide at 3.33 L/ha applied at the 2-12 leaf stage of horse-nettle or up to the 18 leaf stage of tall waterhemp or 2 applications of 1.67 L/ha applied at least 2 weeks apart. **For control of tall waterhemp use the higher rate if weeds are beyond the 6 leaf stage.**

**(9) Alfalfa should have 9 or more leaves and be at least 10-15 cm tall. Bromegrass should have at least 3-5 leaves and be at least 10-15 cm tall.**

**(10) With the 4.67 L/ha rate some short term yellowing may occur in the sprayer overlap areas, but this effect is temporary and will not influence growth or yield.**

**Pre-Harvest Interval(s):**

- 7-10 days for soybean forage and 13–15 days for soybean hay.
- A plant back interval of 120 days is required for those crops on the XtendiMAX with VaporGrip Technology Herbicide label.

2020-1550

2020-08-04

## **Equipment – DO NOT APPLY THIS PRODUCT TO ROUNDUP READY 2 XTEND SOYBEANS USING AERIAL SPRAY EQUIPMENT**

Apply XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE to weeds < 10 cm

**Water Volume** – apply this product in a minimum of 100 Liters of spray solution per hectare.

### **Spray Drift Management**

Do not allow herbicide solution to mist, drip, drift or splash onto desirable vegetation because severe injury or destruction to desirable broadleaf plants could result. The following drift management requirements must be followed to ensure application accuracy from ground application onto agricultural field crops.

### **Controlling Droplet Size**

The most effective way to reduce drift potential is to apply large droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if the application is made improperly, or under unfavorable environmental conditions (see the “**Wind Speed and Direction**”, “**Temperature and Humidity**” and “**Temperature Inversions**” sections of this label).

- **Nozzle type.** Use only spray nozzles that produce very coarse to ultra coarse spray droplets and minimal amounts of fine spray droplets as defined by the American Society of Agricultural and Biological Engineers (ASABE S-572.1). Do not use conventional flat fan nozzles that produce medium droplets.

Check nozzle manufacturer’s recommendations to determine the proper droplet spectrum, operating pressure, boom height, nozzle spacing and ground speed that will deliver the desired droplet size and spray volume of at least 100 L/ha for the very coarse to ultra coarse nozzle that is selected.

- **Spray Pressure.** Adjust pressure for selected nozzles according to the nozzle manufacturer to maintain very coarse to ultra-coarse droplets. Use sufficient spray pressure with air induction nozzles to ensure a good spray pattern, while maintaining very coarse to ultra coarse droplets; use at least 30 psi to ensure proper pattern overlap. Confirm that sprayer rate controller hardware (if so equipped) does not increase pressure above the desired range. Calibrate the flow rate for the selected nozzles on the equipment used to apply this product.

- **Spray Volume.** Apply this product in a minimum of 100 Liters of spray solution per hectare. Use a higher spray volume when treating dense vegetation. Higher spray volumes also allow the use of larger nozzle orifices (sizes) which produce coarser spray droplets.

- **Equipment Ground Speed.** Select a ground speed under 25 KM/PH that will deliver the desired spray volume while maintaining the desired spray pressure. Slower speeds generally result in better spray coverage and deposition on the target area.
- **Spray boom Height.** Spray at the appropriate boom height based on nozzle selection and nozzle spacing (not more than 20 inches above target pest or crop canopy). Set boom to lowest effective height over the target pest or crop canopy based on equipment manufacturer's directions. Automated boom height controllers are recommended with large booms to better maintain optimum nozzle to canopy height. Excessive boom height will increase the potential for spray drift.

### **Temperature and Humidity**

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

**Temperature Inversions.** Do not apply during a temperature inversion because off-target movement potential is high.

- During a temperature inversion, the atmosphere is very stable and vertical air mixing is restricted, which causes small, suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions.
- Temperature inversions are characterized by increasing temperatures with altitude and are common on evenings and nights with limited cloud cover and light to no wind. Cooling of air at the earth's surface takes place and warmer air is trapped above it. They begin to form as the sun sets and often continue into the morning.
- Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.
- The inversion will dissipate with increased winds (above 5 KM/H) or at sunrise when the surface air begins to warm.

### **Wind Speed and Direction**

- Drift potential is lowest between wind speeds of 5 to 16 Km per hour.
- If the wind speed is 5 km/hr or less and fog is present, indicating a temperature inversion, do not apply this product.
- If fog is not present, conduct a smoke test. Smoke that moves upward confirms there is no inversion present whereas smoke that layers and moves laterally in a concentrated cloud, indicates a temperature inversion exists. Do not apply this product during a temperature inversion. Wait until the wind speed is greater than 5 km/hr to ensure that any inversion has lifted.
- Do not spray this product when the wind is blowing in the direction of a sensitive area

2020-1550  
2020-08-04

at a wind speed greater than 16 km/hr.

• For XTENDIMAX WITH VAPORGRIP TECHNOLOGY HERBICIDE wind speed and direction restrictions see below table:

Wind speed	Application conditions and restrictions
<5 km/hr	Do not apply XtendiMAX with VaporGrip Technology Herbicide if temperature inversion exists
5-16 km/hr	Optimum XtendiMAX with VaporGrip Technology Herbicide application conditions.
16 - 25 km/hr	Do not apply product when wind is blowing toward sensitive areas
> 25 km/hr	Do not apply XtendiMAX with VaporGrip Technology Herbicide

**NOTE:** Local terrain can influence wind patterns. Every applicator must be familiar with local wind patterns and how they affect drift.

#### Sensitive Areas

This product should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for species at risk, or sensitive crop plants) is minimal (e.g. when the wind is blowing away from sensitive areas). Applicators should survey the surrounding area before making an application of this product.

Failure to follow the requirements in this label, could result in severe injury or destruction to desirable sensitive crops and trees, particularly beans, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tobacco, tomatoes, and other broadleaf plants when contacting their roots, stems or foliage.

#### Application Awareness

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR

The interaction of equipment and weather related factors must be monitored to maximize performance and on-target spray deposition. The applicator is responsible for considering all of these factors when making a spray decision.

#### Proper spray system equipment cleanout

Minute quantities of dicamba may cause injury to **non-Roundup Ready 2 Xtend soybeans** and other sensitive crops (see the “Sensitive Areas” section of this label for a listing of sensitive crops).

Clean equipment immediately after using this product, using a triple rinse procedure as follows:

1. After spraying, drain the sprayer (including boom and lines) immediately. Do not allow the spray solution to remain in the spray boom lines overnight prior to flushing.

2020-1550

2020-08-04

2. Flush tank, hoses, boom and nozzles with clean water.
3. Inspect and clean all strainers, screens and filters.
4. Prepare a cleaning solution with a commercial detergent or sprayer cleaner or ammonia according to the manufacturer's directions.
5. Take care to wash all parts of the tank, including the inside top surface. Start agitation in the sprayer and thoroughly re-circulate the cleaning solution for at least 15 minutes. All visible deposits must be removed from the spraying system.
6. Flush hoses, spray lines and nozzles for at least 1 minute with the cleaning solution.
7. Repeat above steps for two additional times to accomplish an effective triple rinse.
8. Remove nozzles, screens and strainers and clean separately in the cleaning solution after completing the above procedures.
9. Appropriately dispose of rinsate from steps 1-7 in compliance with all applicable laws and regulations.
10. Drain sump, filter and lines.
11. Rinse the complete spraying system with clean water.

All rinse water must be disposed of in compliance with municipal, provincial, and federal guidelines.

#### **Additives and tank mixes**

- Nozzle selection is one of the most important parameters for drift reduction. A drift reduction additive may be used with this product to further reduce fine droplets.
- Not all drift reduction additives are compatible with every nozzle type and pesticide / adjuvant combination. Check with the additive manufacturer to insure that the drift additive will work properly with the spray nozzle, spray pressure and your specific spray solution.
- Read and carefully observe all precautions, limitations and all other information on the product label.
- A quality nonionic surfactant (NIS) of at least 70% active may be added to the spray solution at 0.25 percent surfactant concentration. Read and carefully observe all caution statements and other information on the surfactant label.
- Do not add acidifying buffering agents, acidic pH adjusting agents or adjuvants other than agriculturally approved NIS to the spray solution.
- Do not use crop oil concentrates (COC) and methylated seed oils (MSO) as adjuvants when this product is applied with glyphosate-based agricultural herbicides. When XtendiMAX with VaporGrip Technology Herbicide is used with another herbicide that requires the use of a COC or MSO adjuvant follow the label instructions of that product.
- In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact Bayer

2020-1550  
2020-08-04

CropScience at 1-888-283-6847 for information before mixing any pesticide or fertilizer that is not specifically recommended on this label. The user assumes the risk of losses that result from the use of tank mixes that do not appear on this label or that are not specifically recommended by Bayer CropScience Inc.

- Apply XtendiMAX with VaporGrip Technology Herbicide or tank mixtures with XtendiMAX with VaporGrip Technology Herbicide at a minimum spray volume rate of 100 L/ha.

## **7.9 WEED CONTROL IN CORN VARIETIES WITH ROUNDUP READY® 2 TECHNOLOGY**

**WARNING: APPLY POWERMAX HERBICIDE ON ONLY CORN VARIETIES THAT ARE DESIGNATED AS CONTAINING ROUNDUP READY® 2 TECHNOLOGY (I.E. CONTAINS A ROUNDUP READY GENE).**

**NOTE: CORN VARIETIES CONTAINING ROUNDUP READY® 2 TECHNOLOGY ARE TOLERANT OF GLYPHOSATE, THE ACTIVE INGREDIENT IN POWERMAX HERBICIDE. ALWAYS USE PEDIGREED (I.E. CERTIFIED) CORN SEED DESIGNATED AS CONTAINING ROUNDUP READY® 2 TECHNOLOGY. CORN WHICH IS NOT DESIGNATED AS CONTAINING ROUNDUP READY® 2 TECHNOLOGY MAY BE DAMAGED OR DESTROYED BY THIS TREATMENT.**

DO NOT APPLY BY AIR

<b>RATE (L/ha)</b>	<b>GROWTH STAGE OF CROP</b>	<b>WEEDS CONTROLLED♦</b>	<b>COMMENTS (use 100-200 L/ha water volumes)</b>
1.67	Up to and including 8 leaf stage	Velvetleaf, common ragweed, common lamb's-quarters, redroot pigweed, smooth pigweed, cocklebur, green smartweed, lady's-thumb, Pennsylvania smartweed, Eastern black nightshade, wild mustard, wild buckwheat, foxtail (green, yellow, giant), barnyard grass, crabgrass (smooth, large), quackgrass, fall	<sup>1</sup> A single application of 1.67 L/ha will provide suppression only.  <sup>2</sup> For control of common milkweed, yellow nutsedge, round-leaved mallow and field bindweed, a second sequential application may be at least 2 weeks after the first application.  • A second 1.67 L/ha application may be used for late weed

<b>RATE (L/ha)</b>	<b>GROWTH STAGE OF CROP</b>	<b>WEEDS CONTROLLED◆</b>	<b>COMMENTS (use 100-200 L/ha water volumes)</b>
		<p>panicum, wild proso millet, wild oats, volunteer barley, volunteer wheat, stinkweed, wild mustard, Russian thistle, non-Roundup Ready canola (rapeseed), hemp-nettle, kochia, chickweed, corn spurry, wild tomato, cleavers, shepherd's purse, cow cockle, night-flowering catchfly, stork's-bill, flixweed, narrow-leaved hawk's-beard</p> <p>common milkweed<sup>1,2</sup>, yellow nutsedge<sup>1,2</sup>, round-leaved mallow<sup>2</sup>, field bindweed<sup>2</sup>, perennial sow thistle, Canada thistle, wire-stemmed muhly</p>	<p>flushes emerging after the initial treatment.</p> <ul style="list-style-type: none"> <li>• Any second application must be applied no later than the 8 leaf stage of the corn.</li> <li>• Common milkweed should be 15-60 cm in height and actively growing.</li> <li>• Yellow nutsedge should be 5-15 cm in height and actively growing.</li> <li>• Perennial sow thistle and Canada thistle should be from the rosette stage to 50 cm in height and actively growing.</li> <li>• Wire-stemmed muhly should be 10-20 cm in height and actively growing.</li> <li>• Plants not fully emerged at the time of application will escape treatment.</li> </ul>
3.33	Up to and including 6 leaf stage	All weeds listed above	<ul style="list-style-type: none"> <li>• Only one application per season at 3.33 L/ha.</li> <li>• Common milkweed should be 15-60 cm in height and actively growing.</li> <li>• Yellow nutsedge should be 5-15 cm in height and actively growing.</li> </ul>

<b>RATE (L/ha)</b>	<b>GROWTH STAGE OF CROP</b>	<b>WEEDS CONTROLLED◆</b>	<b>COMMENTS (use 100-200 L/ha water volumes)</b>
			<ul style="list-style-type: none"> <li>Plants not fully emerged at the time of application will escape treatment.</li> </ul>

◆ Weeds will be more easily controlled and early crop competition avoided with applications made when the weeds are small. Control of weeds greater than 25 cm in height will be inconsistent, although some weeds may be controlled.

### 7.9.1 TANK MIXTURES

For tank mixtures, add herbicide according to instructions on the product label, and then add POWERMAX Herbicide according to instructions on this label (section 5). Refer to the tank mix herbicide labels for further safety precautions, use recommendations and product handling instructions.

#### DO NOT APPLY BY AIR

<b>RATE</b>	<b>GROWTH STAGE OF CROP</b>	<b>WEEDS CONTROLLED◆</b>	<b>COMMENTS (Use 100-200 L/ha water volumes)</b>
1.67 L/ha POWERMAX Herbicide  +  0.75 – 1.0 kg ai/ha atrazine*	Up to and including the 5-leaf stage.	Residual control of lamb's-quarters, redroot pigweed, common ragweed.	Tank-mix should be used when only a single application timing is desired. Use the higher rate of atrazine for heavier weed infestations.
1.67 L/ha POWERMAX Herbicide  +  2.5 – 3.7 L/ha Marksman Herbicide	Up to and including the 5-leaf stage.	Residual control of lamb's-quarters, redroot pigweed, common ragweed, velvetleaf.	Tank-mix should be used when only a single application timing is desired. Use the higher rate of Marksman for heavier weed infestations.
1.67 L/ha POWERMAX Herbicide  +	Before the corn is 15 cm tall (leaf extended) and/or	Volunteer Roundup Ready canola – up to the 4 leaf stage.	Tank mix is most effective when treating small (4 leaf

2020-1550  
2020-08-04

<b>RATE</b>	<b>GROWTH STAGE OF CROP</b>	<b>WEEDS CONTROLLED ♦</b>	<b>COMMENTS (Use 100-200 L/ha water volumes)</b>
0.56 – 1.12 L/ha 2,4-D Herbicide**	before the 6 leaf stage.		or less) canola plants.
Two applications:  First application: 1.67 L/ha POWERMAX Herbicide + 0.56 L/ha 2,4-D Herbicide**  Second application: 1.67 L/ha POWERMAX Herbicide + 0.42-0.56 L/ha 2,4-D Herbicide**	Before the corn is 15 cm tall (leaf extended) and/or before the 6 leaf stage.	Volunteer Roundup Ready canola – up to the 4 leaf stage.	Tank mix is most effective when treating small (4 leaf or less) canola plants.
1.67 L/ha POWERMAX Herbicide + 13.3 g/ha Peak 75WG Herbicide + 0.3 L/ha Banvel II Herbicide + non ionic surfactant (0.2% v/v)	Spike up to and including the 5 leaf stage.	Volunteer Roundup Ready canola – up to the 4 leaf stage.	Tank mix is most effective when treating small (4 leaf or less) canola plants.
1.67 L/ha POWERMAX Herbicide + 1.1 L/ha Dyvel DSp Liquid Herbicide	Before the corn is 15 cm tall (leaf extended)	Volunteer Roundup Ready canola – up to the 4 leaf stage.	Tank mix is most effective when treating small (4 leaf or less) canola plants.
1.67 L/ha	3 - 8 leaf stage of corn	Eastern black nightshade,	Add Agral 90 at 0.2% v/v

2020-1550  
2020-08-04

<b>RATE</b>	<b>GROWTH STAGE OF CROP</b>	<b>WEEDS CONTROLLED ♦</b>	<b>COMMENTS (Use 100-200 L/ha water volumes)</b>
POWERMAX Herbicide + 0.21 L/ha Callisto® 480SC Herbicide		velvetleaf, redroot pigweed, common ragweed (suppression only) plus emerged annual and perennial weeds	Apply up to the 8 leaf stage of broadleaf weeds  Some perennial weeds may not be controlled with these rates.
1.67 L/ha POWERMAX Herbicide + 0.21 L/ha Callisto® 480SC Herbicide + 0.58 L/ha Aatrex Liquid 480 Herbicide	3 - 8 leaf stage of corn	Eastern black nightshade, velvetleaf, redroot pigweed, common ragweed plus emerged annual and perennial weeds	Add Agral 90 at 0.2% v/v  Apply up to the 8 leaf stage of broadleaf weeds  Some perennial weeds may not be controlled with these rates
1.67 L/ha POWERMAX Herbicide + 2.5 L/ha Primextra® II Magnum® Herbicide	Apply up to and including 6 leaf stage of corn.	Annual grasses and broadleaf weeds, emerged annual or perennial weeds	This tank mix requires the use of a surfactant. AGRAL 90 or Ag-Surf may be used.  Do NOT apply this tank-mix to soils with less than 1% or more than 10% organic matter.
1.67 L/ha POWERMAX Herbicide + 0.625 L/ha Banvel II Herbicide	Spike to 5 leaf	Weeds controlled by POWERMAX Herbicide plus improved control of Velvetleaf and extended control of late germinating, deep rooted annuals	

2020-1550  
2020-08-04

<b>RATE</b>	<b>GROWTH STAGE OF CROP</b>	<b>WEEDS CONTROLLED◆</b>	<b>COMMENTS (Use 100-200 L/ha water volumes)</b>
		on the Banvel II Herbicide label.	
1.67 L/ha POWERMAX Herbicide + 285 g/ha Distinct Herbicide + Non ionic surfactant + 28% UAN	2 to 6 leaf	Weeds controlled by POWERMAX Herbicide plus extended control of late emerging weeds listed on the Distinct Herbicide label.	Non-ionic surfactant applied at 0.2% v/v  28% UAN applied at 1.25% v/v
1.67 L/ha POWERMAX Herbicide + 1.25 L/ha Dual II Magnum Herbicide + 1.0 kg ai/ha atrazine*	Spike to 6 leaf	Weeds controlled by POWERMAX Herbicide plus extended control of annual grass and broadleaf weeds on the tank mix partner labels.	
1.67 L/ha POWERMAX Herbicide + 1.1 L/ha Frontier Herbicide + 1.0 kg ai/ha atrazine*	Emergence to 3 leaf	Weeds controlled by POWERMAX Herbicide plus extended control of annual grass and broadleaf weeds on the tank mix partner labels.	
1.67 L/ha POWERMAX Herbicide + 4.2 L/ha Prowl 400EC Herbicide + 1.0 kg ai/ha atrazine*	Up to and including the 4 leaf stage of corn	Weeds controlled by POWERMAX Herbicide plus extended control of annual grass and broadleaf weeds on the tank mix partner labels.	

2020-1550  
2020-08-04

<b>RATE</b>	<b>GROWTH STAGE OF CROP</b>	<b>WEEDS CONTROLLED ♦</b>	<b>COMMENTS (Use 100-200 L/ha water volumes)</b>
1.67 L/ha POWERMAX Herbicide + 0.21 L/ha Callisto® 480SC Herbicide + Non ionic surfactant	3 to 8 leaf stage of corn	Weeds controlled by POWERMAX Herbicide plus extended control of eastern black nightshade, velvetleaf, redroot pigweed, and common ragweed.	Add non ionic surfactant at 0.2% v/v
1.67 L/ha POWERMAX Herbicide + 2.5 - 3.0 L/ha Primextra II Magnum Herbicide	Spike to 6 leaf stage of corn	Weeds controlled by POWERMAX Herbicide plus extended control of annual grass and broadleaf weeds on the Primextra II Magnum label.	

\* 0.75 to 1.0 kilogram active ingredient atrazine per hectare is equivalent to 1.56 to 2.08 litres per hectare of Atrazine 480™ or Aatrex Liquid 480™.

\*\* 500 g ai/litre of 2,4-D formulation. Adjust rates accordingly for other 2,4-D formulations. Use only low volatile ester or amine formulations of 2,4-D. Some corn hybrids may be injured by an application of 2,4-D. It is recommended that the corn seed provider be contacted regarding the tolerance of the corn hybrid to be treated, to 2,4-D prior to application of this tank mix.

♦ Weeds will be more easily controlled and early crop competition avoided with applications made when the weeds are small. Control of weeds greater than 25 centimetres in height will be inconsistent, although some weeds may be controlled.

Aatrex and Peak are registered trademarks of a Syngenta group company.  
Marksman, Banvel II and Dyvel DS are registered trademarks of BASF Corporation.

#### **7.10 WEED CONTROL IN SWEET CORN VARIETIES WITH ROUNDUP READY 2 TECHNOLOGY**

**WARNING: APPLY POWERMAX HERBICIDE ON ONLY SWEET CORN VARIETIES THAT ARE DESIGNATED AS CONTAINING ROUNDUP READY 2 TECHNOLOGY (I.E. CONTAINS A ROUNDUP READY GENE).**

**NOTE: SWEET CORN VARIETIES CONTAINING ROUNDUP READY 2 TECHNOLOGY ARE TOLERANT OF GLYPHOSATE, THE ACTIVE INGREDIENT IN POWERMAX HERBICIDE. ALWAYS USE PEDIGREED (I.E. CERTIFIED) SWEET CORN SEED DESIGNATED AS CONTAINING ROUNDUP READY 2 TECHNOLOGY. SWEET CORN WHICH IS NOT DESIGNATED AS CONTAINING ROUNDUP READY 2 TECHNOLOGY MAY BE DAMAGED OR DESTROYED BY THIS TREATMENT.**

**WEED CONTROL:**

<b>RATE (L/ha)</b>	<b>GROWTH STAGE OF CROP</b>	<b>WEEDS CONTROLLED♦</b>	<b>COMMENTS (use 100-200 L/ha water volumes)</b>
1.67	Up to and including 8 leaf stage	See Weeds Controlled in Section 7.7 Table	<ul style="list-style-type: none"><li>• See Comments in Section 7.7 Table</li><li>• A second 1.67 L/ha application may be used for late weed flushes emerging after the initial treatment.</li><li>• Any second application must be applied no later than the 8 leaf stage of the corn.</li></ul>
3.33	Up to and including 6 leaf stage	See Weeds Controlled in Section 7.7 Table	<ul style="list-style-type: none"><li>• See Comments in Section 7.7 Table</li><li>• Only one application per season at 3.33 L/ha.</li></ul>

♦ Weeds will be more easily controlled and early crop competition avoided with applications made when the weeds are small. Control of weeds greater than 25 cm in height will be inconsistent, although some weeds may be controlled.

♦ Plants not fully emerged at the time of application will escape treatment.

**TANK MIXES - Do not apply Tank Mixes to sweet corn varieties with Roundup Ready 2 Technology**

Allow a minimum of 30 days between application of this product and harvest.

DO NOT APPLY BY AIR

**7.11 WEED CONTROL IN ROUNDUP READY® SUGAR BEETS**

**WARNING: APPLY POWERMAX HERBICIDE ON ROUNDUP READY® SUGAR BEET VARIETIES ONLY.**

2020-1550  
2020-08-04

**NOTE: ALWAYS USE PEDIGREED (CERTIFIED) SUGAR BEET SEED DESIGNATED AS ROUNDUP READY®. SUGAR BEETS WHICH ARE NOT DESIGNATED AS ROUNDUP READY® WILL BE DAMAGED OR DESTROYED BY THIS TREATMENT.**

DO NOT APPLY BY AIR.

For weed control in Roundup Ready® sugar beets apply 0.83 – 1.67 L/ha of POWERMAX Herbicide to emerged weeds. Refer to “**Annual Weed Control**” and “**Perennial Weed Control**” (Sections 7.1 and 8.1, respectively) for a listing of weeds controlled.

Apply POWERMAX Herbicide to emerged weeds up to 15 cm in height.

Up to four applications of POWERMAX Herbicide may be applied to Roundup Ready® sugar beets. Allow a minimum of 10 days between applications.

**Do not exceed a total maximum quantity of 7.31 L/ha of this product per season (e.g. the first application of up to 2.30 L/ha plus 3 applications of up to 1.67 L/ha).**

Do not harvest Roundup Ready® sugar beets within 30 days after the final application of POWERMAX Herbicide.

#### **7.12 WEED CONTROL IN ROUNDUP READY ALFALFA VARIETIES (DO NOT APPLY TO ALFALFA GROWN FOR SEED PRODUCTION)**

**WARNING: APPLY POWERMAX HERBICIDE TO ROUNDUP READY ALFALFA VARIETIES ONLY.**

**NOTE: ALWAYS USE PEDIGREED (I.E. CERTIFIED) ALFALFA SEED DESIGNATED AS ROUNDUP READY. ALFALFA SEED WHICH IS NOT DESIGNATED AS ROUNDUP READY WILL BE DAMAGED OR DESTROYED BY THIS TREATMENT.**

**ROUNDUP READY ALFALFA VARIETIES ARE TOLERANT OF GLYPHOSATE, THE ACTIVE INGREDIENT IN POWERMAX HERBICIDE.**

DO NOT APPLY BY AIR.

Applications can be made from emergence until 5 days prior to cutting.

A sequential treatment may be applied to Roundup Ready alfalfa varieties for control of late weed flushes.

Allow a minimum of 5 days between application and cutting of alfalfa.

Additional applications of this product should be at least 25 days apart.

2020-1550  
2020-08-04

**Total number of in-crop applications not to exceed 3 per growing season.**

**New Stand Establishment (Seedling Year):** Due to the biology and breeding constraints of alfalfa, up to 10 percent of the seedlings may not contain a Roundup Ready gene and will not survive or thrive after the first application of this product. To limit the undesirable effects of stand gaps created by the loss of alfalfa plants not containing a Roundup Ready gene, an application of this product should be applied at or before the 4 trifoliolate leaf stage of alfalfa during the establishment (seedling) year.

Note: Where Roundup Ready alfalfa is grown with a companion or cover crop, or is overseeded with a second species, in-crop (over-the-top) applications of this product will eliminate the non-Roundup Ready (non-glyphosate tolerant) species.

**WEED CONTROL IN ROUNDUP READY ALFALFA VARIETIES**

<b>RATE (L/ha)</b>	<b>GROWTH STAGE OF CROP</b>	<b>WEEDS CONTROLLED</b>	<b>COMMENTS (Apply in 50 –100 L/ha water)</b>
1.67 single application	Emergence until 5 days prior to cutting	<p><u>Annual Grasses</u> Wild oats, green foxtail, volunteer barley, volunteer wheat, barnyard grass, giant and yellow foxtail, fall Panicum, wild proso millet, smooth and large crabgrass</p> <p><u>Annual Broadleaves</u> Stinkweed, redroot pigweed, wild mustard, Russian thistle, lamb's-quarters, non-Roundup Ready volunteer canola (rapeseed), hempnettle, lady's-thumb, kochia, chickweed, corn spurry, wild tomato, cleavers, wild buckwheat, shepherd's purse, cow cockle, night-flowering catchfly, smartweed, stork's-bill, flixweed, narrow-leaved hawk's beard, smooth pigweed, cocklebur, Eastern black nightshade, velvetleaf, biennial wormwood<sup>1</sup>.</p> <p><u>Perennials (season-long control)</u> Quackgrass, Canada thistle, and perennial sow thistle, foxtail barley, dandelion.</p>	<p>All weeds should be actively growing at time of application.</p> <p><sup>1</sup>Biennial wormwood should be at 2-8 leaf stage.</p>
3.33 single application	Emergence until 5 days prior to cutting	<p><u>All the above weeds plus:</u> <u>Annual Broadleaves</u> Round-leaved mallow <u>Perennials (season-long control):</u> Foxtail barley<sup>2</sup>, dandelion<sup>2</sup>, common milkweed<sup>3</sup>, field bindweed, yellow nutsedge<sup>4</sup>, horsenettle<sup>5</sup>, tall waterhemp<sup>6</sup>, bur cucumber<sup>7</sup></p>	<p><sup>2</sup>3.33 L/ha rate is for large, more established plants, heavy infestation or if plants are stressed.</p> <p><sup>3</sup>Common milkweed should be 15-60 cm in height.</p>

			<p><sup>4</sup>Yellow nutsedge should be 5-15 cm in height.</p> <p><sup>5</sup>Horse-nettle from the 2 to 12 leaf stage).</p> <p><sup>6</sup>Tall waterhemp up to and including the 18-leaf stage.</p> <p><sup>7</sup>Bur cucumber from the 1-18 leaf stage.</p>
--	--	--	--

### **7.13 HYBRID CORN SEED PRODUCTION USING THE RHS® SYSTEM WITH ROUNDUP READY 2 TECHNOLOGY**

#### **DO NOT APPLY BY AIR**

The RHS designation indicates that the corn contains technology that allows for tassel-only susceptibility to this product. Use of this product on corn hybrids or inbreds that are not designated as RHS or as corn containing Roundup Ready® 2 Technology may result in severe crop injury and yield loss.

#### **Tassel Control**

This product may be used as an over-the-top broadcast application for tassel control in RHS corn inbred recipient lines in seed production fields planted with corn containing Roundup Ready 2 Technology as the pollen donor.

**USE INSTRUCTIONS:** This product may be applied for tassel control up from the 8 to the 13 leaf stage before flowering at use rates from 1.67 to 2.34 L/ha per application. Up to two applications for tassel control are permitted.

#### **Weed Control**

**Refer Only to Section:** 7.8 WEED CONTROL IN CORN VARIETIES WITH ROUNDUP READY 2 TECHNOLOGY

**Tank mixes:** See section 7.8.1 TANK MIXTURES for use rates, timings and restrictions. Note that only those tank mixtures for which the tank mixture partner herbicide products are registered for use on seed (inbred) corn may be used for weed control on RHS corn inbred recipient lines and corn inbred donor lines containing Roundup Ready 2 Technology.

2020-1550  
2020-08-04

## **8.0 PERENNIAL WEED CONTROL**

**ALWAYS READ PRECAUTIONS, GENERAL INFORMATION AND MIXING AND APPLICATION SECTIONS (3.0, 4.0 AND 5.0) PRIOR TO SPECIFIC APPLICATION INFORMATION IN ANY LABEL SECTION. DO NOT APPLY BY AIR.**

When applied as recommended under the conditions described, this product will control the perennial weeds listed in the following table.

### 8.1 PERENNIAL WEED CONTROL WITH POWERMAX HERBICIDE

WEED	APPLICATION			COMMENTS
	GROWTH STAGE	RATE (L/ha)	WATER VOLUME (L/ha)	
<b>Quackgrass</b> (control, light to moderate infestations)	3 to 4 green leaves or more	1.67	50 - 300	<ul style="list-style-type: none"> <li>• Apply in clean water using flat fan nozzles.</li> <li>• Allow 3 or more days after treatment before tillage.</li> <li>• Refer to “<b>Quackgrass</b>” notes in section 8.2.1 for more information.</li> <li>• For higher volumes (i.e., 150 – 300 L/ha) an approved surfactant must be added at 0.5 L per 100 L of clean water (0.5% v/v). Refer to list in section 8.2.2. See also below.</li> </ul>
<b>Quackgrass</b> (long term control, heavy infestations, high water volumes)	3 to 4 green leaves or more	1.67 – 4.67	50 - 300	<ul style="list-style-type: none"> <li>• Allow 3 or more days after treatment before tillage.</li> <li>• Rates higher than 1.67 L/ha will provide more consistent, longer term control, especially with heavier infestations and/or higher water volumes (i.e., 150 – 300 L/ha).</li> <li>• Refer to “<b>Quackgrass</b>” notes in section 8.2.1 for more information.</li> </ul>
<b>Canada Thistle</b>	Rosette stage (summerfallow)	1.67	50 - 100	<ul style="list-style-type: none"> <li>• Apply in clean water using flat fan nozzles.</li> <li>• Allow 10 or more days after treatment before tillage.</li> <li>• Refer to “<b>Canada Thistle</b>” notes in section 8.2.3 for more information.</li> </ul>
<b>Canada Thistle</b>	Bud stage or beyond	3.17 – 4.67	100 - 300	<ul style="list-style-type: none"> <li>• Allow 5 or more days after treatment before tillage.</li> </ul>
<b>Field Bindweed</b>	Full bloom or beyond	4.67 - 8	100 - 300	<ul style="list-style-type: none"> <li>• Allow 7 or more days after treatment before tillage.</li> </ul>

WEED	APPLICATION			COMMENTS
	GROWTH STAGE	RATE (L/ha)	WATER VOLUME (L/ha)	
<b>Common Milkweed*</b>	Bud to full bloom (preharvest)	1.67	50 – 100	<ul style="list-style-type: none"> <li>• See “<b>Preharvest Treatment</b>” (section 9.9) for more information.</li> <li>• Allow 7 or more days after treatment before tillage.</li> <li>• Reduced control may occur after full bloom.</li> <li>• Common milkweed may not all be in the correct stage, therefore, repeat treatments may be required.</li> </ul>
	Bud to full bloom	8	100 - 300	
<b>Toadflax</b>	Vegetative Stage (summerfallow)	1.67	50 - 100	<ul style="list-style-type: none"> <li>• Apply in clean water using flat fan nozzles.</li> <li>• Allow 7 or more days after treatment before tillage in summerfallow.</li> <li>• For more information, see “<b>Toadflax Control</b>” (section 8.2.4), or “<b>Preharvest Treatment</b>” (Section 9.9).</li> </ul>
	Bud to full bloom (preharvest)			
<b>Alfalfa</b>	Early bud to full bloom stage	2.47 – 3.33	50 - 300	<ul style="list-style-type: none"> <li>• Allow 5 or more days after treatment before tillage. Use the higher rates when alfalfa populations are high or when heavy grass infestations are also present.</li> <li>• For spring applications and control in minimum tillage systems using a 2,4-D tank mix, see section 8.2.6.</li> </ul>
	Fall applications only			

WEED	APPLICATION			COMMENTS
	GROWTH STAGE	RATE (L/ha)	WATER VOLUME (L/ha)	
<b>Dandelion</b>	< 15 cm	1.67	50 – 100	<ul style="list-style-type: none"> <li>• Allow 3 or more days after treatment before tillage for all rates.</li> <li>• Use the higher rate when infestations are heavy.</li> <li>• Refer to “<b>Dandelion</b>” notes in section 8.2.5 for more information.</li> <li>• Allow 7 or more days after treatment before tillage. For more information, see “<b>Preharvest Treatment</b>” (section 9.9).</li> </ul>
	> 15 cm	2.47 – 3.33	50 – 300	
	Rosette to full bloom (preharvest)	1.67	50 - 100	
<b>Foxtail Barley</b>	Seedling to heading	1.67 – 3.33	50 - 100	<ul style="list-style-type: none"> <li>• Allow a minimum of 1 day after treatment before tillage or seeding.</li> <li>• Use higher rates for larger, more established plants, heavy infestations or if plants are stressed.</li> </ul>
<b>Other Perennials</b> (see listing section 6.2)	Early heading or early bud stage	4.67 - 8	100 - 300	<ul style="list-style-type: none"> <li>• Allow 7 or more days after treatment before tillage.</li> </ul>

\*NOTE: For spot treatment, mix 80 millilitres of product in 5 litres of clean water per 100 m<sup>2</sup> (1.67 – 8 litres per hectare is approximately equivalent to 17 – 80 mL/100m<sup>2</sup>, respectively).

## 8.2 SPECIAL NOTES FOR PERENNIAL WEED CONTROL

### 8.2.1 QUACKGRASS

For **season-long control on fall tilled ground**: Apply 1.67 litres per hectare of this product in spring prior to seeding. Apply in 50 to 100 litres per hectare of clean water as described in the preceding table. Delay application until the majority of quackgrass plants have 4 to 5 green leaves. This stage usually occurs 1 to 4 weeks later on fall tilled ground than on undisturbed ground. Reduced control may result on ground tilled deeper than 15 centimetres.

2020-1550  
2020-08-04

**NOTE:** This treatment will provide season-long control of quackgrass on fall tilled ground. Reduced control will be experienced versus this product on non-fall tilled ground. Repeat treatments may be necessary.

**Applications on forages** should be followed by tillage 3 days or later and should be made when good growing conditions exist.

**If a frost has occurred**, wait several days to determine if the quackgrass has recovered. Quackgrass can be treated after a mild frost provided there are 3 to 4 green leaves actively growing at the time of application. Do not apply after the first damaging frost in the fall.

### **8.2.2 SURFACTANT INFORMATION**

The following is a list of approved surfactants for use with POWERMAX HERBICIDE for control of quackgrass:

Agral 90	Companion
Ag Surf	

**Always refer to surfactant label for specific instructions regarding use of that product.**

### **8.2.3 CANADA THISTLE**

**Control of Canada Thistle at the rosette stage:** to ensure the proper timing of application the following steps must be followed:

1. Conduct summerfallow tillage as usual and perform the last tillage operation between July 15<sup>th</sup> and August 1<sup>st</sup>.
2. Allow the thistles to regrow for a minimum of 5 weeks until they are a minimum of 15 centimetres in diameter and in the rosette stage of growth.

**NOTE:** Canada thistle can be treated after a mild frost provided the leaves are still green and actively growing at the time of application. Do not apply after the first damaging frost in the fall.

2020-1550

2020-08-04

## **POWERMAX HERBICIDE PLUS BANVEL II HERBICIDE TANK MIXTURES**

For control of Canada thistle (and perennial sow thistle) in summerfallow or in postharvest stubble, apply 1.13 litres per hectare POWERMAX HERBICIDE plus 1.25 litres per hectare Banvel II Herbicide in 100 – 200 litres per hectare of clean water. In addition, add 350 millilitres per hectare of a non-ionic surfactant registered for use with this product, such as Agral 90, Ag-Surf or Companion.

For best results in summerfallow, cultivate in the spring and apply when the majority of thistles are 15 centimetres to 25 centimetres tall and before the bud stage. Cultivate 3 weeks after application.

In postharvest stubble, apply this tank mixture to actively growing thistles at least 2 weeks prior to a damaging frost.

**NOTE:** Grow only cereals, canola (including rapeseed), soybeans, field corn, sweet corn, or white beans after application of this tank mixture.

If application is made after September 1<sup>st</sup>, or if soil moisture levels are extremely low after application, crop injury may occur in the spring following application.

### **8.2.4 TOADFLAX**

#### **Control of Toadflax in a Summerfallow Vegetative Stage**

To ensure the proper timing of application, the following steps must be followed:

1. Conduct summerfallow tillage as usual and perform the last tillage operation between July 10<sup>th</sup> to July 21<sup>st</sup>.
2. Allow toadflax to regrow for a minimum of 4 to 5 weeks until they are minimum of 15 centimetres tall and at a lush green vegetative stage.

**NOTE:** Toadflax can be treated after a mild frost provided the leaves are still green and actively growing at the time of application. Do not apply after the first damaging frost.

### **8.2.5 DANDELION**

2020-1550  
2020-08-04

Applications should be made up to and including bloom for best results. Follow-up control measures should be used to manage new dandelions germinating from seed to maintain control throughout the season.

2020-1550

2020-08-04

### **8.2.6 ALFALFA CONTROL WITH 2,4-D TANK MIX**

The addition of 2,4-D may improve alfalfa control in situations where control may be more difficult to obtain, such as in minimum tillage systems where populations are heavy, and with spring applications.

For fall control of established stands of alfalfa, apply 1.67 to 3.33 litres per hectare POWERMAX HERBICIDE and 1.2 to 2.4 litres per hectare of any 500 grams per litre 2,4-D amine or low volatile ester formulation in 100 to 200 litres of water per hectare. (Adjust product rates accordingly for other 2,4-D formulations).

For spring applications, use only the low rate of 2,4-D (i.e., 1.2 litres per hectare) and 1.67 to 3.33 litres per hectare POWERMAX HERBICIDE. Only cereal crops not underseeded to legumes may be planted following spring applications of this tank mix, and a 14 day interval between application and planting is required.

Use the higher POWERMAX HERBICIDE rates when perennial grasses are prevalent.

### **8.2.7 ALL PERENNIAL WEEDS**

**Weed Stages:** Weeds must be at the proper stage for effective control. Refer to “**Perennial Weed Control with POWERMAX HERBICIDE**” (section 8.1).

**Nozzle Type:** For best results with conventional boom equipment apply this product with 50 to 300 litres per hectare of clean water using flat fan nozzles and no more pressure than 275 kPa.

**Rhizome Dormancy:** Reduced control may result if rhizomes have become dormant. Dormancy may occur if soil fertility is low and/or the land has not been tilled for several years.

**Mowing Effects:** Mowing prior to application will reduce effectiveness unless weeds are allowed to regrow to the proper stage before application.

**Tillage Effects:** Fall or spring tillage prior to spring applications and tillage between harvesting and fall applications will reduce the effectiveness on perennial weeds. Follow-up tillage after application should be delayed 5 to 7 days for best results. See “**Weed Control**” tables (sections 7.1 and 8.1) for specific tillage interval for each weed.

**Rainfall Effects:** Heavy rainfall immediately after application may wash the chemical off the foliage and a repeat treatment may be required. Do not apply if rainfall is forecast for the time of application.

2020-1550

2020-08-04

**Regrowth from Germinating Seeds:** This product only controls emerged plants. Repeat treatments or other weed control measures may be required to control weeds regenerating from seeds or other underground parts.

**Frost Effects:** Heavy frosts prior to application may reduce control. Do not apply after the first damaging frost in the fall.

## **9.0 CROPLAND SITUATIONS**

**ALWAYS READ PRECAUTIONS, GENERAL INFORMATION AND MIXING AND APPLICATION SECTIONS (3.0, 4.0 and 5.0) PRIOR TO SPECIFIC APPLICATION INFORMATION IN ANY LABEL SECTION. DO NOT APPLY BY AIR EXCEPT FOR PREHARVEST AERIAL APPLICATION (SECTION 9.9.2).**

This product can be applied as a broadcast spray or spot treatment prior to planting all crops, postharvest to annual crops, preharvest in wheat, barley, oats, canola (rapeseed), flax (including low linolenic acid varieties), lentils, peas, soybeans, dry beans and forages, and in summerfallow. It may also be applied as a broadcast spray in Roundup Ready® Corn 2, Roundup Ready® soybeans or Roundup Ready canola varieties (sections 7.5, 7.6 and 7.7). It may be applied as a directed spray in orchards, vineyards, blueberries and strawberries, and using selective equipment in soy and dry beans, orchards, vineyards, cranberries and strawberries (refer to specific sections below for more information). **For specific instructions on weed control in the following cropping situations, always refer to “Annual and Perennial Weed Control” (sections 7.0 and 8.0) for more information.**

### **9.1 PRIOR TO PLANTING – ALL CROPS**

This product may be applied prior to planting all crops for control of emerged weeds listed on this label. Ensure weeds are at the desired stage at the time of application. This product does not provide preemergent weed control and newly germinating weeds may be a problem in the crop. **APPLY BEFORE SEEDING OR TRANSPLANTING.**

#### **9.1.1 PRIOR TO PLANTING – TANK MIXES\* - SOYBEANS**

**\*TANK MIXES – REFER TO THE RESPECTIVE PRODUCT LABELS WHEN TANK MIXING FOR USE RATES, CAUTIONS/WARNINGS, MIXING INSTRUCTIONS, RE-CROPPING RECOMMENDATIONS AND OTHER DETAILS.**

**WHERE TANK MIX PARTNER LABELS REFER ONLY TO THE OLDER (360 G/L) GLYPHOSATE PRODUCTS, E,G, ROUNDUP ORIGINAL OR ROUNDUP**

2020-1550  
2020-08-04

**TRANSORB, ENSURE THAT THE LABEL RATE IS ADJUSTED TO  
COMPENSATE FOR THIS MORE CONCENTRATED PRODUCT.**

2020-1550

2020-08-04

### **POWERMAX HERBICIDE plus Pursuit Herbicide**

POWERMAX HERBICIDE plus Pursuit Herbicide can be applied prior to or after seeding, but before crop emergence. POWERMAX HERBICIDE will control emerged weeds listed on this label when applied as directed (refer to Annual and Perennial Weed control sections in the POWERMAX HERBICIDE product label). Pursuit Herbicide will control weeds germinating from seed.

ONLY SOYBEANS, WHITE BEANS, KIDNEY BEANS, PROCESSING PEAS, FIELD CORN, SPRING BARLEY, SPRING WHEAT AND WINTER WHEAT MAY BE PLANTED THE SEASON FOLLOWING A PURSUIT APPLICATION. WINTER WHEAT MAY BE PLANTED THE SAME YEAR AS A PURSUIT APPLICATION TO SOYBEANS, BUT NOT EARLIER THAN 100 DAYS AFTER THE APPLICATION.

DO NOT APPLY AFTER CROP EMERGENCE

### **POWERMAX HERBICIDE plus metribuzin (Sencor 75 DF Herbicide, Sencor 500F Flowable Herbicide, Sencor 480F Flowable Herbicide, Sencor Soybean Flowable Herbicide, or Lexone DF Herbicide)**

For burndown and residual control of selected annual weeds taller than 4 cm in soybeans, apply POWERMAX HERBICIDE in tank mix with Sencor 75 DF Herbicide, Sencor 500F Flowable Herbicide, Sencor 480F Flowable Herbicide, Sencor 480 Soybean Flowable Herbicide or Lexone DF Herbicide as a preplant surface or pre-emergence application before crop emergence.

### **POWERMAX HERBICIDE plus Dual Magnum Herbicide or Dual II Magnum Herbicide**

For burndown and residual control of selected annual weeds in soybeans. Apply POWERMAX HERBICIDE in tank mix with Dual Magnum Herbicide or Dual II Magnum Herbicide at 1.15– 1.75 L/ha as a preplant surface (up to 30 days before planting) or pre-emergence application before crop emergence.

Perennial weeds such as quack grass may not be controlled with lower rates of POWERMAX HERBICIDE. Use higher rates of POWERMAX HERBICIDE if perennial weeds are present.

### **POWERMAX HERBICIDE plus Dual Magnum Herbicide or Dual II Magnum Herbicide plus metribuzin (Sencor 75DF Herbicide, Sencor 500F Flowable Herbicide, Sencor 480F Flowable Herbicide, Sencor Soybean Flowable Herbicide or Lexone DF Herbicide)**

2020-1550

2020-08-04

For burndown and residual control of selected annual weeds in soybeans.

Apply as a preplant surface (up to 30 days before planting) or pre-emergence application before crop emergence. Perennial weeds such as quack grass may not be controlled with lower rates of POWERMAX HERBICIDE.

### **POWERMAX HERBICIDE plus Broadstrike Dual Magnum Soybean Herbicide**

Broadstrike Dual Magnum Soybean Herbicide at 1.56 L/ha may be tank mixed with POWERMAX HERBICIDE at 1.7 L/ha for control of existing annual weeds and certain perennial weeds including quack grass. This tank mix may be applied preplant surface or pre-emergence in minimum till or no-till conditions. When mixing, add the Broadstrike Dual Magnum Soybean Herbicide component first.

### **POWERMAX HERBICIDE plus Frontier Herbicide**

For burndown and residual control of selected annual weeds apply POWERMAX HERBICIDE plus Frontier Herbicide preplant surface or pre-emergence.

### **POWERMAX HERBICIDE plus linuron**

For burndown and residual control of selected annual weeds apply POWERMAX HERBICIDE plus linuron after seeding but before crop emergence.

### **POWERMAX HERBICIDE plus Axiom DF Herbicide**

Preplant Surface:

For use in conservation tillage, minimum-tillage or no-tillage crop production systems, when weeds are present at the time of application, apply the Axiom DF Herbicide treatment in tank mixture with POWERMAX HERBICIDE. Apply Axiom DF Herbicide in a minimum of 200 L/ha of total volume.

Preemergence:

POWERMAX HERBICIDE plus Axiom DF Herbicide may be applied to the soil surface as a broadcast spray after planting of the crop, but prior to weed or crop emergence.

**For conservation tillage systems:** Apply this tank mixture in a minimum of 200 L/ha of total volume.

## **9.1.2 PRIOR TO PLANTING – TANK MIXES\* - CORN**

**\*TANK MIXES – REFER TO THE RESPECTIVE PRODUCT LABELS WHEN TANK MIXING FOR USE RATES, CAUTIONS/WARNINGS, MIXING INSTRUCTIONS, RE-CROPPING RECOMMENDATIONS AND OTHER DETAILS.**

2020-1550

2020-08-04

**WHERE TANK MIX PARTNER LABELS REFER TO ONLY TO OLDER (360 G/L) GLYPHOSATE PRODUCTS, E.G. ROUNDUP ORIGINAL OR ROUNDUP TRANSORB, ENSURE THAT THE LABEL RATE IS ADJUSTED TO COMPENSATE FOR THIS MORE CONCENTRATED PRODUCT.**

**POWERMAX HERBICIDE plus Dual Magnum Herbicide or Dual II Magnum Herbicide**

For burndown and residual control of selected annual weeds in corn. Apply POWERMAX HERBICIDE in tank mix with Dual Magnum or Dual II Magnum at 1.25 to 1.75 L/ha as a preplant surface (up to 30 days before planting) or pre-emergence application before crop emergence.

NOTE: The use on corn is for EASTERN CANADA ONLY.

Perennial weeds such as quack grass may not be controlled with lower rates of POWERMAX HERBICIDE. Use higher rates of POWERMAX HERBICIDE if perennial weeds are present.

**POWERMAX HERBICIDE plus Dual Magnum Herbicide or Dual II Magnum Herbicide plus Aatrex Liquid 480 Herbicide**

For burndown and residual control of selected annual weeds in corn. Apply POWERMAX HERBICIDE in tank mix with Dual Magnum Herbicide or Dual II Magnum Herbicide at 1.25 – 1.75 L/ha plus Aatrex Liquid 480 Herbicide at 2.1 - 3.1 L/ha as a preplant surface (up to 30 days before planting) or pre-emergence application before crop emergence.

NOTE: The use on corn is for EASTERN CANADA ONLY.

Perennial weeds such as quack grass may not be controlled with lower rates of POWERMAX HERBICIDE. Use higher rates of POWERMAX HERBICIDE if perennial weeds are present.

**POWERMAX HERBICIDE plus Primextra II Magnum Herbicide**

For burndown and residual control of selected annual weeds in corn apply POWERMAX HERBICIDE plus Primextra II Magnum preplant surface or pre-emergence application before crop emergence. This tank mixture requires the use of a surfactant, either Agral 90 or Ag-Surf. See mixing instructions for more information.

Perennial weeds such as quack grass may not be controlled with lower rates of POWERMAX HERBICIDE. Use higher rates of POWERMAX HERBICIDE if perennial weeds are present.

2020-1550

2020-08-04

### **POWERMAX HERBICIDE plus Fieldstar Herbicide**

For burndown and residual control of selected annual weeds apply POWERMAX HERBICIDE plus Fieldstar Herbicide as a preplant surface or pre-emergence application before crop emergence.

### **POWERMAX HERBICIDE plus Frontier MAX Herbicide**

For burndown and residual control of selected annual weeds apply POWERMAX HERBICIDE plus Frontier Herbicide as a preplant surface or pre-emergence application before crop emergence.

### **POWERMAX HERBICIDE plus Prowl 60 WDG Herbicide**

For burndown and residual control of selected annual weeds apply POWERMAX HERBICIDE plus Prowl 60 WDG herbicide after seeding but before crop emergence.

### **POWERMAX HERBICIDE plus Linuron 400 L herbicide**

For burndown and residual control of selected annual weeds apply POWERMAX HERBICIDE plus Linuron 400 L herbicide after seeding but before crop emergence.

### **POWERMAX HERBICIDE plus Converge Pro Herbicide or Converge 75 WDG Herbicide**

Surface Preplant:

CONVERGE 75 WDG Herbicide can be applied to the soil surface up to 14 days prior to planting. CONVERGE 75 WDG Herbicide must be tankmixed with atrazine when applied as a surface preplant application. When weed growth is present at the time of application, POWERMAX HERBICIDE can be added to the Converge Pro Herbicide or Converge 75 WDG Herbicide + atrazine treatment for burndown control of these weeds. Do not incorporate.

Preemergence:

Converge Pro Herbicide or Converge 75 WDG Herbicide can also be applied after planting to just prior to crop emergence. Atrazine and/or POWERMAX HERBICIDE can be tank mixed with pre-emergent applications of Converge Pro Herbicide or Converge 75 WDG Herbicide .

Apply Converge Pro Herbicide at 165-220 mL per hectare, or Converge 75 WDG Herbicide at 105-140 g per hectare, tankmixed with POWERMAX HERBICIDE at 1.67 L per hectare for burndown control of emerged weeds in all tillage management systems and improved control of established dandelion in zero-tillage management systems. A three-way tankmix of Converge Pro Herbicide or Converge 75 WDG Herbicide +

2020-1550

2020-08-04

atrazine + POWERMAX HERBICIDE can be used to provide residual control of the weeds listed in the Converge Pro Herbicide or Converge 75 WDG Herbicide + atrazine section.

### **POWERMAX HERBICIDE plus Axiom DF Herbicide**

#### **Preplant Surface:**

For use in conservation tillage, minimum-tillage or no-tillage crop production systems, when weeds are present at the time of application, apply the Axiom DF Herbicide treatment in tank mixture with POWERMAX HERBICIDE. Apply Axiom DF Herbicide in a minimum of 200 L/ha of total volume.

#### **Preemergence:**

POWERMAX HERBICIDE plus Axiom DF Herbicide may be applied to the soil surface as a broadcast spray after planting of the crop, but prior to weed or crop emergence.

#### **For conservation tillage systems:**

Apply this tankmix in a minimum of 200 L/ha of total volume.

Sencor and Axiom are registered trademarks of Bayer.

Lexone is a registered trademark of E.I. duPont de Nemours and Company.

Dual, Magnum and Primextra are registered trademarks of Syngenta group company.

Broadstrike and Fieldstar are trademarks of Dow Agrosiences LLC.

Frontier is a registered trademark of BASF Corporation.

## **9.2 POSTHARVEST STUBBLE TREATMENT**

This product may be applied in the fall as a postharvest stubble treatment for control of perennial weeds such as quackgrass and Canada thistle. Allow weeds to regrow to the desired stage (20 to 25 centimetres tall for quackgrass and Canada thistle) before application and ensure they have a high proportion of green colouration. Straw should be removed or evenly spread to allow for proper regrowth and spray coverage. Heavy frosts prior to application may decrease control.

## **9.3 SPOT TREATMENT (IN-CROP)**

This product can be applied as an in-crop spot treatment in barley, corn, oats, soybeans, wheat, strawberry, blueberry, forage grasses and legumes including seed production. Applications should be made using the same rates and at the same growth stages as listed in the “**Weed Control**” tables (sections 7.1 and 8.1) or use a 0.67 percent solution for annual weeds and quackgrass and a 1.34 percent solution for other perennial weeds (a 0.67 percent solution equals 0.67 litres of POWERMAX HERBICIDE in 100 litres of spray solution). 0.67 and 1.34 percent solutions should be applied to wet, but not run-off.

2020-1550

2020-08-04

Applications can be made using a boom sprayer, hose and handgun, or hand sprayer in accordance with instructions in “**Application Equipment**” (section 5.2).

**9.3.1 Grazing Restrictions:** Applications can be made up to heading of small grains, initial pod set on soy and dry beans, silking of corn and emergence of seed heads. The crop in the treated area will be killed. Take care to avoid drift for the same reason. **DO NOT APPLY IF CROP GROWTH HAS ADVANCED BEYOND SEED SET. ALLOW 3 TO 5 DAYS FOR POWERMAX HERBICIDE TO TRANSLOCATE INTO ALL PLANT PARTS BEFORE GRAZING OR HARVESTING TREATED AREAS IN FORAGES.**

#### **9.4 SUMMERFALLOW TREATMENT**

This product, or labeled tank mixtures, may be applied in summerfallow to control weeds listed on this label. Ensure weeds are at the desired growth stage and actively growing at application for best results. Reduced control may result if weeds are drought stressed. Weeds will continue to germinate from seed throughout the growing season. Repeat treatments may be necessary to control later germinating weeds.

#### **9.5 MINIMUM AND ZERO TILLAGE CROPPING SYSTEMS (ALL FIELD CROPS, INCLUDING CEREALS, OILSEEDS, PULSES, FORAGES, CORN AND POTATOES)**

This product may be applied prior to seeding or after seeding, but before crop emergence for control of emerged weeds in minimum and zero tillage cropping systems for all field crops. Applications made too far in advance of seeding may allow weeds to emerge between application and crop emergence, as this product does not provide residual weed control.

##### **Minimum and Zero Tillage Tank Mixtures**

**9.5.1 POWERMAX HERBICIDE plus 2,4-D amine or ester** can be applied prior to seeding or after seeding, but before crop emergence in **wheat, winter wheat, barley and rye**. Refer to “**Annual Weed Control with POWERMAX HERBICIDE Tank Mixtures**” table for information (section 7.2).

**9.5.2 POWERMAX HERBICIDE plus bromoxynil (Pardner)** can be applied prior to seeding or after seeding, but before crop emergence in wheat, barley and oats. Refer to “**Annual Weed Control with POWERMAX HERBICIDE Tank Mixtures**” table for information (section 7.2).

**9.5.3 POWERMAX HERBICIDE plus Pursuit®** can be applied prior to, or after seeding, but before crop emergence in soybeans. POWERMAX HERBICIDE will control emerged weeds listed on this label when applied as directed (refer to “**Annual**

2020-1550

2020-08-04

**and Perennial Weed Control”** section 7.0 and 8.0). Pursuit will control weeds germinating from seed. Add the recommended rates of both products in 100 litres of water per hectare, following the instructions on the Pursuit herbicide label.

**ALWAYS REFER TO THE PURSUIT LABEL FOR FURTHER INFORMATION ON WEEDS CONTROLLED, APPLICATION DIRECTIONS, AND USE PRECAUTIONS. ONLY SOYBEANS, FIELD CORN, SPRING BARLEY, SPRING WHEAT AND WINTER WHEAT MAY BE PLANTED THE SEASON FOLLOWING A PURSUIT APPLICATION. WINTER WHEAT MAY BE PLANTED THE SAME YEAR AS A PURSUIT HERBICIDE APPLICATION TO SOYBEANS, BUT NOT EARLIER THAN 120 DAYS AFTER THE APPLICATION.**

**DO NOT APPLY AFTER CROP EMERGENCE.**

Pursuit is a registered trademark of BASF Agrochemical Products B.V. Netherlands.

**9.5.4 POWERMAX HERBICIDE plus MCPA** can be applied prior to seeding in wheat, barley, rye, oats, corn (field and sweet; MCPA amine only), flax and field peas (MCPA amine only). Refer to “**Annual Weed Control with POWERMAX HERBICIDE Tank Mixtures**” table for information (section 7.2).

**9.5.5 POWERMAX HERBICIDE plus Buctril M®** can be applied prior to seeding in wheat, rye, corn, barley, oats, flax, canary seed and seedling grasses (including brome grass, crested wheatgrass, intermediate wheat grass, slender wheatgrass, tall wheatgrass, Russian wild rye, timothy, orchard grass, creeping red fescue, meadow fescue, meadow foxtail, seedling tall fescue, seedling meadow bromegrass, seedling streambank wheatgrass and reed canary grass. Refer to “**Annual Weed Control with POWERMAX HERBICIDE Tank Mixtures**” table for information (section 7.2).

**9.5.6 POWERMAX HERBICIDE plus MCPA amine** can be applied prior to seeding in lentil and chickpea. Refer to “**Annual Weed Control with POWERMAX HERBICIDE Tank Mixtures**” table for information (section 7.2).

**9.5.7 POWERMAX HERBICIDE plus Express Toss-N-Go Herbicide Or Express Toss-N-Go® Dry Flowable 75% Herbicide** in pre-seed situations, wheat and barley may be seeded after a minimum of 24 hours after application. Refer to “**Annual Weed Control with POWERMAX HERBICIDE Tank Mixtures**” table for information (section 7.2).

**ALWAYS REFER TO THE EXPRESS® TOSS-N-GO HERBICIDE OR EXPRESS TOSS-N-GO DRY FLOWABLE 75% HERBICIDE LABEL FOR FURTHER INFORMATION ON APPLICATION DIRECTIONS, TANK MIXING, AND USE PRECAUTIONS.**

2020-1550

2020-08-04

**9.5.8 POWERMAX HERBICIDE plus Banvel II Herbicide can be applied prior to seeding in** wheat, barley, rye, oats and field corn only (do not apply prior to seeding sweet corn). **Refer to “Annual Weed Control with POWERMAX HERBICIDE Tank Mixtures” table for information (section 7.2).**

## **9.6 FORAGES LEGUMES AND GRASSES**

This product may be applied for control of emerged weeds prior to emergence of forage legumes and grasses. If the forages are to be under-seeded with a cover crop, this product must be applied prior to planting the cover crop.

## **9.7 PASTURE RENOVATION**

Use this product to control or suppress existing vegetation for zero-tillage seeding of legumes into established sod for pasture renovation. Delay spraying until weed growth is at least 20 centimetres in height and a maximum number of seedlings or shoots have emerged. Application can be made immediately before, during or after seeding, but before crop emergence.

## **9.8 FORAGE SEED PRODUCTION**

For spot treatment control of perennial weed problems such as quackgrass and Canada thistle in seed fields, apply as directed to vegetation that is at least 20 to 25 centimetres in height but before emergence of seed head. The crop in the treated areas will be killed. Take care to avoid drift outside target areas for the same reason.

## **9.9 PREHARVEST TREATMENT**

### **CONTROL OF QUACKGRASS, CANADA THISTLE, MILKWEED, TOADFLAX AND DANDELION; SEASON-LONG CONTROL OF PERENNIAL SOW THISTLE, AND HARVEST MANAGEMENT**

For control of quackgrass, Canada thistle, common milkweed, toadflax and dandelion; and season-long control of perennial sow thistle, POWERMAX HERBICIDE can be applied prior to harvest of wheat, barley (including malting barley), oats, canola (rapeseed) (including Roundup Ready® varieties), flax (including low linolenic acid varieties), lentils, peas, dry beans, soybeans (including Roundup Ready® varieties) and forages. **DO NOT** apply to crops if grown for seed production.

This treatment may also provide harvest management benefits, by drying down crop and weed vegetative growth, for example, where late flushes of annual weeds, green vegetative crop growth, or late tillering may interfere with harvest operations.

2020-1550  
2020-08-04

EXTREMELY COOL, WET AND/OR CLOUDY WEATHER CONDITIONS BETWEEN THE TIME OF APPLICATION AND THE ANTICIPATED HARVEST DATE MAY SLOW DOWN ACTIVITY OF THIS PRODUCT, THEREBY DELAYING CROP DRYDOWN AND HARVEST DATE. Preharvest treatment to Roundup Ready varieties of canola and soybean provides weed control only.

POWERMAX HERBICIDE should be applied preharvest at 1.67 litres per hectare in 50 to 100 litres per hectare of clean water, by ground application only. Apply only when the crop has 30 percent or less grain moisture content. This stage typically occurs 7 to 14 days before harvest. For forage crops, apply this product at 1.67 to 3.33 litres per hectare 3 to 7 days prior to the last cut before rotation or forage renovation. Consult the table “**Guidelines for Timing of Preharvest Applications**” (section 9.9.1) for visual indicators of this stage in each crop. For the best weed control results, quackgrass should be actively growing and have at least 4 to 5 green leaves. Canada thistle and perennial sow thistle should be actively growing and at or beyond the bud stage for best results. Common milkweed should be at the bud to bloom stage and actively growing for best results. Applications for weed control (not for harvest management) must be made at the correct stage of both weed and crop growth.

Apply only during the period 7 to 14 days (or 3 to 7 days for forage applications) before harvest to ensure best weed control and to maximize harvest management benefits. Earlier application may reduce crop yield and/or quality, and may lead to excess glyphosate residues in the crop.

DO NOT APPLY BY AIR.

### 9.9.1 GUIDELINES FOR TIMING OF PREHARVEST APPLICATIONS

CROP(S)	PERCENT GRAIN MOISTURE	VISUAL SYMPTOMS
WHEAT/BARLEY/OATS	Less than 30	Hard dough stage; a thumbnail impression remains on seed.
CANOLA (including Roundup Ready varieties)	Less than 30	Pods are green to yellow; most seeds are yellow to brown.
FLAX (INCLUDING LOW LINOLENIC ACID VARIETIES)	Less than 30	Majority (75% - 80%) of bolls are brown.
PEAS	Less than 30	Majority (75% - 80%) of pods are brown.
LENTILS	Less than 30	Lowermost pods (bottom 15%) are brown and seeds rattle.
DRY BEANS	Less than 30	Stems are green to brown in colour; pods are mature (yellow to brown in colour); 80% - 90% leaf drop (original leaves).
SOYBEANS (including Roundup Ready varieties)	Less than 30	Stems are green to brown in colour; pod tissue is dry and brown in appearance; 80% - 90% leaf drop.

2020-1550  
2020-08-04

<b>CROP(S)</b>	<b>PERCENT GRAIN MOISTURE</b>	<b>VISUAL SYMPTOMS</b>
FORAGES	Not applicable	Normal stage for forage harvesting.

**NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS: (PREHARVEST TREATMENT OF CHICKPEA, DRIED LUPIN AND DRIED FAVA BEAN).**

The DIRECTIONS FOR USE for this product described below were developed by persons other than Bayer CropScience and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program. Bayer CropScience itself makes no representation or warranty with respect to performance (efficacy) and/or crop tolerance (phytotoxicity) claims for this product when used on the crop listed below.

Accordingly, the Buyer and User assume all risks related to performance and crop tolerance liability arising, and agree to hold Bayer CropScience harmless from any claims based on efficacy and/or phytotoxicity in connection with the uses described below.

**DIRECTIONS FOR USE**

**Preharvest Treatment of Chickpea, Dried Lupin and Dried Fava Bean**

For control of quackgrass, Canada thistle, common milkweed, toadflax and dandelion; and season-long control of perennial sow thistle and harvest management, POWERMAX HERBICIDE can be applied prior to harvest of chickpea, dried lupin and dried fava bean. DO NOT apply to crops if grown for seed production.

POWERMAX HERBICIDE should be applied preharvest at 1.67 litres per hectare in 50 to 100 litres per hectare of clean water, by ground application only. Apply only when the crop has 30 percent or less grain moisture content. This stage typically occurs 7 to 14 days before harvest. For further information see guidelines above. The Pre-harvest interval is 7 days.

**GUIDELINES FOR TIMING OF PREHARVEST APPLICATIONS**

<b>CROP(S)</b>	<b>PERCENT GRAIN MOISTURE</b>	<b>VISUAL SYMPTOMS</b>
Chickpea	Less than 30	Stems are green to brown in colour; pods are mature (yellow to brown in colour); 80%-90% leaf drop (original leaves)
Dried Lupin		
Dried Fava Bean		

--	--	--

**ALWAYS REFER TO THE PRODUCT LABEL FOR FURTHER INFORMATION ON WEEDS CONTROLLED, APPLICATION DIRECTIONS, AND USE PRECAUTIONS**

## 9.9.2 PREHARVEST AERIAL APPLICATION

Refer to the general guidelines for aerial application in Sections 5.2 and 5.3 as well as specific instructions in this section.

### **RESTRICTED USE**

#### **AERIAL PREHARVEST APPLICATION PRAIRIE PROVINCES ONLY (including PEACE RIVER REGION OF B.C.)**

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

**NATURE OF RESTRICTION:** This product is to be used only in the manner authorized. For use only by aerial applicators and aerial application services approved by the provincial regulatory agency to apply this product with aerial application equipment. To qualify for consideration of provincial approval, the following requirements must be demonstrated to the provincial regulatory agency:

1. Aircraft used in the application of this product must have been configured and calibrated to acceptable standards at a recognized calibration (patterning) clinic within 20 months of the date of application. The spray system must not have been subjected to major changes (new nozzles, booms or configurations) since the calibration, and must meet critical drift management standards e.g. maximum boom width 65% of wing span; nozzle type, size and orientation to minimize drift and deliver droplet size VMD in the coarse (400 – 600 microns) or very coarse (600 – 1000 microns) range.
2. Aircraft used in the application of this product must carry a minimum of \$25,000 drift insurance in addition to any provincial requirements for general comprehensive insurance coverage.

3. Applicators using this product must have successfully completed a POWERMAX Herbicide aerial application training course provided by Bayer CropScience Inc.
4. Aerial application services applying this product must employ on staff at least one pilot applicator with at least 250 hours of actual aerial application time and a minimum of 100 hours within the last 24 month period. All pilots who do not meet the minimum experience standard must work under the *direct daily supervision* of a qualified pilot.

Refer to general directions and precautions concerning aerial application, sections 5.2, and 5.3, Buffer Zones.

### DIRECTIONS FOR USE

POWERMAX HERBICIDE may be applied with aerial application equipment for control of quackgrass, Canada thistle, common milkweed, toadflax and dandelion, and season-long control of perennial sow thistle. POWERMAX HERBICIDE can be applied prior to harvest of wheat, barley (including malting barley), oats, canola (rapeseed), flax (including low linolenic acid varieties), lentils, peas, dry beans and soybeans. Do not use on forages. **DO NOT apply to any crops if grown for seed production.**

This treatment may also provide harvest management benefits, by drying down crop and weed vegetative growth, for example, where late flushes of annual weeds, green vegetative crop growth, or late tillering may interfere with harvest operations.

EXTREMELY COOL, WET AND/OR CLOUDY WEATHER CONDITIONS BETWEEN THE TIME OF APPLICATION AND THE ANTICIPATED HARVEST DATE MAY SLOW DOWN ACTIVITY OF THIS PRODUCT, THEREBY DELAYING CROP DRYDOWN AND HARVEST DATE.

POWERMAX HERBICIDE should be applied at 1.67 L/ha in 20 – 50 L/ha of clean water with aerial application equipment. Apply only when the crop has 30% of less grain moisture content. This stage typically occurs 7 to 14 days before harvest. Consult the table “**Guidelines for Timing of Preharvest Applications**” (Section 9.9.1) for visual indicators of this stage in each crop. For the best weed control results quackgrass should be actively growing and have at least 4 to 5 green leaves. Canada thistle and perennial sow thistle should be actively growing and at or beyond the bud stage for best results. Common milkweed should be at the bud to bloom stage and actively growing for best results. Applications for weed control (not for harvest management) must be made at the correct stage of both weed and crop growth.

2020-1550  
2020-08-04

Apply only during the period 7 – 14 days before harvest to ensure best weed control and to maximize harvest management benefits. Earlier application may reduce crop yield and/or quality, and may lead to excess glyphosate residues in the crop.

## 9.10 TREE PLANTINGS

### SHELTERBELTS AND NURSERY STOCK (WOODY ORNAMENTALS)

This product may be used to control listed annual or perennial weeds prior to planting, or as a post directed spray in established nurseries or shelterbelts of the following species:

#### DECIDUOUS

**Ash**

*Fraxinus spp.*

**Caragana**

*Caragana spp.*

**Cherry**

*Prunus spp.*

**Elm**

*Ulmus spp.*

**Lilac**

*Syringa spp.*

**Maple**

*Acer spp.*

**Mountain Ash**

*Sorbus spp.*

**Poplar**

*Populus spp.*

**Russian Olive**

*Elaeagnus spp.*

**Willow**

*Salix spp.*

#### CONIFEROUS

**Fir**

*Abies spp.*

**Juniper**

*Juniperus spp.*

**Pine**

*Pinus spp.*

**Spruce**

*Picea spp.*

**Yew**

*Taxus spp.*

**NOTE:** This product is not recommended for use as an over-the-top broadcast spray in forest tree nurseries or in Christmas tree plantations. Application in such sites should be limited to directed sprays. DO NOT treat Christmas tree plantations in the year of anticipated harvest.

## 9.11 TREE, VINE, BERRY AND OTHER CROPS

This product is recommended for annual and perennial weed control in established vineyards or orchards, in blueberry, cranberry and strawberry, or for site preparation prior to transplanting tree and vine crops. Applications may be made with boom equipment, shielded sprayers, hand held and high volume orchard guns, or with wiper applicator equipment (orchards, vineyards, cranberry and strawberry only). See “**Mixing and Application Equipment Information**” (section 5.2) and the following table for specific information on the use of equipment.

2020-1550  
2020-08-04

Repeat treatments may be necessary to control weeds originating from underground parts of untreated weeds or from seeds. This product does not provide residual or pre-emergent weed control. For subsequent weed control, follow a program using residual herbicides or use repeated applications of this product. Do not apply more than 23 litres of this product per hectare per year.

**EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE SOLUTION, SPRAY, DRIFT, OR MIST WITH FOLIAGE OR GREEN BARK OF TRUNK, BRANCHES, SUCKERS, FRUIT, CANES OF BLUEBERRY BUSHES, OR OTHER PARTS OF TREES OR VINES. CONTACT OF THIS PRODUCT WITH OTHER THAN MATURED BROWN BARK CAN RESULT IN SERIOUS CROP DAMAGE.**

Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed or cut and have not been allowed to regrow to the recommended stage for treatment.

#### **WEED CONTROL IN TREE, VINE, BERRY AND OTHER CROPS**

<b>CROP</b>	<b>RATE (L/ha)</b>	<b>PRE-HARVEST INTERVAL (days)</b>	<b>MAX. APPL. PER YEAR</b>	<b>WEEDS CONTROLLED</b>	<b>COMMENTS (Refer to sections 7.1 and 8.1 for specific rates for weed control)</b>
Apples, Apricot, Cherry (sweet/sour), Peaches, Pears, Plums	1.5 - 8	30	3	Annual and perennial weeds	
Apples, Grapes	Tank Mix 1.5 – 8 + simazine 2.0 – 4.5 kg ai/ha	-	1	Annual and perennial weeds	<ul style="list-style-type: none"> <li>• Will provide season-long preemergent control.</li> <li>• Do not apply to coarse, sandy or gravelly soil.</li> <li>• Use according to the more restrictive label direction for each product in the mix.</li> <li>• DO NOT apply to orchards or vineyards that have been established less than 1</li> </ul>

<b>CROP</b>	<b>RATE (L/ha)</b>	<b>PRE- HARVEST INTERVAL (days)</b>	<b>MAX. APPL. PER YEAR</b>	<b>WEEDS CONTROLLED</b>	<b>COMMENTS (Refer to sections 7.1 and 8.1 for specific rates for weed control)</b>
					or 3 years, respectively. • Simazine rate is equivalent to 2.25 – 5.0 kg/ha Princep® Nine-T®, or 4.0 – 9.0 kg/ha Simadex®
Grapes	1.5 - 8	14	3	Annual and perennial weeds.	<ul style="list-style-type: none"> <li>• Remove all sucker growth from the spray zone before spraying, except for the Concord variety of grape.</li> <li>• Suckering should be conducted within 2 weeks prior to application.</li> <li>• Do not apply to vines which have been established less than 3 years.</li> </ul>
Highbush (cultivated) blueberry	1.87 – 3.73	30	1	Quackgrass	• Use as a directed spray, with no more than 275 kPa pressure.
Lowbush blueberry	0.67 – 1.34% solution (spot application)	Apply in non-bearing year only	1	Woody brush (section 6.3)	<ul style="list-style-type: none"> <li>• Apply as a directed spray in mid-summer of the vegetative (non-bearing) year.</li> <li>• See section 9.3 for instructions on spot treatments.</li> </ul>
Filberts, Hazelnut (established plantations)	1.5 – 2.33	14	-	Annual Weeds	• Use as a directed spray, with no more than 275 kPa pressure.
Walnut, Chestnut, Japanese Heartnut	1.5 - 8	-	2	Annual and perennial weeds	• Apply late spring and fall, postharvest but prior to a damaging frost.

<b>CROP</b>	<b>RATE (L/ha)</b>	<b>PRE- HARVEST INTERVAL (days)</b>	<b>MAX. APPL. PER YEAR</b>	<b>WEEDS CONTROLLED</b>	<b>COMMENTS (Refer to sections 7.1 and 8.1 for specific rates for weed control)</b>
					<ul style="list-style-type: none"> <li>• Apply in 200 – 300 L water as a directed spray, using no more than 275 kPa pressure.</li> <li>• Apply alternatively as a 1.34% wiper solution (see “<b>Wiper Applications</b>” section 9.12).</li> </ul>
Cranberry	13.4% solution (0.62 L POWERMAX HERBICIDE + 4L water)	30	1	Annual and perennial weeds	<ul style="list-style-type: none"> <li>• Apply using wick or wiper applicators (section 9.12).</li> </ul>
Strawberry	0.67 – 1.34% solution (spot application) 22% solution (wiper application)	30	1	Emerged perennial weeds	<ul style="list-style-type: none"> <li>• Apply when weeds are at a susceptible growth stage (see sections 8.1 and 8.2).</li> <li>• See section 9.3 for instructions on spot treatments.</li> <li>• See section 9.12 for instructions on wiper applications.</li> </ul>
Sugar Beets	0.67 – 1.34% solution (spot application)	Treated crop <b>MUST NOT</b> be harvested	1	Dodder species	<ul style="list-style-type: none"> <li>• Apply when dodder is vigorously growing but before flowering.</li> <li>• See section 9.3 for instructions on spot treatments.</li> </ul>
Asparagus	0.83 – 1.67	7	1	Fall seeded ryegrass	<ul style="list-style-type: none"> <li>• Apply in spring before emergence of crop shoots.</li> </ul>

Princep and Nine-T are registered trademarks of Syngenta group company.  
Simadex is a registered trademark of Bayer.

2020-1550  
2020-08-04

**SHORT ROTATION INTENSIVE CULTURE (SRIC) POPLAR (*Populus spp*)**

DO NOT APPLY BY AIR.

This product may be used to control listed annual or perennial weeds prior to planting, or as a post directed spray in established crops of short rotation intensive culture (SRIC) Poplar species (*Populus spp.*)

**EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF HERBICIDE SOLUTION, SPRAY, DRIFT, OR MIST WITH FOLIAGE OR GREEN BARK OF TRUNK, BRANCHES, OR OTHER PARTS OF TREES. CONTACT OF THIS PRODUCT WITH OTHER THAN MATURED BROWN BARK CAN RESULT IN SERIOUS CROP DAMAGE.**

Reduced control may result when applications are made to annual or perennial weeds that have been mowed, grazed or cut and have not been allowed to regrow to the recommended stage for treatment.

POWERMAX HERBICIDE may be applied prior to planting or as a post directed spray in established short rotation intensive culture crops. Apply POWERMAX HERBICIDE up to 8 L/ha in 50 – 100 liters or 150 – 300 L/h for quackgrass control by ground application only. Applications can be made 1-3 times per year during establishment however, not to exceed the limit of 8 L/ha per year. Shielded sprayers must be utilized when applying post directed spray solutions. Allow a 6-8 week interval between spray applications. Apply to actively growing weeds.

**NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR INDICATED SPECIAL USE APPLICATIONS: (NORTH AMERICAN GINSENG).**

The DIRECTIONS FOR USE for this product for the uses described below were developed by persons other than Bayer CropScience and accepted for registration by Health Canada under the User Requested Minor Use Label Expansion program. Bayer CropScience itself makes no representation or warranty with respect to performance (efficacy) and/or crop tolerance (phytotoxicity) claims for this product when used on the crop listed below.

Accordingly, the Buyer and User assume all risks related to performance and crop tolerance arising, and agree to hold Bayer CropScience harmless from any claims based on efficacy and/or phytotoxicity in connection with the uses described below.

**DIRECTIONS FOR USE**

**ALWAYS REFER TO THE PRODUCT LABEL FOR FURTHER INFORMATION ON WEEDS CONTROLLED, APPLICATION DIRECTIONS, AND USE PRECAUTIONS.**

**NORTH AMERICAN GINSENG**

**New Gardens (British Columbia only):** Apply this product in the fall after seeding but before freeze-up in new gardens only to control volunteer cereals. Apply when weeds are at the growth stages listed on the product label. Use a single application of 1.67 litres per hectare in 50 to 100 litres water per hectare. **DO NOT USE A FALL APPLICATION IN ESTABLISHED/EXISTING GARDENS.**

**Existing/Established Gardens:** Apply this product in the spring before the crop has emerged above the soil. Apply when weeds are at the growth stages described in the product label. A maximum of two 1.67 litres per hectare applications in 50 to 100 litres water per hectare may be made in a season. **DO NOT USE A FALL APPLICATION IN ESTABLISHED/EXISTING GARDENS.**

**9.12 SELECTIVE EQUIPMENT**

**WIPER APPLICATORS**

This product may be applied with a wiper applicator, after dilution and thorough mixing with water, to listed weeds in soy and dry beans, grapes, orchards, cranberries, lowbush blueberries and strawberries. Applications must be made before initial pod set in soy and dry beans. (It may also be used in any industrial, tree planting and non-crop site specified on this label. See sections 9.10 and 10.1).

A wiper applicator applies the herbicide solution onto weeds by rubbing the weed with an absorbent material containing the herbicide solution. Wiper applicators include either roller or wick devices which physically wipe appropriate concentrations or amounts of this product directly onto the weed. Equipment must be designed, maintained and operated to prevent the herbicide solution from contacting desirable vegetation. Performance may be improved by reducing speed in areas of heavy weed infestations to insure adequate wiper saturation. Best results may be obtained if 2 applications are made in opposite directions.

**AVOID CONTACT WITH DESIRABLE VEGETATION.** Contact of the herbicide solution with desirable vegetation may result in damage or destruction. Applicators used above desired vegetation should be adjusted so that wiper contact point is at least 5 centimetres above the desirable vegetation. Droplets or foam of the herbicide solution settling on desirable vegetation may result in discoloration, stunting or destruction.

2020-1550

2020-08-04

Applications should be made when the weeds are a minimum of 15 centimetres above the desirable vegetation. Best results may be obtained when more of the weed is exposed to the herbicide solution. Weeds not contacted by the herbicide solution will not be affected. This may occur in dense clumps, severe infestations, or when the height of the weeds varies so that not all weeds are contacted. In these instances, repeat treatments may be necessary. See the “**Weed Control**” tables (sections 7.1 and 8.1) for recommended stage of growth for specific weeds.

## NOTES

- **Maintain equipment in good operating condition. Avoid leakage or dripping onto desirable vegetation.**
- **Adjust height of applicator to insure proper contact with weeds.**
- **Keep wiping surfaces clean.**
- **Maintain recommended roller RPM on roller applicators while in use.**
- **Keep wiper material at proper degree of saturation with herbicide solution.**
- **DO NOT use wiper equipment when weeds are wet.**
- **DO NOT operate equipment at ground speeds below 4 and greater than 10 kilometres per hour. Weed control may be affected by speed of application equipment. As weed density increases, reduce equipment ground speed to insure good coverage of weeds.**
- **Be aware that on sloping ground the herbicide solution may migrate, causing dripping on the lower end and drying on the upper end of the wiper applicator.**
- **Variation in equipment design may affect weed control. With wiper applicators, the wiping material and its orientation must allow delivery of sufficient quantities of the recommended herbicide solution directly to the weed.**
- **Care must be taken with all types of wipers to insure that the absorbent material does not become over-saturated, causing the herbicide to drip onto desirable vegetation.**
- **With all equipment, drain and clean wiper parts immediately after using this product, by thoroughly flushing with water.**

2020-1550  
2020-08-04

**For Roller Applicators** – Mix 0.33 to 0.67 litres of this product in 10 litres water to prepare a 3 to 7 percent solution. Roller speed should be maintained at 50 to 150 RPM.

**For Wick or other Wiper Applicators** – Mix 0.57 litres of this product in 2 litres of water to prepare a 22 percent solution.

## 10.0 NON-CROPLAND USES

### INDUSTRIAL, RIGHTS-OF-WAY, RECREATIONAL, AND PUBLIC AREAS.

**ALWAYS READ PRECAUTIONS, GENERAL INFORMATION AND MIXING AND APPLICATION SECTIONS (3.0, 4.0 AND 5.0) PRIOR TO SPECIFIC APPLICATION INFORMATION IN ANY LABEL SECTION.**

This product can be used to control annual and perennial weeds and woody brush and trees listed on this label in non-crop areas such as railroad, pipeline, highway, power and telephone rights-of-way, petroleum tank farms and pumping installations; roadsides; storage areas; lumberyards; fence rows; industrial plant sites; parking areas; school yards, parks, golf courses, other public areas; airports and similar industrial or non-crop areas.

**NOTE:** For all industrial, rights-of-way, recreational and public areas, repeat treatments may be necessary to control regeneration or new growth.

When applied as recommended under the conditions described, this product will control weeds in non-cropland areas as listed in the following table.

### 10.1 WEED CONTROL IN NON-CROPLAND AREAS WITH POWERMAX HERBICIDE

WEEDS	GROUND APPLICATION			COMMENTS
	BOOM APPLICATION		HAND HELD HIGH VOLUME APPLICATION % SOLUTION	
	RATE* (L/ha)	WATER VOL.* (L/ha)		
<b>Annual grasses and broadleaves</b>	1.5–2.33	50-100	0.67	• Actively growing weeds.
<b>Perennial Weeds</b>				• Actively growing weeds.
Quackgrass	1.67 3.17-4.67	50-300 50-300	0.67 1.34	• Add 0.5% v/v of a recommended surfactant when using water volumes greater than 150 L (see section 8.2.2).
Canada Thistle (bud stage)	3.17-4.67	100-300	1.34	

2020-1550  
2020-08-04

WEEDS	GROUND APPLICATION			COMMENTS
	BOOM APPLICATION		HAND HELD HIGH VOLUME APPLICATION % SOLUTION	
	RATE* (L/ha)	WATER VOL.* (L/ha)		
Purple Loosestrife	4	300-600	0.67-1.34 (or 22% for wiper application)	<ul style="list-style-type: none"> <li>• Higher rate for long term control and for heavy infestations.</li> <li>• See section 10.2.3 for instructions on purple loosestrife applications.</li> <li>• Summer through fall is optimum.</li> </ul>
Other Perennials	4.67-8	100-300	1.34	
<b>Brush and Trees</b> Birch, Cherry, Poplar, Western Snowberry, Willow	2-4	100-300	0.67-1.34	<ul style="list-style-type: none"> <li>• Summer through early fall (see section 10.2).</li> </ul>
Maple, Raspberry/ Salmonberry, Alder	4	100-300	1.34	<ul style="list-style-type: none"> <li>• Late summer through fall.</li> <li>• Fall is optimum.</li> </ul>
<b>Turf Renovation</b> Annual and perennial weeds	1.67-8	100-300	0.67-1.34	<ul style="list-style-type: none"> <li>• Use higher end of the rate range for perennials.</li> </ul>
<b>Roadside Vegetation</b> (1-2m wide along shoulders) Annual weeds (refer to tank mix sections on product labels for specific weeds controlled)	<b>1)</b> 0.5 – 0.67 + 1.25 – 2.5 L DyCleer® or <b>2)</b> 0.5 – 0.67 + 0.30 L DyCleer + 1.2 L 2,4-D amine 500	25-150	-	<ul style="list-style-type: none"> <li>• Refer to “Annual Weed Control” table (section 7.1) for appropriate product rate for specific weeds.</li> <li>• For 2,4-D amine formulations with a different guarantee, adjust the rate accordingly.</li> <li>• No application to standing water.</li> </ul>
<b>Residual Control</b> Annual and perennial weeds	1.67 – 8 +	200-400	-	<ul style="list-style-type: none"> <li>• Do not apply to coarse, sandy or gravelly soil. One application per year.</li> </ul>

2020-1550  
2020-08-04

WEEDS	GROUND APPLICATION			COMMENTS
	BOOM APPLICATION		HAND HELD HIGH VOLUME APPLICATION % SOLUTION	
	RATE* (L/ha)	WATER VOL.* (L/ha)		
(the simazine component of this tank mixture will provide season long control of most germinating broadleaf weeds and grasses. It may also provide postemergent activity on certain annual weeds).	4.0 -9.0 L Simadex Simazine Flowable			<ul style="list-style-type: none"><li>• Use according to the most restrictive label directions for each product in the mixture.</li><li>• For other simazine formulations registered for industrial/ non-cropland areas, use equivalent rates; i.e., 2.0 – 4.5 kg simazine/ha.</li></ul>

\* For more information on rates, water volumes and application, refer to “**Annual and Perennial Weed Control**” (sections 7.1 and 8.1, respectively).

DyCleer is a registered trademark of Syngenta group company.  
Simadex is a registered trademark of Bayer.

## **10.2 APPLICATION INFORMATION FOR NON-CROPLAND USES**

### **FOLIAR APPLICATIONS**

Spray coverage should be uniform and complete. Do not spray to the point of run-off. Do not allow spray drift to contact desirable vegetation as severe injury or destruction may occur. For woody brush and trees, early season applications may take 30 to 45 days for symptoms to develop on target species. Late season application may be made to species that have some autumn colors provided no major leaf drop has occurred. Control will be observed the following spring.

**EXTREME CARE MUST BE EXERCISED TO AVOID CONTACT OF SPRAY WITH FOLIAGE OF DESIRABLE TURF GRASSES, TREES, SHRUBS, OR OTHER DESIRABLE VEGETATION SINCE SEVERE DAMAGE OR DESTRUCTION MAY RESULT.**

This product does not provide residual weed control. For subsequent weed control, follow a label approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

#### **10.2.1 GROUND APPLICATIONS: For all non-cropland uses**

For woody brush and trees, apply 2 to 4 litres of this product per hectare. Use ground boom or boomless, or mist blower equipment, or apply as a 0.67 to 1.34 percent solution using hand held, high volume equipment. Apply as directed in the recommended volume of clean water to foliage of actively growing vegetation. Use the 4 litres per hectare rate for Maple, Alder and Willow\* species, as well as for hard to control perennial weed species. (\*suppression only).

Spray coverage should be uniform and complete. Do not spray to the point of run-off. Do not allow spray drift to contact desirable vegetation as severe injury or destruction may occur. If weeds have been mowed or tilled, do not treat until regrowth has reached the recommended stages.

#### **10.2.2 PURPLE LOOSESTRIFE CONTROL**

- **DO NOT TREAT PLANTS OVER OPEN WATER. POWERMAX HERBICIDE is not registered for direct application to bodies of water.**
- Treat when plants are actively growing at or beyond the bloom stage. If using hand held equipment, spray-to-wet.
- For wiper applications see section 9.12.

2020-1550

2020-08-04

- Where feasible, remove flower heads before treatment to ensure prevention of seed set.
- For large (>1.6 ha) monocultures of loosestrife, work from the periphery inward in successive years to allow competing vegetation to invade the treated area.
- A long-term control strategy should include measures to control both established plants and seedlings. Sprayed areas should be monitored to determine the appropriate follow-up management. Early detection and treatment of second and third generation seedlings is important to prevent re-infestation of purple loosestrife. Desirable native plant communities will then have a chance to become re-established.

### **10.3 SELECTIVE APPLICATION FOR ALL NON-CROPLAND USES**

Selective equipment such as WIPER and ROLLER applicators can be used to control emerged weeds in non-crop areas and tree plantings. See “**Selective Equipment**” (section 9.12) for more information.

### **10.4 TURF GRASS**

When applied as directed, under conditions described, this product controls most existing vegetation. Apply this product at rates specified in “**Weed Control in Non-Cropland Areas**” (section 10.1).

### **DO NOT DISTURB SOIL OR UNDERGROUND PLANT PARTS BEFORE TREATMENT.**

Where existing vegetation is growing in a field or unmowed situation, apply this product to actively growing weeds at the stages of growth given in “**Weeds Controlled**” (sections 7.1 and 8.1, respectively). Where existing vegetation is growing under mowed turfgrass management, apply this product after omitting at least one regular mowing to allow sufficient growth for good interception of the spray and proper translocation into underground plant parts. Tillage or renovation techniques such as vertical mowing, coring or slicing should be delayed for 7 days after application to allow proper translocation into underground plant parts.

For maximum control of existing vegetation, delay establishment to determine if regrowth from escaped underground plant parts occurs. When repeat treatments are necessary, sufficient regrowth must be attained prior to application. Desirable turfgrass may be established following the above procedures.

2020-1550

2020-08-04

## 10.5 INJECTION APPLICATIONS -- FOR ALL NON-CROPLAND USES

Woody vegetation may be controlled by injection application of this product. Apply using suitable equipment, which must penetrate into living tissue, at a rate of at least 0.33 millilitres (either undiluted or 1:1 with water) per 5 centimetres tree diameter at breast height (DBH). The cuts should be spaced evenly around the tree and below all major branches. Application may be made at any time of year, except when cold temperatures prevent adequate penetration of injection equipment, or in the spring during periods of heavy sap flow. Control of tree species with tree diameters greater than 20 centimetres may not be acceptable at this rate.

Total control may not be evident for 1 to 2 years following treatment.

A partial list of species controlled includes:

### **Alder**

*Alnus spp.*

### **Birch**

*Betula spp.*

### **Cedar**

*Thuja spp.*

### **Cherry**

*Prunus spp.*

### **Douglas Fir**

*Pseudotsuga spp.*

### **Hemlock**

*Tsuga spp.*

### **Maple\***

*Acer spp.*

### **Pine**

*Pinus spp.*

### **Poplar**

*Populus spp.*

### **Willow**

*Salix spp.*

\* This treatment may only provide suppression of Bigleaf Maple. Late fall applications will provide optimum suppression of Bigleaf Maple.

## 10.6 CUT STUMP APPLICATION

Woody vegetation may be controlled by the application of this product to freshly cut stumps to prevent regrowth. Because the treatment uses a concentrated solution, application must be made using low-pressure equipment e.g., squirt bottle or similar device. This product must be applied immediately to the surface of the freshly cut stump i.e., within 5 minutes for optimum control at the prescribed rates. Only the cambial tissues of the cut surface should be treated. Apply the herbicide solution at a rate equivalent to at least 0.33 millilitres product for every 5 centimetres DBH. Do not cover the remaining area nor any exposed roots, as this product does not penetrate bark well. This treatment may be used at any time of year, except during periods of heavy sap flow or when low temperatures prevent solution application due to freezing. A water soluble colourant may be added to the solution as a means of indicating which surfaces have been treated. Total control may not be evident until 1 to 2 years after treatment.

2020-1550  
2020-08-04

See “**Injection Applications**” (section 10.5) of this label for a partial list of species controlled.