

2024-0027  
2024-12-12

## Container base label

# CONCISE<sup>®</sup> PLANT GROWTH REGULATOR

COMMERCIAL

SOLUTION

For greenhouse use only

**Active Ingredient:** Uniconazole -P .....0.055%  
Contains 1,2-Benzisothiazolin-3-one at 0.04 % as a preservative

REGISTRATION NO. 32342 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

**CAUTION  
POISON**

KEEP OUT OF REACH OF CHILDREN

**Manufactured by:**  
**FINE AGROCHEMICALS LIMITED**  
Hill End House, Whittington  
Worcester WR5 2RQ  
United Kingdom  
[www.fine.eu](http://www.fine.eu)

**Distributed by:**  
**N.M. Bartlett Inc.**  
4509 Bartlett Rd.  
Beamsville, Ontario  
L3J 0Z1, Canada  
Tel : 905-563-8261

Net Contents: 1-10 litres

## PRECAUTIONS

- **KEEP OUT OF REACH OF CHILDREN**
- Wash concentrate from skin or eyes immediately
- After use, wash hands and other exposed skin
- Remove contaminated clothing immediately after handling this product. Wash separately from other laundry before reuse
- For mixing, loading, application, clean-up and repair wear waterproof rain gear (e.g. Tyvek coveralls) with a hood, chemical-resistant gloves, goggles, socks, rubber boots and a NIOSH approved organic-vapour-removing cartridge with a prefilter approved for pesticides, or a NIOSH-approved canister approved for pesticides. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Contamination of water in ponds, streams, rivers and areas with desirable vegetation should be avoided

## FIRST AID

**IF SWALLOWED:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth to mouth if possible. Call a poison control centre or doctor for further treatment advice

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

Note to Physicians: Emergency Information – call 1-800-892-0099

**TOXICOLOGICAL INFORMATION:** Treat symptomatically.

## ENVIRONMENTAL PRECAUTIONS

This product demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

## **STORAGE**

Store this product away from food or feed. Keep pesticide in original container. Store the container tightly closed. Do not put concentrate or dilute solutions into food or drink container. Store in cool, dry and well-ventilated place away from seed, plants, fertilizer or other pesticides. Do not store diluted spray.

## **DISPOSAL**

DO NOT reuse this container for any purpose. This is a recyclable container and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

## **NOTICE TO USER**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label.

# CONCISE®

## PLANT GROWTH REGULATOR

COMMERCIAL

SOLUTION

For greenhouse use only

**Active Ingredient:** Uniconazole -P .....0.055%  
Contains 1,2-Benzisothiazolin-3-one at 0.04 % as a preservative

REGISTRATION NO. 32342 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING  
IF ANY LABEL INFORMATION IS MISSING, CONTACT THE DISTRIBUTOR BEFORE  
USING THIS PRODUCT

**CAUTION**  
**POISON**

KEEP OUT OF REACH OF CHILDREN

**Manufactured by:**  
**FINE AGROCHEMICALS LIMITED**  
Hill End House, Whittington  
Worcester WR5 2RQ  
United Kingdom  
[www.fine.eu](http://www.fine.eu)

**Distributed by:**  
N.M. Bartlett Inc.  
4509 Bartlett Rd.  
Beamsville, Ontario  
L3J 0Z1, Canada  
Tel : 905-563-8261

Net Contents: 1-10 litres

## PRECAUTIONS

- **KEEP OUT OF REACH OF CHILDREN**
- Wash concentrate from skin or eyes immediately
- After use, wash hands and other exposed skin
- Remove contaminated clothing immediately after handling this product. Wash separately from other laundry before reuse
- For mixing, loading, application, clean-up and repair wear waterproof rain gear (e.g. Tyvek coveralls) with a hood, chemical-resistant gloves, goggles, socks, rubber boots and a NIOSH approved organic-vapour-removing cartridge with a prefilter approved for pesticides, or a NIOSH-approved canister approved for pesticides. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Contamination of water in ponds, streams, rivers and areas with desirable vegetation should be avoided

## FIRST AID

**IF SWALLOWED:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth to mouth if possible. Call a poison control centre or doctor for further treatment advice

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

Note to Physicians: Emergency Information – call 1-800-892-0099

**TOXICOLOGICAL INFORMATION:** Treat symptomatically.

## ENVIRONMENTAL PRECAUTIONS

This product demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

## **STORAGE**

Store this product away from food or feed. Keep pesticide in original container. Store the container tightly closed. Do not put concentrate or dilute solutions into food or drink container. Store in cool, dry and well-ventilated place away from seed, plants, fertilizer or other pesticides. Do not store diluted spray.

## **DISPOSAL**

DO NOT reuse this container for any purpose. This is a recyclable container and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

## **NOTICE TO USER**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label.

## **DIRECTIONS FOR USE**

### **IMPORTANT NOTES**

- Concise is an extremely active plant growth regulator. Avoid excessive doses by exercising care when diluting and applying the product. Overdosing could result in plant retardation and other adverse effect on flowering, plant appearance and marketability. Avoid spray drift onto non-target species.
- Apply to well watered, vigorously growing plants. Do not mix with other products such as pesticides, fertilizers, or adjuvants.

### **PRECAUTIONS**

- FOR GREENHOUSE USES ONLY
- Do not apply Concise in a manner inconsistent with the recommendations on

the label.

- Do not use in greenhouses with soil floors consisting of sand or sandy loam soil.
- Do not reuse soil from plants treated with Concise .
- Do not use on cranesbill geranium (*Geranium spp.*)
- Concise is recommended to be applied during morning or late afternoon hours.
- Do not allow releases, effluent or runoff from greenhouses containing this product to enter lakes, streams, ponds or other waters.
- Concise retards elongation almost immediately after application. Assessment of rate response can often be made about one week after treatment
- For Poinsettias and Chrysanthemums only, if response is inadequate after the initial application within the recommended rate range, a repeat application can be made.
- Differences in plant response to Concise due to plant surface area in relation to soil surface area are possible
- Within the recommended rates of use, there can be differences in response at extreme high or low temperature rates. A higher rate may be required at high temperatures than at low temperatures to achieve the same effect.
- DO NOT apply using handheld mist blowers/airblast or handheld fogging equipment.
- DO NOT apply to ornamentals grown for cut flowers.

## **GENERAL INFORMATION**

Concise can be used to reduce plant height by limiting internode elongation through the inhibition of gibberellin synthesis. The product retards the growth of Poinsettias, Chrysanthemums, Easter Lilies, Geranium and Bedding Plant Species. Dilute with water according to the directions below. Apply as foliar spray with standard spray equipment.

Effective dose is strongly dependent on the application volume. Rates expressed are based on a spray volume of 2 litres applied uniformly to 10 square metres of bench area. For best response, use low pressure and larger pressure droplets.

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Fine Agrochemicals Ltd at 888-474-3463 for information before applying any tank mix that is not specifically recommended on this label.

Fine Agrochemicals Ltd has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) of all potential tank-mixes under all environmental conditions or for all crop varieties. Tank mixes that are not specifically listed on this label should be tested on a small area first, under local conditions and using standard practices, to confirm the tank mix is suitable for widespread application.

Cultural and environmental factors such as growing media, water/fertiliser management, temperature, light, relative day and night temperatures, and growing house composition influence the plant response to Concise . First time users of Concise should treat a limited number of plants at the lowest recommended rate. Adjust rates within the recommended rate range as required for best effect. For any given rate, the effect of temperature can be significant. Within the recommended rate range, a higher rate may be required at higher temperatures than at lower temperatures for the same effect.

Concise has been tested on a limited range of cultivars. Plant response to Concise is strongly influenced by genetic background. First time users of Concise should treat a limited number of plants at the lowest recommended rate, observe plant response and make rate adjustments accordingly.

### **POINSETTIA**

Apply Concise as a foliar spray when lateral shoots are 3.5 to 5.0 cm in length (about 10 to 14 days after pinching). Use a rate of 4 to 16 ml/l (2-8 ppm) and apply uniformly at a spray volume of 2 litres per 10 square meters of bench area. Apply one or two treatments at 7 day intervals until the desired growth characteristics are achieved. Bract size and diameter may be slightly reduced with treatment, and days to flowering may be delayed. Multiple applications of the lower label rate may provide more satisfactory response. Applications later in the season (after short-day initiation) may have negative effects on bract size, plant quality, and marketability. **DO NOT APPLY TO POINSETTIAS AFTER INITIATION OF SHORT DAYS.**

### **CHRYSANTHEMUMS**

Concise can be applied to pot Chrysanthemums as well as hardy Chrysanthemums. Use as a foliar spray 7-14 days after pinching when the length of the breaks is 3.5 to 5.0cm. Use a rate of 10 -20 ml/l (5-10ppm) and apply uniformly in a spray volume of 2 litres per 10 square meter of bench area. A single repeat spray can be applied 7-21 days after the initial application to increase the level of growth reduction.

Multiple applications may have more effect than the same amount of product applied as a single treatment. A solution of 40ml/l (20ppm) can be applied in a spray volume of 2 litres per 10 square meters of bench area, but **DO NOT** make a repeat application following the 20ppm spray

### **EASTER LILIES**

Apply Concise as a foliar spray when shoots are 7.5 cm tall. Use a rate of 6-60ml/l

(3-30 ppm) and apply uniformly in a volume of 2 litres per 10 square metres of bench area. A single repeat spray of 3-5 ppm may be applied 2-3 weeks after initial spray of 3-5 ppm. DO NOT make a repeat application following a spray of greater than 5 ppm. Avoid late-season application. Insufficient height reductions may result following application of Concise to Nellie White Easter lilies grown under zero or negative DIF (day temperatures the same as or lower than night temperatures)

NOTE: Easter lilies that have been treated with Concise may have lower requirement for water.

### GERANIUM AND BEDDING PLANT SPECIES

Apply as a foliar spray.

Geranium and bedding plant species for which Concise is acceptable with stated restrictions

Host species	Acceptable rate (ppm uniconazole-P)*	Acceptable maximum number of applications**	Acceptable application timing*** (plant height)
Bedding plants			
Celosia ( <i>Celosia plumose</i> )	2.5-10	2	5-10 cm
Coleus ( <i>Coleus sp.</i> )	5-10	1	2-5 cm
Impatiens ( <i>Impatiens walleriana</i> or <i>I. balsamina</i> )	2.5-10	2	3-10 cm
Marigold ( <i>Calendula officinalis</i> , <i>Tagetes erectus</i> , <i>T. patula</i> and <i>T. signata</i> )	5-15	2	5-10 cm
Pansy ( <i>Viola wittrockiana</i> )	1-6	1; if >3ppm was applied 2; both applications of 1-3ppm	3-5 cm
Periwinkle ( <i>Vinca minor</i> )	1-3	2	5-10 cm
Petunia ( <i>Petunia</i> hybrid)	10-30	1	2-5 cm
Salvia ( <i>Salvia splendens</i> , <i>S. horminum</i> , <i>S. farinacea</i> , <i>S. coccinea</i> and <i>S. sclarea</i> )	1-10	2	3-10 cm
Snapdragon ( <i>Antirrhinum majus</i> )	10-30	1: if >15 ppm was applied 2: both applications of 10-15 ppm	5-10 cm
Viola ( <i>V. tricolor</i> , <i>V. cornuta</i> and <i>V. williamsii</i> )	1-5	1	2-5 cm

Host species	Acceptable rate (ppm uniconazole-P)*	Acceptable maximum number of applications**	Acceptable application timing*** (plant height)
Geranium (Zonal: <i>Pelargonium hortorum</i> , and Ivy: <i>Pelargonium peltatum</i> )	2-4 if grown from seed; 2-8 if vegetatively propagated	2	2-4 ppm on seed geraniums when plant height is 5 -16 cm.

\*Concise is extremely active at very low rates. Growers are advised to start with lower rates for younger seedlings. Apply only by foliar spray.

\*\* To some crops (as indicated), a second application may be made as required, but no earlier than 7 days after the first application.

\*\*\* The application timing proposed was 5-16 cm height for all proposed bedding plant species. The application timing shown in this column was supported by the submitted data, however, the applicant and registrant may further restrict the application timing, but only within the accepted application window.

## PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

Concise<sup>®</sup> is a registered trademark of Fine Agrochemicals Limited