

(Sleeve)

GROUP 3 FUNGICIDE

HORNET™ 432 F

Foliar Fungicide

SUSPENSION

FOR SUPPRESSION OF *FUSARIUM* HEAD BLIGHT (SCAB) AND CONTROL OF FOLIAR DISEASES ON WHEAT (SPRING, WINTER AND DURUM), BARLEY AND OATS

COMMERCIAL

ACTIVE INGREDIENT: Tebuconazole 432 g/L

Contains 1,2-benzisothiazolin-3-one at 0.048% as a preservative.

REGISTRATION NO. **32500** PEST CONTROL PRODUCTS ACT

READ THE LABEL AND PAMPHLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

CAUTION: EYE IRRITANT

DANGER: SKIN IRRITANT

NET CONTENTS: 4.73 LITRES to 200 LITRES (BULK)

Product Information: 1-800-868-5444

Nufarm Agriculture Inc.
5101, 333 - 96th Ave NE
Calgary, Alberta T3K 0S3

In case of spills, poisoning or fire, use the 24 hour telephone emergency response number 1-800-424-9300

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist.

Wear coveralls over long pants, long-sleeved shirt, chemical resistant gloves and boots during mixing, loading, application, clean-up and repair activities. Coveralls are not required during application within closed cabs or cockpits.

Follow manufacturer's instructions for cleaning/maintaining the Personal Protective Clothing (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

User Safety Recommendations:

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

FIRST AID

In case of poisoning, call physician or Poison Control Centre immediately. Have patient lie down and keep quiet.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

HORNET 432 F Foliar Fungicide

The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.

To Physician: No specific antidote. Treat symptomatically.

EMERGENCY TELEPHONE NUMBERS

For spills or transportation accidents, Chemtrec, 1-800-424-9300

For health or environmental emergencies, ProPharma Group, 1-877-325-1840

For product and use information, Nufarm Agriculture Inc., 1-800-868-5444

ENVIRONMENTAL HAZARDS

Tebuconazole is persistent and will carryover. It is recommended that any products containing tebuconazole not be used in areas treated with this product during the previous season.

Toxic to birds, small wild animals, aquatic organisms, and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE and APPLICATION IN FIELDS ADJACENT TO AQUATIC AREAS. As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests. Do not apply to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff is hazardous to aquatic organisms in neighboring areas. To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body. To reduce the run off refer to the recommendations under "APPLICATION IN FIELDS ADJACENT TO AQUATIC AREAS".

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

LEAK OR SPILL PROCEDURE: Handle and open container in a manner as to prevent spillage. If container is leaking invert to prevent leakage. If the container is leaking or material is spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away. For decontamination procedures or any other assistance that may be necessary, you may contact the 24 hour emergency response number 1-800-424-9300.

DISPOSAL:

Recyclable Container Disposal: Do not reuse this container for any purpose. This is a recyclable container and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

HORNET 432 F Foliar Fungicide

Returnable Container: Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable Container: For disposal, this empty container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not use this container for any other purpose.

Non-Returnable Container:

1. Triple or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.

Disposal of Unused, Unwanted Product:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills.

TM HORNET is a trademark of Nufarm Agriculture Inc.

(Booklet)

GROUP 3 FUNGICIDE

HORNET™ 432 F

Foliar Fungicide

SUSPENSION

FOR SUPPRESSION OF *FUSARIUM* HEAD BLIGHT (SCAB) AND CONTROL OF FOLIAR DISEASES ON WHEAT (SPRING, WINTER AND DURUM), BARLEY AND OATS

COMMERCIAL

ACTIVE INGREDIENT: Tebuconazole 432 g/L

Contains 1,2-benzisothiazolin-3-one at 0.048% as a preservative.

REGISTRATION NO. **32500** PEST CONTROL PRODUCTS ACT

READ THE LABEL AND PAMPHLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

CAUTION: EYE IRRITANT

DANGER: SKIN IRRITANT

NET CONTENTS: 4.73 LITRES to 200 LITRES (BULK)

Product Information: 1-800-868-5444

Nufarm Agriculture Inc.
5101, 333 - 96th Ave NE
Calgary, Alberta T3K 0S3

In case of spills, poisoning or fire, use the 24 hour telephone emergency response number
1-800-424-9300

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

DIRECTIONS FOR USE

IMPORTANT: Read this entire label before using HORNET™ 432 F Foliar Fungicide. **Ensure no bystanders are present during the application operation.** Only protected handlers may be in the area during application. Use mechanical flaggers only. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Spray Volume: HORNET 432 F Foliar Fungicide should be applied in a minimum of 100 litres of spray solution per hectare by ground sprayer or 47 litres of spray solution per hectare by aircraft spray equipment. Check equipment calibration frequently.

Chemigation: Do not apply this product through any type of irrigation system.

Mixing: Add the required amount of HORNET 432 F Foliar Fungicide into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, HORNET 432 F Foliar Fungicide should be thoroughly dispersed prior to the addition of other materials. HORNET 432 F Foliar Fungicide is recommended to be used with a non-ionic surfactant at the surfactant's labeled rate.

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

Airblast application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** direct spray above plants to be treated. Turn off outward pointing nozzles at row ends and outer rows. **DO NOT** apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side.

Aerial application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 km/h at flying height at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. To reduce drift caused by turbulent wingtip vortices, the nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotorspan.

Buffer zones:

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands). In addition to the buffer zones specified in the table below, users must also observe buffer zones specified under APPLICATION IN FIELDS ADJACENT TO AQUATIC AREAS.

| Method of application | Crop | | Buffer Zones (metres) Required for the Protection of: |
|-----------------------|--|-----------------------|---|
| | | | Terrestrial Habitats |
| Field sprayer | Wheat (spring, winter, durum), Barley, Oats | | 1 |
| Aerial | Wheat (spring, winter, durum), Barley, Oats | Fixed and rotary wing | 15 |

The spray drift buffer zones required for the protection of terrestrial habitats for this product can be modified based on weather conditions and spray equipment configuration by accessing the Buffer Zone Calculator on the Pest Management Regulatory Agency web site.

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

HORNET 432 F Foliar Fungicide

| Cereals (Wheat, Barley and Oats): | | | |
|--|---|---|--|
| CROP | DISEASES | DOSAGE OF HORNET 432 F FOLIAR FUNGICIDE | REMARKS |
| Wheat (spring, winter and durum) | For suppression* of <i>Fusarium</i> head blight (scab) (<i>Gibberella zeae</i> / <i>Fusarium graminearum</i>) For control of Septoria glume blotch (<i>Stagonospora nodorum</i>) | 292 mL/ha | * HORNET 432 F Foliar Fungicide should only be applied when the risk of <i>Fusarium</i> head blight infection is high. Consult your local extension authority regarding the need for HORNET 432 Fuse. Head blight is extremely difficult to control. For that reason only suppression of <i>Fusarium</i> head blight (scab) is claimed. <ul style="list-style-type: none"> • <i>Fusarium</i> head blight (scab) outbreaks occur when the weather is warm and wet at the flowering to soft dough stages. • The application of HORNET 432 F Foliar Fungicide for protection against <i>Fusarium</i> head blight (scab) should be considered when these weather conditions are forecasted for this stage of wheat development. • Timing of application is critical: For optimum suppression of <i>Fusarium</i> Head Blight and control of Septoria glume blotch, apply HORNET 432 F Foliar Fungicide within the time period from when at least 75% of the wheat heads on the main stem are fully emerged to when 50% of the heads on the main stem are in flower. • Spray coverage is essential: Ensure thorough coverage of all wheat heads. • Spray equipment must be set up to provide good coverage to wheat heads. • HORNET 432 F Foliar Fungicide may be applied by ground or air equipment • GROUND APPLICATION: Apply specified dosage in a minimum of 100 L of water per hectare. • AERIAL APPLICATION: Apply specified dosage in a minimum of 47 L of water per hectare. • HORNET 432 F Foliar Fungicide is recommended to be used with non-ionic surfactant at the surfactant's labeled rate. |
| | Rusts (Leaf, Stem and Stripe) (<i>Puccinia triticina</i> , <i>P. graminis</i> , <i>P. striiformis</i>) Septoria (Leaf Blotch) (<i>Septoria tritici</i>) Tan Spot (<i>Pyrenophora tritici-repentis</i>) | 220-292 mL/ha | <ul style="list-style-type: none"> • Apply HORNET 432 F Foliar Fungicide to leaf foliage at the first sign or very early stage of disease, especially if weather conditions are conducive to disease development, up to the end of the flowering stage. • Where a rate range is specified, use of the higher rate should be considered when weather conditions are conducive to heavy disease development. • HORNET 432 F Foliar Fungicide may be applied by ground or air equipment • GROUND APPLICATION: Apply specified dosage in a minimum of 100 L of water per hectare. • AERIAL APPLICATION: Apply specified dosage in a minimum of 47 L of water per hectare. • HORNET 432 F Foliar Fungicide is recommended to be used with non-ionic surfactant at the surfactant's labeled rate. |
| | Powdery Mildew (<i>Erysiphe graminis</i>) | 292 mL/ha | |

HORNET 432 F Foliar Fungicide

| | | | |
|--------|---|---------------|---|
| Barley | <p>Net Blotch (<i>Pyrenophora teres</i>)</p> <p>Spot Blotch (<i>Cochliobolus sativus</i>)</p> <p>Scald (<i>Rhynchosporium secalis</i>)</p> <p>Rusts (Leaf, Stem and Stripe) (<i>Puccinia hordei</i>, <i>P. graminis</i>, <i>P. striiformis</i>)</p> <p>Septoria Leaf Blotch (<i>Septoria passerinii</i>)</p> <p>Powdery Mildew (<i>Erysiphe graminis</i>)</p> | 220-292 mL/ha | <ul style="list-style-type: none"> • Apply HORNET 432 F Foliar Fungicide at the very early stages of disease development. • Where a rate range is specified, use of the higher rate should be considered when weather conditions are conducive to heavy disease development. • HORNET 432 F Foliar Fungicide may be applied by ground or air equipment • GROUND APPLICATION: Apply specified dosage in a minimum of 100 L of water per hectare. • AERIAL APPLICATION: Apply specified dosage in a minimum of 47 L of water per hectare. • HORNET 432 F Foliar Fungicide is recommended to be used with non-ionic surfactant at the surfactant's labeled rate. |
| Oats | <p>Crown Rust (<i>Puccinia coronata</i>)</p> <p>Stem Rust (<i>Puccinia graminis</i>)</p> | 220 mL/ha | <ul style="list-style-type: none"> • Apply HORNET 432 F Foliar Fungicide at the very early stages of disease development. • Where a rate range is specified, use of the higher rate should be considered when weather conditions are conducive to heavy disease development. • HORNET 432 F Foliar Fungicide may be applied by ground or air equipment • GROUND APPLICATION: Apply specified dosage in a minimum of 100 L of water per hectare. • AERIAL APPLICATION: Apply specified dosage in a minimum of 47 L of water per hectare. • HORNET 432 F Foliar Fungicide is recommended to be used with non-ionic surfactant at the surfactant's labeled rate. |

Restrictions: A maximum of one application of HORNET 432 F Foliar Fungicide may be applied per crop season to wheat, barley and oats. Applications may not be made within 36 days of harvest. Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with HORNET 432 F Foliar Fungicide. Straw cut after harvest may be fed or used for bedding.

SEQUENTIAL APPLICATIONS IN WHEAT AND BARLEY:

HORNET 432 F Foliar Fungicide may be applied in sequence with Prosaro 250EC Fungicide. Please refer to respective product labels for specific use directions, pertinent recommendations, restrictions and precautions.

Recommended Applications:

| Application Timing/Crop Stage/Product/Product Rates for single application | | Minimum Interval between Applications and Maximum Seasonal Application Rates | |
|--|-----------------------------|--|--|
| Vegetative stage | Heading/Anthesis | Minimum Interval between applications | Maximum seasonal tebuconazole rate (g a.i./ha) |
| HORNET 432 F 220-292 mL/ha | Prosaro 250 EC 800 mL/ha | 7 days | 226 |

Resistance Management Advisory for Cereal Crops:

Repeated application of standalone DMI Fungicides should not be used on the same crop in one season against risky pathogens such as cereal powdery mildew in areas of high disease pressure for that particular pathogen. Mixture products, tank-mixtures or alternation with fungicides having a different mode of action have been shown to protect against the development of resistant forms of disease.

HORNET 432 F Foliar Fungicide

Tank Mix Combinations with Herbicides:

| HORNET 432 F Foliar Fungicide can be tank-mixed with the herbicides noted below, for control of early season leaf diseases and weeds on the respective labels. Weeds and crop must be at the correct growth stage as specified in both the HORNET 432 F Foliar Fungicide and the tank mix partner label. Always read and follow the label for HORNET 432 F Foliar Fungicide and that of the tank mix partner(s) regarding the directions for use, restrictions and precautions. Do not tank mix HORNET 432 F Foliar Fungicide with pesticides, fertilizers or any other chemical additives unless recommended on this label. | | | |
|--|---|--|--|
| CROP | DISEASES and WEEDS | TANKMIXES | REMARKS |
| Wheat | Leaf Diseases on the HORNET 432 F Foliar Fungicide label and weeds on the respective herbicide labels | HORNET 432 F Foliar Fungicide at 220 mL/ha + Buctril M at 1L/ha | Apply HORNET 432 F Foliar Fungicide and herbicide at labeled rates for the respective leaf diseases and weeds in wheat. Apply once when plants are at the 4-leaf stage up to the early flag leaf stage. |
| | | HORNET 432 F Foliar Fungicide at 220 mL/ha + Puma ¹²⁰ Super at 385 or 770 mL/ha + Buctril M at 1L/ha | For use in the prairie provinces and Peace River region of British Columbia only. Apply HORNET 432 F Foliar Fungicide and herbicides at labeled rates for the respective leaf diseases and weeds in wheat. |
| | | HORNET 432 F Foliar Fungicide at 220 mL/ha + Puma ¹²⁰ Super at 385 or 770 mL/ha + Refine Extra at 20 g/ha | Apply once when plants are at the 4-leaf stage up to the 3 rd tiller stage. DO NOT apply HORNET 432 F Foliar Fungicide mixed with Puma ¹²⁰ Super 2-3 days prior to or following cold temperatures (3°C or lower) as crop injury may occur. |
| Barley | Leaf Diseases on the HORNET 432 F Foliar Fungicide label and weeds on the respective herbicide labels | HORNET 432 F Foliar Fungicide at 220 mL/ha + Puma ¹²⁰ Super at 385 or 770 mL/ha + Refine Extra at 20 g/ha | For use in the prairie provinces and Peace River region of British Columbia only. Apply HORNET 432 F Foliar Fungicide and herbicides at labeled rates for the respective leaf diseases and weeds in barley. Apply once when plants are at the 4-leaf stage to the 5-leaf, 2 nd tiller stage. DO NOT apply HORNET 432 F Foliar Fungicide mixed with Puma ¹²⁰ Super 2-3 days prior to or following cold temperatures (3°C or lower) as crop injury may occur. |

Tank Mix Combinations with Insecticides:

HORNET 432 F Foliar Fungicide can be tank-mixed with Lorsban 4E Insecticide for control of Orange wheat blossom midge (*Sitodiplosis mosellana*) in wheat at labelled rates. Read carefully and follow use directions and use precautions on both the HORNET 432 F Foliar Fungicide and Lorsban 4E Insecticide label. When tank mixing HORNET 432 F Foliar Fungicide and Lorsban 4E Insecticide, always add HORNET 432 F Foliar Fungicide to the tank first, then add Lorsban 4E Insecticide. Do not add any surfactants.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, please note that HORNET 432 F Foliar Fungicide contains a Group 3 fungicide. Any fungal population may contain individuals naturally resistant to HORNET 432 F Foliar Fungicide and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

Where possible, rotate the use of HORNET 432 F Foliar Fungicide or other Group 3 fungicides with different groups that control the same pathogens.

Do not apply more than the indicated maximum number of applications specified for each crop in the DIRECTIONS FOR USE. Avoid using consecutive applications of other fungicides in the same group in a season.

Use tank mixtures with fungicides from a different group that is effective on the target pathogen when such use is permitted.

Fungicide use should be based on an IPM program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.

Where possible, make use of predictive disease models to effectively time fungicide applications.

Monitor treated fungal populations for resistance development. Notify Nufarm Agriculture Inc. if reduced sensitivity of the pathogen to HORNET 432 F Foliar Fungicide is suspected.

If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and disease problems in your area.

For further information and to report suspected resistance, contact Nufarm Agriculture Inc. at 1-800-868-5444 or at www.nufarm.ca.

APPLICATION IN FIELDS ADJACENT TO AQUATIC AREAS

OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, AND ESTUARIES.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of

aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

This product should be used only in alternate years.

Do not apply by ground or air within 30 metres of aquatic areas listed above.

Do not cultivate within 3 metres of an aquatic area to allow growth of a vegetative filter strip.

The aquatic buffer zone of 30 meters **may not** be modified by the Buffer Zone Calculator on the Pest Management Regulatory Agency web site.

SPRAY DRIFT MANAGEMENT FOR AERIAL AND GROUND APPLICATIONS

For the protection of non-target habitats, overspray or drift to any body of water or other environmentally sensitive habitats must be avoided. Do not apply under conditions where drift to an unprotected person(s), occupied dwelling, or to food, forage, or other plantings can occur.

The interaction of many equipment- and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

1. **SPRAY BOOM:** For aerial applications, the **spray boom** should be mounted on the aircraft so as to minimize drift caused by wing tip vortices. The minimum practical boom length should be used and must not exceed 65% of the wing span or rotor diameter.
2. **DROPLET SIZE:** An important factor influencing drift is the droplet size. Small droplets (<150 to 200 microns) drift to a greater extent than large droplets. Within typical equipment specifications, applications should be made to deliver the largest **droplet spectrum** that provides sufficient control and coverage. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure.
3. **SPRAY HEIGHT:** For aerial applications, spray should be released at the lowest height consistent with efficacy and flight safety. Applications more than 3 metres above the crop canopy should be avoided.
4. **WIND:** Do not apply during periods of dead calm, when winds are gusty or when wind speed is greater than 16 km/hour at flying height at the site of application. Use extreme caution when any body of water or other environmentally sensitive habitat is on downwind side of aircraft.
5. **TEMPERATURE INVERSIONS:** Do not make aerial or ground applications during temperature inversions. Drift potential is high during temperature inversions. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however if fog is not present, inversions can also be identified by the movement of smoke from a ground source. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical mixing.
6. **HUMIDITY AND TEMPERATURE:** Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift. Avoid spraying during conditions of low humidity and/or high temperature.

AERIAL APPLICATION LABEL INSTRUCTIONS

Directions For use

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices, or equivalent electronic positioning systems (GPS). The use of a spotter plane is recommended. Use mechanical flaggers only.

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the *National Aerial Pesticide Application Manual*, developed by the Federal/Provincial/Territorial committee on Pest management and Pesticides.

Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear coveralls, chemical resistant gloves, and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call Nufarm Agriculture Inc. at 1-800-868-5444 or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

Volume: Apply the recommended rate in a minimum spray volume of 47 litres per hectare.

ROTATIONAL CROPS

Treated areas may be replanted immediately following harvest with any crop listed on this label. For crops not listed on this label, do not plant back within 120 days of last application.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist.

Wear coveralls over long pants, long-sleeved shirt, chemical resistant gloves and boots during mixing, loading, application, clean-up and repair activities. Coveralls are not required during application within closed cabs or cockpits.

Follow manufacturer's instructions for cleaning/maintaining the Personal Protective Clothing (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

User Safety Recommendations:

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

FIRST AID

In case of poisoning, call physician or Poison Control Centre immediately. Have patient lie down and keep quiet.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.

To Physician: No specific antidote. Treat symptomatically.

EMERGENCY TELEPHONE NUMBERS

For spills or transportation accidents, Chemtrec, 1-800-424-9300

For health or environmental emergencies, ProPharma Group, 1-877-325-1840

For product and use information, Nufarm Agriculture Inc., 1-800-868-5444

ENVIRONMENTAL HAZARDS

Tebuconazole is persistent and will carryover. It is recommended that any products containing tebuconazole not be used in areas treated with this product during the previous season.

HORNET 432 F Foliar Fungicide

Toxic to birds, small wild animals, aquatic organisms, and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE and APPLICATION IN FIELDS ADJACENT TO AQUATIC AREAS. As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests. Do not apply to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff is hazardous to aquatic organisms in neighboring areas. To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body. To reduce the run off refer to the recommendations under "APPLICATION IN FIELDS ADJACENT TO AQUATIC AREAS".

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

LEAK OR SPILL PROCEDURE: Handle and open container in a manner as to prevent spillage. If container is leaking invert to prevent leakage. If the container is leaking or material is spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away. For decontamination procedures or any other assistance that may be necessary, you may contact the 24 hour emergency response number 1-800-424-9300.

DISPOSAL:

Recyclable Container Disposal: Do not reuse this container for any purpose. This is a recyclable container and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

Returnable Container: Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable Container: For disposal, this empty container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not use this container for any other purpose.

Non-Returnable Container:

1. Triple or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.

HORNET 432 F Foliar Fungicide

Disposal of Unused, Unwanted Product: For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills.

TM HORNET is a trademark of Nufarm Agriculture Inc.
All other products mentioned are trademarks of their respective companies.