

GROUP	3	4	7	12	FUNGICIDES
-------	---	---	---	----	------------

VIBRANCE® FLEXI CANOLA

SEED TREATMENT FUNGICIDE

COMMERCIAL – AGRICULTURAL

SUSPENSION

Fungicide seed treatment for use on labelled crops to control seed and soil-borne diseases.

ACTIVE INGREDIENTS:

Difenoconazole	40 g/L
Metalaxyl-M and S-isomer	12.5 g/L
Sedaxane.....	8.4 g/L
Fludioxonil.....	4.2 g/L

Contains 1,2-benzisothiazolin-3-one at 0.004%, 5-chloro-2-methyl-4-isothiazolin-3-one at 0.00056% and 2-methyl-4-isothiazolin-3-one at 0.00019% as preservatives.

**READ THE LABEL AND ATTACHED PAMPHLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

Warning, contains the allergen soy.

REGISTRATION NUMBER: **32624**
PEST CONTROL PRODUCTS ACT

NET CONTENTS: **1 L – 1050 L**

Syngenta Canada Inc.
140 Research Lane, Research Park
Guelph, Ontario N1G 4Z3
Telephone: 1-877-964-3682

This product contains no colourant. All seed treated with this product must be conspicuously coloured at the time of treatment. Regulations pertaining to the “Seeds Act” must be strictly adhered to when using this product.

For use with commercial seed treaters (facilities and mobile treaters) with closed transfer systems only. Closed transfer includes closed mixing, loading, calibrating and closed treatment equipment. No open transfer of VIBRANCE® FLEXI CANOLA is permitted.

LABEL

1.0 NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label.

2.0 FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

3.0 TOXICOLOGICAL INFORMATION

There is no specific antidote. Treat symptomatically.

4.0 PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS.

Do not contaminate food, feed, or domestic water supplies.

Apply this product to the seed only in a way that it will not contact workers or other persons, either directly or through drift. Only workers wearing personal protective equipment may be in the area where treating, bagging, sewing, and/or stacking is occurring.

VIBRANCE® FLEXI CANOLA is for closed transfer commercial seed treatment (in facilities or with mobile treaters) only on canola. Closed transfer includes closed mixing, loading, calibrating and closed treatment equipment. No open transfer is permitted. Treat seed in a well-ventilated area. Keep treated seeds out of reach of children and animals.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please

contact Syngenta Canada Inc. at 1-87-SYNGENTA / 1-877-964-3682.

5.0 PERSONAL PROTECTIVE EQUIPMENT (PPE)

Commercial Seed Treatment:

Workers involved in clean-up or contacting contaminated seed-treating equipment must wear chemical-resistant coveralls over a long-sleeved shirt and long pants, chemical-resistant gloves, socks and chemical-resistant footwear. Workers involved in treating seeds or repair or maintenance of seed treating equipment must wear coveralls over a long-sleeved shirt and long pants, chemical-resistant gloves, socks and chemical-resistant footwear. Workers involved in bagging, sewing, handling, and stacking bags of treated seed, or when transferring seed to a storage bin must wear coveralls over long-sleeved shirt and long pants, chemical-resistant gloves, a NIOSH-approved N95 filtering facepiece respirator (dust mask) that is properly fit tested, and socks and shoes. Forklift operators must wear coveralls over long-sleeved shirt, long pants, chemical-resistant gloves, and socks plus shoes. For good hygiene practice, it is also recommended to wear a NIOSH-approved N95 filtering facepiece respirator (dust mask) that is properly fit tested during all job activities.

Planting Treated Seed:

Wear long-sleeved shirt, long pants, shoes plus socks and chemical-resistant gloves when handling and planting treated seed. For good hygiene practice, it is also recommended to wear a NIOSH-approved N95 filtering facepiece respirator (dust mask) that is properly fit tested during all job activities. When using closed-cab planting equipment, chemical-resistant gloves are not required inside cab.

All Activities:

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Wash splashes from skin IMMEDIATELY with plenty of water. Remove PPE immediately after handling this product. Wash the outside of gloves with soap and water before removing. As soon as possible, wash thoroughly and change into clean clothing. After treating, wash hands and shower thoroughly with soap and water.

Heavily contaminated or drenched clothing and other absorbent materials contaminated with this product should be discarded. Do not reuse them.

Users should wash hands and exposed skin before eating, drinking, chewing gum, using tobacco, applying cosmetics or using the toilet.

While using product, do not eat, drink or use tobacco (including smoking).

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Wear freshly laundered clothes daily.

Do not wear contaminated shoes.

6.0 ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms. Do not contaminate food, feed, domestic or irrigation water supplies, lakes, streams and ponds.

Treated seed is TOXIC to birds and small wild mammals. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil or other surfaces.

7.0 USE RESTRICTIONS

1. Treated seed must not be used for food, feed or oil processing.
2. Store away from food, feed and out of the reach of children and animals.
3. Do not graze or feed livestock on treated areas for 45 days after planting.
4. Do not plant any crop other than barley, oats, rye, triticale, wheat, canola, rapeseed, or mustard within 60 days to fields in which seed treated with VIBRANCE FLEXI CANOLA were planted.
5. All bags containing treated seed for sale or use in Canada must be labeled or tagged as follows: **“This seed has been treated with difenoconazole, metalaxyl-M and S-isomer, fludioxonil and sedaxane fungicides. Wear long-sleeved shirt, long pants, shoes plus socks and chemical-resistant gloves when handling and planting treated seed. When using closed-cab planting equipment, chemical-resistant gloves are not required inside cab. For good hygiene practice, it is also recommended to wear a NIOSH-approved N95 filtering facepiece respirator (dust mask) that is properly fit tested during all job activities. Keep treated seed out of reach of children and animals. Do not use for food, feed or oil processing. Store this product away from food and feed. Treated seed is TOXIC to birds. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface.”**

8.0 STORAGE

Keep in original container, tightly closed, during storage. Store in a cool, dry, well-ventilated area away from other pesticides, feed, food and out of the reach of children and animals. Store this product away from from seed, food or feed.

9.0 DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

For Refillable Containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

For Returnable Containers

DO NOT reuse this container for any purpose. For disposal, the empty container may be returned to the point of purchase (distributor/dealer).

For Recyclable Containers:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

VIBRANCE® is a trademark of a Syngenta Group Company.

GROUP	3	4	7	12	FUNGICIDES
-------	---	---	---	----	------------

VIBRANCE® FLEXI CANOLA

SEED TREATMENT FUNGICIDE

COMMERCIAL – AGRICULTURAL

SUSPENSION

Fungicide seed treatment for use on labelled crops to control seed and soil-borne diseases.

ACTIVE INGREDIENTS:

Difenoconazole	40 g/L
Metalaxyl-M and S-isomer	12.5 g/L
Sedaxane.....	8.4 g/L
Fludioxonil.....	4.2 g/L

Contains 1,2-benzisothiazolin-3-one at 0.004%, 5-chloro-2-methyl-4-isothiazolin-3-one at 0.00056% and 2-methyl-4-isothiazolin-3-one at 0.00019% as preservatives.

**READ THE LABEL AND ATTACHED PAMPHLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

Warning, contains the allergen soy.

REGISTRATION NUMBER: **32624**
PEST CONTROL PRODUCTS ACT

Syngenta Canada Inc.
140 Research Lane, Research Park
Guelph, Ontario N1G 4Z3
Telephone: 1-877-964-3682

This product contains no colourant. All seed treated with this product must be conspicuously coloured at the time of treatment. Regulations pertaining to the “Seeds Act” must be strictly adhered to when using this product.

For use with commercial seed treaters (facilities and mobile treaters) with closed transfer systems only. Closed transfer includes closed mixing, loading, calibrating and closed treatment equipment. No open transfer of VIBRANCE® FLEXI CANOLA is permitted.

PAMPHLET

TABLE OF CONTENTS

	<u>Section Number</u>
NOTICE TO USER	1.0
FIRST AID	2.0
TOXICOLOGICAL INFORMATION	3.0
PRECAUTIONS.....	4.0
PERSONAL PROTECTIVE EQUIPMENT (PPE).....	5.0
ENVIRONMENTAL PRECAUTIONS.....	6.0
USE RESTRICTIONS.....	7.0
STORAGE	8.0
DISPOSAL	9.0
PRODUCT INFORMATION	10.0
DIRECTIONS FOR USE	11.0
General Information	11.1
CROP USE DIRECTIONS	12.0
RESISTANCE MANAGEMENT RECOMMENDATIONS	13.0

1.0 NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label.

2.0 FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

3.0 TOXICOLOGICAL INFORMATION

There is no specific antidote. Treat symptomatically.

4.0 PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS.

Do not contaminate food, feed, or domestic water supplies.

Apply this product to the seed only in a way that it will not contact workers or other persons, either directly or through drift. Only workers wearing personal protective equipment may be in the area where treating, bagging, sewing and/or stacking is occurring.

VIBRANCE® FLEXI CANOLA is for closed transfer commercial seed treatment (in facilities or with mobile treaters) only on canola. Closed transfer includes closed mixing, loading, calibrating and closed treatment equipment. No open transfer is permitted. Treat seed in a well-ventilated area. Keep treated seeds out of reach of children and animals.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please

contact Syngenta Canada Inc. at 1-87-SYNGENTA / 1-877-964-3682.

5.0 PERSONAL PROTECTIVE EQUIPMENT (PPE)

Commercial Seed Treatment:

Workers involved in clean-up or contacting contaminated seed-treating equipment must wear chemical-resistant coveralls over a long-sleeved shirt and long pants, chemical-resistant gloves, socks and chemical-resistant footwear. Workers involved in treating seeds or repair or maintenance of seed treating equipment must wear coveralls over a long-sleeved shirt and long pants, chemical-resistant gloves, socks and chemical-resistant footwear. Workers involved in bagging, sewing, handling and stacking bags of treated seed, or when transferring seed to a storage bin must wear coveralls over long-sleeved shirt and long pants, chemical-resistant gloves, a NIOSH-approved N95 filtering facepiece respirator (dust mask) that is properly fit tested, and socks and shoes. Forklift operators must wear coveralls over long-sleeved shirt, long pants, chemical-resistant gloves, and socks plus shoes. For good hygiene practice, it is also recommended to wear a NIOSH-approved N95) filtering facepiece respirator (dust mask) that is properly fit tested during all job activities.

Planting Treated Seed:

Wear long-sleeved shirt, long pants, shoes plus socks and chemical-resistant gloves when handling and planting treated seed. For good hygiene practice, it is also recommended to wear a NIOSH-approved N95 filtering facepiece respirator (dust mask) that is properly fit tested during all job activities. When using closed-cab planting equipment, chemical-resistant gloves are not required inside cab.

All Activities:

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Wash splashes from skin IMMEDIATELY with plenty of water. Remove PPE immediately after handling this product. Wash the outside of gloves with soap and water before removing. As soon as possible, wash thoroughly and change into clean clothing. After treating, wash hands and shower thoroughly with soap and water.

Heavily contaminated or drenched clothing and other absorbent materials contaminated with this product should be discarded. Do not reuse them.

Users should wash hands and exposed skin before eating, drinking, chewing gum, using tobacco, applying cosmetics or using the toilet.

While using product, do not eat, drink or use tobacco (including smoking).

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Wear freshly laundered clothes daily.

Do not wear contaminated shoes.

6.0 ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms. Do not contaminate food, feed, domestic or irrigation water supplies, lakes, streams and ponds.

Treated seed is TOXIC to birds and small wild mammals. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil or other surfaces.

7.0 USE RESTRICTIONS

1. Treated seed must not be used for food, feed or oil processing.
2. Store away from food, feed and out of the reach of children and animals.
3. Do not graze or feed livestock on treated areas for 45 days after planting.
4. Do not plant any crop other than barley, oats, rye, triticale, wheat, canola, rapeseed, or mustard within 60 days to fields in which seed treated with VIBRANCE FLEXI CANOLA were planted.
5. All bags containing treated seed for sale or use in Canada must be labeled or tagged as follows: **“This seed has been treated with difenoconazole, metalaxyl-M and S-isomer, fludioxonil and sedaxane fungicides. Wear long-sleeved shirt, long pants, shoes plus socks and chemical-resistant gloves when handling and planting treated seed. When using closed-cab planting equipment, chemical-resistant gloves are not required inside cab. For good hygiene practice, it is also recommended to wear a NIOSH-approved N95 filtering facepiece respirator (dust mask) that is properly fit tested during all job activities. Keep treated seed out of reach of children and animals. Do not use for food, feed or oil processing. Store this product away from food and feed. Treated seed is TOXIC to birds. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface.”**

8.0 STORAGE

Keep in original container, tightly closed, during storage. Store in a cool, dry, well-ventilated area away from other pesticides and out of the reach of children and animals. Store this product away from seed, food or feed.

9.0 DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

For Refillable Containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

For Returnable Containers

DO NOT reuse this container for any purpose. For disposal, the empty container may be returned to the point of purchase (distributor/dealer).

For Recyclable Containers:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the treatment tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

10.0 PRODUCT INFORMATION

VIBRANCE FLEXI CANOLA is a seed treatment fungicide containing difenoconazole, metalaxyl-M and S-isomer, fludioxonil and sedaxane. This product controls a broad spectrum of seed and soil-borne diseases in canola, rapeseed and mustard.

Note: Treatment of highly mechanically scarred or damaged seed, or seed known to be of low vigour and poor quality, except for the purpose of curative control of existing disease pests, may result in reduced germination, reduced emergence and/or reduction of seed and seedling vigour. Treat a small quantity of seed using equipment similar to that planned for treating the total seed lot. Conduct germination tests on a small portion of the seed before committing the total seed lot to a selected seed treatment. Due to seed quality and seed storage conditions beyond the control of Syngenta Canada Inc., no claims are made to guarantee the germination of carry-over treated seed.

11.0 DIRECTIONS FOR USE

11.1 General Information

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

VIBRANCE FLEXI CANOLA is a seed treatment formulation for use in commercial seed treatment (in facilities or with mobile treaters) using closed transfer seed treatment equipment that accurately meters, mixes and applies a flowable seed treatment. The equipment must provide uniform coverage of VIBRANCE FLEXI CANOLA on the seed. Uneven seed coverage may not give the desired level of disease control.

Thoroughly mix the recommended amount of VIBRANCE FLEXI CANOLA with the required amount of water for the slurry treater and dilution rate to be used. Slurry volumes will vary depending on seed size. Follow the manufacturer's application instructions for the seed treatment equipment being used. Maintain constant agitation of the slurry during the treatment. Allow the seed to dry before bagging, storing or seeding.

Depending on planting equipment, seed treated with VIBRANCE FLEXI CANOLA, or a combination of VIBRANCE FLEXI CANOLA and other seed treatment products may not flow through planting equipment at the same rate as untreated seed. Recalibrate the equipment before planting treated seed.

This product contains no colourant. An appropriate colourant must be added when this product is applied. Users are responsible for ensuring that the treated seed, when dried and ready for bagging, has an unnatural colour. Regulations pertaining to the "Seeds Act" must be strictly adhered to when using this product.

12.0 CROP USE DIRECTIONS

Apply 600 mL of VIBRANCE FLEXI CANOLA per 100 kg of seed to the crops listed below. At this rate, VIBRANCE FLEXI CANOLA delivers 24 g of difenoconazole, 2.5 g of fludioxonil, 7.5 g of metalaxyl-M and S-isomer, and 5 g of sedaxane per 100 kg of seed.

12.1 Canola

CROPS	Canola Rapeseed Mustard (Condiment and Oilseed)
DISEASE CONTROLLED	Seed-borne Blackleg (<i>Leptosphaeria maculans</i>) Seed-borne Alternaria (<i>Alternaria spp.</i>) Seedling disease complex (damping-off, seedling blight, seed rot, and root rot) caused by <i>Pythium spp.</i> , <i>Fusarium spp.</i> and <i>Rhizoctonia spp.</i>
RATE	600 mL / 100 kg seed (39 g ai total/100 kg seed)
APPLICATION METHOD	Seed Treatment
NOTES	For control of early season seed and soil-borne diseases based on 200,000 seeds per kg (5.0 g/1000 seeds), VIBRANCE FLEXI CANOLA delivers 0.39 g ai/kg or 1.8 µg active ingredient per seed (1.2 of µg difenoconazole, 0.3 µg of metalaxyl-M, 0.1 µg of fludioxonil, 0.2 µg of sedaxane)
RESTRICTIONS	This product contains no colourant. An appropriate colourant must be added when this product is applied to the seed.

12.2 Tank Mixes

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Syngenta Canada Inc. at 1-87-SYNGENTA / 1-877-964-3682 for information before applying any tank mix that is not specifically recommended on this label.

Syngenta Canada Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) of all potential tank mixes under all environmental conditions or for all crop varieties. Tank mixes that are not specifically listed on this label should be tested on a small area first, under local conditions and using standard practices, to confirm the tank mix is suitable for widespread application.

To determine the physical compatibility of this product with other products, use a jar test.

Always read and follow label directions including WALES mixing order.

13.0 RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, VIBRANCE FLEXI CANOLA contains Group 3, 4, 7 and 12 fungicides. Any fungal population may contain individuals naturally resistant to VIBRANCE FLEXI CANOLA and other Group 3, 4, 7 and 12 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

Where possible, rotate the use of VIBRANCE FLEXI CANOLA or other Group 3, 4, 7 and 12 fungicides with different groups that control the same pathogens.

Use tank mixtures with fungicides from a different group that is effective on the target pathogen when such use is permitted.

Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.

Monitor treated fungal populations for resistance development. Notify Syngenta Canada Inc. if reduced sensitivity of the pathogen to VIBRANCE FLEXI CANOLA is suspected.

If disease continues to progress after treatment with this product, DO NOT increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pests problems in your area.

For further information and to report suspected resistance, contact Syngenta Canada Inc. at 1-877-SYNGENTA (1-877-964-3682) or at www.syngenta.ca.