

(Container)

# Retain<sup>®</sup> B 333 Herbicide

|       |   |           |
|-------|---|-----------|
| GROUP | 4 | HERBICIDE |
|-------|---|-----------|

Retain B 333 Herbicide is a selective herbicide for postemergent control of annual broadleaved weeds including cleavers and certain ALS-resistant broadleaved weeds such as kochia in spring wheat, durum wheat, winter wheat and spring barley. Retain B 333 Herbicide will also suppress hemp-nettle, common chickweed and wild buckwheat.

COMMERCIAL (AGRICULTURAL)

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING  
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: fluroxypyr, present as 1-methylheptyl ester 333 g/L  
Emulsifiable concentrate

REGISTRATION NO. 32845 PEST CONTROL PRODUCTS ACT

**WARNING**  
**EYE AND SKIN IRRITANT**  
**POTENTIAL SKIN SENSITIZER**

NET CONTENTS: 1 L - bulk

**Loveland Products Canada Inc.**  
789 Donnybrook Drive  
Dorchester, ON N0L 1G5

**Corteva Agriscience Canada Company**  
Suite 240, 115 Quarry Park Rd. SE  
Calgary, AB T2C 5G9

For information: 1-800-328-4678  
In case of emergency call: 1-800-561-8273

®Retain is a registered trademark

## **PRECAUTIONS**

### **KEEP OUT OF REACH OF CHILDREN**

Causes eye irritation. DO NOT get in eyes. May irritate the skin. Avoid contact with skin. Potential skin sensitizer.

**For agricultural/crop scenarios:** DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12-hours. **For non-crop areas:** DO NOT enter or allow worker entry into treated areas until sprays have dried.

DO NOT use in residential areas. Residential areas are defined as any use site where bystanders including children could be exposed during or after application. This includes homes, schools, parks, playgrounds, playing fields, public buildings, or any other area where the general public including children could be exposed.

Apply only when potential for drift to areas of human habitation and human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversion, application equipment and sprayer settings.

### **PROTECTIVE CLOTHING AND EQUIPMENT**

Wear a hat, coveralls over a long-sleeved shirt and long pants, socks, rubber boots or work boots and chemical-resistant gloves during spraying or application. In addition, wear goggles or face shield during mixing, loading, repair and clean-up or when handling the concentrate. Applicators using a closed cab are not required to wear chemical-resistant gloves.

### **OPERATOR USE PRECAUTIONS**

Wear freshly laundered clothing and clean protective equipment daily. At completion of spraying or end of the day take a shower immediately. Wash thoroughly with soap and water before eating, drinking, smoking or using the toilet. Remove contaminated clothing at the end of the work session and store and wash separately from household laundry using detergents and hot water before reuse. If herbicide penetrates clothing, remove immediately; then wash thoroughly and put on clean clothing. Throw away clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Avoid contact with eyes, skin and clothing. Avoid breathing spray mist.

### **PHYSICAL OR CHEMICAL HAZARDS**

**COMBUSTIBLE.** Do not use or store near heat or open flame.

### **FIRST AID**

**Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.**

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**TOXICOLOGICAL INFORMATION**

THIS PRODUCT CONTAINS A PETROLEUM DISTILLATE. Vomiting may cause aspiration pneumonia. The decision of whether to induce vomiting or not should be made by the attending physician. If lavage is performed, suggest endotracheal and/or oesophageal control. Danger from lung aspiration of petroleum based solvents must be weighed against toxicity when considering emptying the stomach. No specific antidote. Employ supportive care. Treatment should be based on the judgment of the physician in response to reactions of the patient.

**AGRICULTURAL CHEMICAL**

Do not ship or store with food, feeds, drugs or clothing.

**ENVIRONMENTAL PRECAUTIONS**

TOXIC to non-target terrestrial plants and aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE. To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

This product contains an active ingredient and aromatic petroleum distillates which are toxic to aquatic organisms.

**STORAGE**

Store in original containers in a secure, dry heated storage. If product is frozen, bring to room temperature and agitate before use. Do not allow contamination of seeds, plants, fertilizers or other pesticides. To prevent contamination, store this product away from food or feed. If containers are damaged or spill occurs, use the product immediately or contain the spill with absorbent materials and dispose of waste.

**DISPOSAL****Recyclable Containers:**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

**Returnable Containers:**

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

**Refillable Containers:**

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.



## **PRECAUTIONS**

### **KEEP OUT OF REACH OF CHILDREN**

Causes eye irritation. DO NOT get in eyes. May irritate the skin. Avoid contact with skin. Potential skin sensitizer.

**For agricultural/crop scenarios:** DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12-hours. **For non-crop areas:** DO NOT enter or allow worker entry into treated areas until sprays have dried.

DO NOT use in residential areas. Residential areas are defined as any use site where bystanders including children could be exposed during or after application. This includes homes, schools, parks, playgrounds, playing fields, public buildings, or any other area where the general public including children could be exposed.

Apply only when potential for drift to areas of human habitation and human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversion, application equipment and sprayer settings.

### **PROTECTIVE CLOTHING AND EQUIPMENT**

Wear a hat, coveralls over a long-sleeved shirt and long pants, socks, rubber boots or work boots and chemical-resistant gloves during spraying or application. In addition, wear goggles or face shield during mixing, loading, repair and clean-up or when handling the concentrate. Applicators using a closed cab are not required to wear chemical-resistant gloves.

### **OPERATOR USE PRECAUTIONS**

Wear freshly laundered clothing and clean protective equipment daily. At completion of spraying or end of the day take a shower immediately. Wash thoroughly with soap and water before eating, drinking, smoking or using the toilet. Remove contaminated clothing at the end of the work session and store and wash separately from household laundry using detergents and hot water before reuse. If herbicide penetrates clothing, remove immediately; then wash thoroughly and put on clean clothing. Throw away clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Avoid contact with eyes, skin and clothing. Avoid breathing spray mist.

### **PHYSICAL OR CHEMICAL HAZARDS**

**COMBUSTIBLE.** Do not use or store near heat or open flame.

### **FIRST AID**

**Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.**

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**TOXICOLOGICAL INFORMATION**

THIS PRODUCT CONTAINS A PETROLEUM DISTILLATE. Vomiting may cause aspiration pneumonia. The decision of whether to induce vomiting or not should be made by the attending physician. If lavage is performed, suggest endotracheal and/or oesophageal control. Danger from lung aspiration of petroleum based solvents must be weighed against toxicity when considering emptying the stomach. No specific antidote. Employ supportive care. Treatment should be based on the judgment of the physician in response to reactions of the patient.

**AGRICULTURAL CHEMICAL**

Do not ship or store with food, feeds, drugs or clothing.

**ENVIRONMENTAL PRECAUTIONS**

TOXIC to non-target terrestrial plants and aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE. To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

This product contains an active ingredient and aromatic petroleum distillates which are toxic to aquatic organisms.

**STORAGE**

Store in original containers in a secure, dry heated storage. If product is frozen, bring to room temperature and agitate before use. Do not allow contamination of seeds, plants, fertilizers or other pesticides. To prevent contamination, store this product away from food or feed. If containers are damaged or spill occurs, use the product immediately or contain the spill with absorbent materials and dispose of waste.

**DISPOSAL****Recyclable Containers:**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

**Returnable Containers:**

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

**Refillable Containers:**

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

## GENERAL INFORMATION

Retain B 333 Herbicide is a selective postemergent herbicide for the control of annual broadleaved weeds in spring wheat, durum wheat, winter wheat and spring barley, not underseeded with legumes. Retain B 333 Herbicide offers a novel mode of action for controlling hard-to-kill annual broadleaved weeds such as kochia (including ALS resistant biotypes) and cleavers. Retain B 333 Herbicide will also provide suppression of hempnettle, wild buckwheat and chickweed.

## MODE OF ACTION

Retain B 333 Herbicide is a systemic auxin-type herbicide and moves within the plant for control of exposed and underground plant tissues. The product controls weeds by disrupting normal plant growth patterns. Symptoms of weeds include epinasty (twisting of the stems) and swollen nodes.

## GENERAL USE PRECAUTIONS

### Sensitive Plants

Do not apply Retain B 333 Herbicide directly to, or otherwise permit it to come in direct contact with susceptible crops or desirable plants including alfalfa, edible beans, canola, flowers and ornamentals, lentils, lettuce, peas, potatoes, radishes, soybeans, sugar beets, sunflowers, tomatoes or tobacco.

### Non-Target Sites

Do not apply where proximity of susceptible crops or other desirable plants is likely to result in exposure to spray or spray drift. See ENVIRONMENTAL PRECAUTIONS section of the label.

### Crop Rotation

Fields previously treated with Retain B 333 Herbicide can be seeded the following year to alfalfa, barley, canola, corn, dry beans, fababeans, flax, forage grasses, lentils, mustard, oats, peas, potatoes, rye, soybeans, sugar beets, sunflowers or wheat or fields can be summerfallowed.

### Tank Mixtures

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact LOVELAND PRODUCTS CANADA INC. at 1-800-328-4678 or [www.lovelandproducts.ca](http://www.lovelandproducts.ca) for information before applying any tank mix that is not specifically recommended on this label.

### Spray Equipment Precaution

Do not apply through any type of irrigation system.

### To Reduce Spray Drift

- Use nozzles delivering higher volumes and coarser droplets.
- Use low pressures (200 to 275 kPa).
- Spray when the wind velocity is 15 km/hr or less.
- Spot treatments should only be applied with a calibrated boom to prevent over-application.

### Equipment Clean-Up

Equipment used to apply Retain B 333 Herbicide should not be used to apply other pesticides to sensitive crops without thorough cleaning. Contact your Retain B 333 Herbicide dealer for detailed equipment cleaning procedures.

## DIRECTIONS FOR USE

READ THE ENTIRE LABEL BEFORE USE. FAILURE TO FOLLOW LABEL INSTRUCTIONS MAY RESULT IN ERRATIC WEED CONTROL OR CROP DAMAGE. DO NOT APPLY TO CROPS UNDERSEEDING WITH LEGUMES.

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

**DO NOT** contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

#### Field sprayer application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

#### Aerial application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length MUST NOT exceed 65% of the wing or rotorspan.

#### Crops Registered

spring wheat  
durum wheat  
winter wheat  
spring barley

#### TANK-MIX COMBINATION – RETAIN B 333 HERBICIDE + 2,4-D ESTER HERBICIDE (Do not apply to oats)

For control of a wide spectrum of broadleaved weeds (see following table) apply Retain B 333 Herbicide tank mixed with 2,4-D ester herbicide. Apply postemergence when weeds are in the seedling stage (2-4 leaf) and when spring wheat, durum wheat, winter wheat and spring barley are in the 4-leaf to flag-leaf stage. Apply to winter wheat in the spring from the 3 tiller stage to just before the flag leaf stage. Application before the 4-leaf stage of wheat and barley may cause severe twisting of leaves and stem, and head deformities which may reduce yield up to 10%. Do not apply later than the flag-leaf stage.

#### Rates of Applications

**I. Low Rate:** To control the following weeds apply a uniform spray containing 0.23 L/ha Retain B 333 Herbicide and 420 g a.e./ha 2,4-D ester herbicide (equivalent to 0.64 L/ha of 2,4-D Ester LV700 Herbicide).

#### Weeds Controlled

|                       |                                      |                           |
|-----------------------|--------------------------------------|---------------------------|
| bluebur               | goat's-beard                         | shepherd's-purse          |
| burdock               | hoary cress*                         | stinkweed                 |
| canola (volunteer)    | kochia***                            | sunflower (annual)        |
| cleavers (1-6 whorls) | lamb's-quarters                      | vetch                     |
| clovers (sweet)       | mustards (except green, dog & tansy) | wild radish               |
| cocklebur             | plantain                             | wild mustard              |
| flixweed              | prickly lettuce                      | wild buckwheat (1-6 leaf) |
| field horsetail*      | ragweed                              |                           |

**II. High Rate:** To control the above weeds plus the additional weeds listed below, apply a uniform spray containing 0.31 L/ha of Retain B 333 Herbicide and 560 g a.e./ha of 2,4-D ester herbicide (equivalent to 0.85 L/ha of 2,4-D Ester LV700 Herbicide).

**Additional Weeds Controlled:**

|                       |                                |                           |
|-----------------------|--------------------------------|---------------------------|
| blue lettuce*         | hairy galinsoga                | Russian thistle           |
| cleavers (1-8 whorls) | hedge bindweed                 | smartweed                 |
| dandelion**           | lady's-thumb                   | stork's-bill (1-8 leaf)   |
| docks                 | leafy spurge*                  | tansy mustard             |
| dog mustard           | oak-leaved goosefoot           | tartary buckwheat         |
| field bindweed*       | redroot pigweed                | volunteer flax (1-12 cm)  |
| field peppergrass     | round-leaved mallow (1-6 Leaf) | wild buckwheat (1-8 leaf) |
| gumweed               |                                |                           |

**Weeds Suppressed:**

|                                  |                              |                           |
|----------------------------------|------------------------------|---------------------------|
| common chickweed*** (up to 8 cm) | hemp-nettle (2-6 leaf stage) | sow-thistle (perennial) * |
| Canada thistle*                  | sow-thistle (annual)         |                           |

\* Top growth control only

\*\* Spring rosettes

\*\*\* Including biotypes resistant to Group 2 herbicides that inhibit the ALS enzyme

**NOTE:** The activity of Retain B 333 Herbicide combined with 2,4-D ester herbicide is influenced by weather conditions. Optimum activity requires active crop and weed growth. The temperature range for optimum activity is 12°C to 24°C. Reduced activity will occur when temperatures are below 8°C or above 27°C. Frost before application (3 days) or shortly after (3 days) may reduce weed control and crop tolerance. Weed control may be reduced during stress conditions, e.g. drought, heat or cold stress, or if weeds have initiated flowering, or if heavy infestations exist.

**Mixing Instructions**

Add the correct amount of Retain B 333 Herbicide to spray tank 1/2 filled with water and agitate. Add the correct amount of 2,4-D ester herbicide (e.g. 2,4-D Ester LV700 Herbicide) and continue to agitate.

**Pre-Harvest/Grazing Intervals (Retain B 333 Herbicide + 2,4-D ester herbicide)**

- Do not permit lactating dairy animals to graze fields within 7 days after application.
- Do not harvest forage or cut hay within 30 days after application.
- Withdraw meat animals from treated fields at least 3 days before slaughter.
- Do not harvest the treated mature crop within 60 days after application.

**TANK-MIX COMBINATIONS – Retain B 333 HERBICIDE + 2,4-D ESTER HERBICIDE + RETAIN A HERBICIDE or RETAIN A SG HERBICIDE (Do not apply to oats)****Spring Wheat (excluding durum) and spring barley**

For control of a wide spectrum of broadleaf weeds, including cleavers, Retain B 333 Herbicide may be tank mixed with 2,4-D ester herbicide and Retain A Herbicide **or** Retain A SG Herbicide. Apply Retain B 333 Herbicide at 0.16 L/ha tank mixed with Retain A Herbicide **or** Retain A SG Herbicide at 30 g/ha, 2,4-D ester herbicide at 280 g a.e./ha (equal to 0.5 L/ha of 2,4-D Ester LV 600 or 0.42 L/ha of 2,4-D Ester LV 700) and a non-ionic surfactant (Agral 90, Ag-Surf or Citowett Plus) at 2 L per 1000 L of spray solution. A spray volume of 100 L/ha should be utilized. Apply postemergence when weeds are in the seedling up to 4-leaf/whorls stage, and when spring wheat (excluding durum) and spring barley are in the 4-leaf to flag-leaf stage. **Do not apply to crops underseeded to legumes.**

**Weeds Controlled or Suppressed****Weeds Controlled:**

|  |                            |                                 |
|--|----------------------------|---------------------------------|
| annual smartweed (green smartweed, lady's thumb) | groundsel (common)         | shepherd's purse                |
| ball mustard                                     | hemp-nettle                | stinkweed                       |
| chickweed (1-6 leaf) <sup>1</sup>                | kochia <sup>3</sup>        | tartary buckwheat               |
| cleavers <sup>2</sup>                            | lamb'-quarters             | volunteer rapeseed <sup>4</sup> |
| corn spurry                                      | narrow-leaved hawk's beard | volunteer sunflower             |
| cow cockle                                       | redroot pigweed            | wild buckwheat <sup>5</sup>     |
| flixweed   | Russian thistle            | wild mustard                    |

<sup>1</sup> Spray before crop canopy prevents thorough coverage of weeds. Chickweed emerging after application will not be controlled.

<sup>2</sup> Excluding Group 2 resistant cleavers

<sup>3</sup> Excluding Group 2 resistant kochia

<sup>4</sup> The tank-mix will not control imazethapyr tolerant canola varieties (e.g. Clearfield varieties or other varieties with the Pursuit Smart trait)

<sup>5</sup> Apply to actively growing wild buckwheat from the cotyledon to 3-leaf stage. Control may be reduced under environmental stress. Large plants may regrow after treatment

**Weeds Suppressed:**

|   |                                  |
|---|----------------------------------|
| Canada thistle <sup>6</sup>                 | sow thistle <sup>6</sup>         |
| round leaved mallow (2-6 leaf) <sup>7</sup> | stork's bill (2-6 leaf)          |
| scentless chamomile                         | toadflax (to 15 cm) <sup>8</sup> |

<sup>6</sup> Apply when most of the thistles have emerged and are actively growing. Apply before bud stage, and before crop canopy prevents thorough coverage of thistles, when thistles are no larger than 15 cm in height for best top growth control. A single application will effectively inhibit the ability of emerged thistles to compete with the crop. Later emerging thistles will not be controlled.

<sup>7</sup> Apply to actively growing round leaved mallow. Will keep mallow stunted, but may not be able to reduce overall populations.

<sup>8</sup> A control program for this weed includes chemical application and frequent tillage.

**Mixing Instructions**

Fill the tank to 1/3 full of clean water. With agitator running, add required amount of Retain A Herbicide **or** Retain A SG Herbicide and continue to agitate for a minimum of 5 minutes to ensure Retain A Herbicide **or** Retain A SG Herbicide is completely dissolved. After it is fully dissolved add the Retain 333 B Herbicide with continuous agitation. Add the 2,4-D ester herbicide and the rest of the water. Then add adjuvant. Use a spray volume of 100 L/ha. Consult the 2,4-D label for additional application instructions and use precautions.

**Preharvest/Grazing Intervals (Retain 333 B Herbicide + Refine Extra + 2,4-D ester)**

- Allow 7 days between application and grazing of lactating dairy animals.
- Do not harvest forage or cut hay within 30 days after application.
- Remove meat animals from treated fields at least 3 days prior to slaughter.
- Do not harvest the treated mature crop within 60 days after application.

**TANK-MIX COMBINATIONS – RETAIN B 333 HERBICIDE + 2,4-D ESTER HERBICIDE + OTHER HERBICIDES (Do not apply to oats)**

Tank mixtures of Retain B 333 Herbicide + 2,4-D ester herbicide with other herbicides will provide control of additional weeds. For products which may be tank mixed with Retain B 333 Herbicide + 2,4-D ester herbicide and the additional weeds controlled see following table. Follow all precautions, minimal interval to harvest and directions for use on the Retain B 333 Herbicide and tank-mix partner labels.

**Tank-Mix Combinations with Retain B 333 Herbicide (0.31 L/ha) + 2,4-D ester herbicide (560 g a.e./ha) or Retain B 333 Herbicide (0.23 L/ha) + 2,4-D ester herbicide (420 g a.e./ha)**

| <b>Herbicide Tank-Mix Partner</b>       | <b>Crops Registered</b>                                | <b>Rate of Tank-Mix Partner</b> | <b>Adjuvant Rate</b>  | <b>Additional Weeds</b>   |
|---|--|---------------------------------|---|---|
| Liquid Achieve SC or Achieve Liquid*♦♦♦ | spring wheat, durum wheat, winter wheat, spring barley | 0.5 L/ha                        | Turbocharge D or Turbocharge 0.5 L/100 L of spray volume (0.5% v/v) | green foxtail, wild oats  |
| Simplicity™♦♦♦                          | spring wheat, durum wheat, winter wheat                | 0.5 L/ha                        | None required   | barnyard grass (1-5 leaf), common chickweed (up to 10 cm), hemp nettle (1-8 leaf), Japanese brome (1-6 leaf), wild oats (up to 4 leaf, 2 tiller), wild buckwheat (1-4 leaf) |

\* This tank-mix may cause temporary crop injury. However, yield will not normally be affected. This tank-mix may reduce control of wild oats compared to Liquid Achieve SC or Achieve Liquid Herbicide alone.

♦♦♦ Tank mixing with Simplicity only applies to the low rate of Retain B 333 Herbicide + 2,4-D ester

### **SPRAY BUFFER ZONES**

A spray buffer zone is NOT required for:

- uses with hand-held application equipment permitted on this label.

For application to rights-of-way, spray buffer zones for protection of sensitive terrestrial habitats are not required; however, the best available application strategies that minimize off-site drift, including meteorological conditions (for example, wind direction, low wind speed) and spray equipment (for example, coarse droplet sizes, minimizing height above canopy), should be used. However, applicators must observe the specified spray buffer zones for protection of sensitive aquatic habitats.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands) and sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

| Method of application | Crop  |             | Spray Buffer Zones (metres) Required for the Protection of: |                  |                                     |                  |                     |
|-----------------------|---|-------------|---|------------------|-------------------------------------|------------------|---------------------|
|                       |   |             | Freshwater Habitat of Depths:                               |                  | Estuarine/marine Habitat of Depths: |                  | Terrestrial habitat |
|                       |   |             | Less than 1 m   | Greater than 1 m | Less than 1 m                       | Greater than 1 m |                     |
| Field sprayer         | Spring wheat, durum wheat, winter wheat and spring barley |             | 1   | 0                | 1                                   | 1                | 3                   |
| Aerial                | Spring wheat, durum wheat, winter wheat and spring barley | Fixed Wing  | 5   | 0                | 1                                   | 1                | 95                  |
|                       |   | Rotary Wing | 3   | 0                | 1                                   | 1                | 80                  |

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

### RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, Retain B 333 Herbicide is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to Retain B 333 Herbicide and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of Retain B 333 Herbicide or other Group 4 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.

- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Loveland Products Canada Inc. at 1-800-328-4678.

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

All products listed are registered trademarks of their respective companies.