

2024-6576
2025-05-01

Container label

GROUP	3	7	11	FUNGICIDE
-------	---	---	----	-----------

Belyan®
FUNGICIDE

For use in Canola, Rapeseed, Mustard (including oilseed/condiment mustard), Flax, Crop Subgroup 6C (Dried Shelled Peas and Beans, except Soybean), Potato, Soybean and Wheat (All Types)

Suspension Concentrate

COMMERCIAL (AGRICULTURAL)

ACTIVE INGREDIENTS: Mefentrifluconazole 133.3 g/L
Pyraclostrobin 177.8 g/L
Fluxapyroxad88.9 g/L

Contains 5-chloro-2-methyl-4-isothiazolin-3-one at 0.00111 % , 2-methyl-4-isothiazolin-3-one at 0.00037%, 1,2-benzisothiazolin-3-one at 0.0400 % and 2-bromo-2-nitropropane-1,3-diol at 0.0240% as preservatives

OR

Contains 1,2-benzisothiazolin-3-one and 2-methyl-4-isothiazolin-3-one at 0.0075%, as preservatives

Warning, contains the allergen soy

CAUTION – SKIN IRRITANT

REGISTRATION NO. 33403

PEST CONTROL PRODUCTS ACT

**IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT
1-800-454-2673**

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS: 0.1 L – Bulk

BASF Agricultural Solutions Canada Inc.
510 - 28 Quarry Park Blvd. SE
Calgary, Alberta T2C 5P9
1-877-371-2273

BELYAN is a registered trade-mark of BASF SE, used under license by BASF Agricultural Solutions Canada Inc.

2024-6576
2025-05-01

PRECAUTIONS

1. **KEEP OUT OF REACH OF CHILDREN.**
2. **DO NOT** take internally.
3. May irritate skin. Avoid contact with skin and clothing.
4. Wash exposed areas of skin thoroughly after handling and before eating, drinking or smoking.
5. **DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) or 12 hours.
6. Wear a long-sleeved shirt and long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eye wear (goggles or face shield) during mixing and loading. Gloves are not required during application within a closed cab and/or cockpit.
7. If clothing becomes contaminated, remove and wash separately from household laundry before reuse.
8. Clean spray equipment thoroughly after use.
9. **DO NOT** contaminate domestic or irrigation water, lakes, streams or ponds by the cleaning of equipment or the disposal of wastes.
10. Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

2024-6576
2025-05-01

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

STORAGE

To prevent contamination, store this product away from food or feed.

DISPOSAL

For Recyclable Containers

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For Returnable Containers

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For Refillable Containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For Non-Returnable Containers

Triple or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank. Follow provincial instructions for any required additional cleaning of the container prior to its disposal. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements.

For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

2024-6576
2025-05-01

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

2024-6576
2025-05-01

Booklet

GROUP	3	7	11	FUNGICIDE
-------	---	---	----	-----------

Belyan®
FUNGICIDE

For use in Canola, Rapeseed, Mustard (including oilseed/condiment mustard), Flax, Crop Subgroup 6C (Dried Shelled Peas and Beans, except Soybean), Potato, Soybean and Wheat (All Types)

Suspension Concentrate

COMMERCIAL (AGRICULTURAL)

ACTIVE INGREDIENTS: Mefentrifluconazole 133.3 g/L
Pyraclostrobin 177.8 g/L
Fluxapyroxad88.9 g/L

Contains 5-chloro-2-methyl-4-isothiazolin-3-one at 0.00111 % , 2-methyl-4-isothiazolin-3-one at 0.00037%, 1,2-benzisothiazolin-3-one at 0.0400 % and 2-bromo-2-nitropropane-1,3-diol at 0.0240%, as preservatives

OR

Contains 1,2-benzisothiazolin-3-one and 2-methyl-4-isothiazolin-3-one at 0.0075%, as preservatives,

Warning, contains the allergen soy

CAUTION – SKIN IRRITANT

REGISTRATION NO. 33403

PEST CONTROL PRODUCTS ACT

**IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT
1-800-454-2673**

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS: 0.1 L – Bulk

BASF Agricultural Solutions Canada Inc.
510 - 28 Quarry Park Blvd. SE
Calgary, Alberta T2C 5P9
1-877-371-2273

BELYAN is a registered trade-mark of BASF SE, used under license by BASF Agricultural Solutions Canada Inc.

2024-6576
2025-05-01

GENERAL INFORMATION

Belyan is a broad-spectrum fungicide containing three active ingredients: pyraclostrobin, fluxapyroxad, and mefentrifluconazole. Optimum disease management is achieved when **Belyan** is applied in a regularly scheduled protective spray program and is used in a rotation program with other non-Group 3, non-Group 7, and non-Group 11 fungicides.

MODE OF ACTION

Each of the components in **Belyan** provides a different mode of action against plant pathogenic fungi. Pyraclostrobin belongs to the group of respiration inhibitors classified by the Pest Management Regulatory Agency (PMRA) as Quinone Outside Inhibitors (QoI) which are classified by the Fungicide Resistance Action Committee (FRAC) as Group 11 fungicides. Fluxapyroxad belongs to the group of respiration inhibitors classified by the PMRA as Succinate Dehydrogenase Inhibitors (SDHI) which are classified by FRAC as Group 7 fungicides. Mefentrifluconazole inhibits demethylation of sterol biosynthesis (DMI) which disrupts cell membrane synthesis and are classified by FRAC as Group 3 fungicides.

PRECAUTIONS

1. **KEEP OUT OF REACH OF CHILDREN.**
2. **DO NOT** take internally.
3. May irritate skin. Avoid contact with skin and clothing.
4. Wash exposed areas of skin thoroughly after handling and before eating, drinking or smoking.
5. **DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) or 12 hours.
6. Wear a long-sleeved shirt and long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eye wear (goggles or face shield) during mixing and loading. Gloves are not required during application within a closed cab and/or cockpit.
7. If clothing becomes contaminated, remove and wash separately from household laundry before reuse.
8. Clean spray equipment thoroughly after use.
9. **DO NOT** contaminate domestic or irrigation water, lakes, streams or ponds by the cleaning of equipment or the disposal of wastes.

2024-6576
2025-05-01

10. Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms. Observe buffer zones specified under DIRECTIONS FOR USE.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

Mefentrifluconazole and fluxapyroxad are persistent and may carryover. It is recommended that any products containing mefentrifluconazole or fluxapyroxad not be used in areas treated with this product during the previous season.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

RESTRICTIONS AND LIMITATIONS

1. All crops on this label can be grazed or fed to livestock, except for canola (rapeseed) forage. For livestock feeding and grazing, observe the minimum preharvest intervals specified for each crop under the crop-specific table.
2. Crop Rotation Restrictions: All crops on this label may be planted immediately after the last application. Crops not on this label must not be planted within 1 year after the last application.
3. **DO NOT** use in greenhouses.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

2024-6576
2025-05-01

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

DIRECTIONS FOR USE (See specific sections for each crop group)

<p>All Crops</p> <p>Apply recommended rates of Belyan as instructed in the following series of crop application rate tables. Apply Belyan with spray equipment as specified. Equipment should be checked frequently for proper calibration.</p> <p>Apply before disease develops. See crop-specific requirements for exceptions.</p> <p>Select the use rate based on relative disease pressure. Use the high rate if weather conditions are favourable for disease development (i.e. high humidity/moisture) and/or when risk for disease development is high (i.e. narrow host rotation with disease history and high potential for inoculum).</p> <p>The addition of a spreading/penetrating adjuvant may be used with Belyan to improve coverage and disease control performance, but is not required.</p> <p>DO NOT make more than two (2) sequential applications of Belyan targeting the same disease before alternating to a labeled fungicide containing a different mode of action.</p>
<p>TANK MIXING</p> <p>In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced efficacy or increased host crop injury. The user should contact BASF Agricultural Solutions Canada Inc. ("BASF") at 1-877-371-2273 for information before mixing any pesticide or fertilizer that is not specifically recommended on this label. The user assumes the risk of losses that result from the use of tank mixes that do not appear on this label or that are not specifically recommended by BASF.</p> <p>When applied as a tank-mix combination, read and observe all label directions, including rates, personal protective equipment, restrictions and precautions for each product used in the tank-mix. Always use in accordance with the most restrictive label restrictions and precautions.</p>

2024-6576
2025-05-01

2024-6576
2025-05-01

Canola/Rapeseed (<i>Brassica</i> spp., <i>Brassica napus</i>), mustard (<i>Brassica</i> spp., <i>Brassica juncea</i>) including oilseed/condiment mustard			
Application Rate Table			
(Ground and Aerial Application)			
Disease Controlled	Application Rate (L/ha)	Pre-harvest interval (PHI) (days)	Maximum per year (L/ha)
Blackleg (<i>Leptosphaeria maculans</i>)	0.45 – 0.56	21	1.12
Apply Belyan beginning at the 2 to 6-leaf stage to control blackleg. Apply an additional application at an interval of 10-14 days if disease persists or weather conditions are favourable for disease development.			

Flax			
Application Rate Table			
(Ground and Aerial Application)			
Disease Controlled	Application Rate (L/ha)	PHI and animal feed interval (days)	Maximum per year (L/ha)
Pasmo (<i>Septoria linicola</i>)	0.45 – 0.56	21	1.12
Apply Belyan beginning at 20-50% flowering to control pasmo (<i>Septoria linicola</i>). Apply an additional application at an interval of 10-14 days if disease persists or weather conditions are favourable for disease development.			

2024-6576
2025-05-01

Crop Subgroup 6C (Dried Shelled Peas and Beans, except Soybean) Application Rate Table (Ground and Aerial Application)			
Disease Controlled	Application Rate (L/ha)	PHI and animal feed interval (days)	Maximum per year (L/ha)
Mycosphaerella blight (<i>Mycosphaerella pinodes</i>) Ascochyta blight (<i>Ascochyta</i> spp.) Rust (<i>Uromyces appendiculatus</i>) Powdery mildew (<i>Erysiphe</i> spp.) Anthracnose (<i>Colletotrichum</i> spp.) Asian soybean rust (<i>Phakopsora pachyrhzi</i>) Downy mildew (<i>Peronospora viciae</i> f. sp. <i>pisi</i>) - Suppression	0.56	30	1.12
Apply Belyan at the beginning of flowering or at the onset of symptoms. Apply an additional application at an interval of 10-14 days if disease persists or weather conditions are favourable for disease development.			
Detailed Crop Listing: Crop Subgroup 6-C - Dried shelled peas and beans (except soybean) dried cultivars of bean (dry beans) <i>Lupinus</i> spp. (includes grain lupin, sweet lupin, white lupin and white sweet lupin); <i>Phaseolus</i> spp. (includes field bean, kidney bean, lima bean (dry), navy bean, pinto bean; tepary bean); <i>Vigna</i> spp. (includes adzuki bean, blackeyed pea, catjang, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean; broad bean [dry]; guar; lablab bean; pea; pigeon pea); dry peas (<i>Pisum</i> spp.), lentil, chickpea, and fababean.			

2024-6576
2025-05-01

Soybean Application Rate Table (Ground and Aerial Application)			
Disease Controlled	Application Rate (L/ha)	PHI and animal feed interval (days)	Maximum per year (L/ha)
Frog eye leaf spot (<i>Cercospora sojina</i>) Septoria brown spot (<i>Septoria glycines</i>) Asian soybean rust (<i>Phakopsora pachyrhzi</i>) Cercospora blight & purple seed stain (PSS) (<i>Cercospora kikuchii</i>)	0.56	21	1.12
Apply Belyan prior to disease development when conditions are favourable for disease development. Apply an additional application at an interval of 10-14 days if disease persists or weather conditions are favourable for disease development.			

Potato Application Rate Table (Ground and Aerial Application)			
Disease Controlled	Application Rate (L/ha)	PHI and animal feed interval (days)	Maximum per year (L/ha)
Early Blight (<i>Alternaria solani</i>)	0.56	7	2.24
Begin applications of Belyan prior to disease development. Apply an additional application at an interval of 7-14 days if disease persists or weather conditions are favourable for disease development.			

2024-6576
2025-05-01

Wheat (All Types) Application Rate Table (Ground and Aerial Application)			
Disease Controlled	Application Rate (L/ha)	PHI and animal feed interval (days)	Maximum per year (L/ha)
Septoria leaf blotch (<i>Septoria tritici</i> or <i>Septoria nodorum</i>) Stripe rust (<i>Puccinia striiformis</i>) Leaf rust (<i>Puccinia recondita</i>) Tan spot (<i>Pyrenophora tritici-repentis</i>) Spot blotch (<i>Cochliobolus sativus</i>) Powdery mildew (<i>Erysiphe graminis f. sp. tritici</i>)	0.56	PHI: Apply no later than the end of flowering. Animal feed interval: 21 days.	1.12
Apply Belyan prior to disease development when conditions are favourable for disease development. Apply an additional application at an interval of 10-14 days if disease persists or weather conditions are favourable for disease development.			
Detailed Crop Listing: wheat (spring, winter, and durum)			

MIXING

1. Ensure the spray tank is clean before use. Follow the clean-out recommendations stated on the label of the product that was previously used.
2. Fill the spray tank one-half full of water and start agitation.
3. Add the required amount of **Belyan** to the tank.
4. Add the required amount of the tank mix partner.
5. Add the recommended amount of adjuvant.
6. Continue agitation while filling the remainder of the spray tank .
7. After use, clean the spray tank according to PRECAUTIONS.

APPLICATION INSTRUCTIONS

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

2024-6576
2025-05-01

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Ground Application

Apply **Belyan** at rates listed in the Application Rate and Timing Tables (crop specific) when conditions are favourable for the development of disease. Use a minimum water volume of 100 L/ha and ensure thorough coverage of foliage. Consult nozzle manufacturers recommendation for spray pressures for specific nozzles.

Aerial Application

Apply **Belyan** at rates listed in application rate and timing tables (crop specific) when conditions are favourable for the development of disease. Use a minimum water volume of 50 L/ha. Ensure thorough coverage of foliage. Consult nozzle manufacturers recommendation for spray pressures for specific nozzles.

Instructions for Aerial Applications

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label. Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment. Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted. It is desirable that the pilot have communication capabilities at each treatment site at the time of application. The field crew and the mixer/ loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label. All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-877-371-2273 or obtain technical advice from the distributor or your provincial agricultural representative.

Restrictions

Field Sprayer Application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American

2024-6576
2025-05-01

Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

Aerial application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 km/h at flying height at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotorspan.

Buffer Zones

Spot treatments using hand-held equipment **DO NOT** require a buffer zone.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop		Buffer Zones (metres) Required for the Protection of:			
			Freshwater Habitat of Depths:		Estuarine/Marine Habitat of Depths:	
			Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m
Field sprayer	Canola, rapeseed, mustard and oriental mustard (oilseed/condiment mustard), flax, Crop Subgroup 6C (dried shelled peas and beans), soybean and wheat		5	1	1	1
	Potato		10	1	1	1
Aerial	Canola, rapeseed, mustard and oriental mustard (oilseed/condiment mustard), flax, Crop Subgroup 6C (dried shelled peas and beans), soybean and wheat	Fixed wing	200	10	5	1
		Rotary wing	200	10	2	1
	Potato	Fixed wing	650	15	5	1
		Rotary wing	375	15	2	1

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

2024-6576
2025-05-01

The buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Buffer Zone Calculator on the Pest Management Regulatory Agency web site.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, please note that **Belyan** contains Group 3, 7 and 11 fungicides. Any fungal population may contain individuals naturally resistant to **Belyan** and other Group 3, 7 or 11 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same locations. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

Where possible, rotate the use of **Belyan** with a fungicide from a different mode of action group other than group 3, 7, 11 that controls the same pathogens. **DO NOT** exceed the total number of applications of **Belyan** per season as stated in the Crop-Specific Restrictions and Limitations.

Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.

Where possible, make use of predictive disease models to effectively time fungicide applications.

Monitor treated fungal populations for resistance development. Notify BASF if reduced sensitivity of the pathogen to **Belyan** is suspected.

If disease continues to progress after treatment with this product, do not increase the use rate.

Discontinue use of this product, and switch to another fungicide with a different site of action, if available.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.

For further information and to report suspected resistance, contact BASF at 1-877-371-2273 or www.agsolutions.ca.

STORAGE

To prevent contamination, store this product away from food or feed.

DISPOSAL

For Recyclable Containers

2024-6576
2025-05-01

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For Returnable Containers

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For Refillable Containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For Non-Returnable Containers

Triple or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank. Follow provincial instructions for any required additional cleaning of the container prior to its disposal. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements.

For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.