

GROUP 2 HERBICIDE

SIERRA® 3.0 AG Herbicide

SUSPENSION CONCENTRATE HERBICIDE

COMMERCIAL (AGRICULTURAL)

FOR POSTEMERGENCE APPLICATION ON SPRING WHEAT (INCLUDING DURUM WHEAT) AND WINTER WHEAT FOR CONTROL OF WILD OAT, GREEN FOXTAIL, YELLOW FOXTAIL, BARNYARD GRASS, JAPANESE BROME, VOLUNTEER TAME OAT AND CERTAIN BROADLEAF WEEDS.

FOR SALE AND USE IN MANITOBA, SASKATCHEWAN, ALBERTA AND INTERIOR OF BRITISH COLUMBIA (INCLUDING THE PEACE RIVER REGION OF BRITISH COLUMBIA) ONLY

ACTIVE INGREDIENT:

Flucarbazone (present as flucarbazone-sodium).....200 g/L

**READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

**DANGER: CORROSIVE TO EYES
POTENTIAL SKIN SENSITIZER**

STORE IN A COOL, DRY PLACE

NET CONTENTS: 250 mL to 1000 L

REGISTRATION NO.: **33538**
PEST CONTROL PRODUCTS ACT

Syngenta Canada Inc.
140 Research Lane, Research Park
Guelph, Ontario N1G 4Z3
Telephone: 1-877-964-3682

LABEL

1.0 NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

2.0 FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Do not give **any** liquid to the person. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

3.0 TOXICOLOGICAL INFORMATION

This product contains petroleum distillate - vomiting may cause aspiration pneumonia.

No specific antidote is available. Treat the patient symptomatically.

4.0 PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

Potential skin sensitizer.

Corrosive to eyes. DO NOT get in eyes.

RESTRICTED ENTRY INTERVAL: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and

sprayer settings.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Syngenta Canada Inc. at 1-87-SYNGENTA/1-877-964-3682.

5.0 PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) and chemical-resistant gloves during mixing and loading. Gloves are not required during application within a closed cab and/or cockpit.

Read and follow handling precautions and protective clothing recommendations on broadleaf herbicide and surfactant tank mix partner labels. Follow manufacturer's instructions for cleaning/maintaining PPE (Personal Protective Equipment). If no such instructions for washables, use detergent and hot water. Keep PPE apart from other laundry and wash separately.

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove personal protective equipment immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

6.0 ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

7.0 STORAGE

Store this product away from food or feed. Store in cool, dry place and in such a manner as to prevent cross contamination with other pesticides and fertilizers. Store in original container and out of reach of children, preferably in a locked storage area.

8.0 SPILL CLEAN-UP

Contact Syngenta Canada Inc. (See EMERGENCY NUMBER) and the provincial regulatory agency in case of a spill and for clean-up of spills. Follow safety precautions as directed for handling the product. If the product is clean, it may be used; otherwise, follow the DISPOSAL instructions.

9.0 DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

For Recyclable Containers

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For Refillable Containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

GROUP 2 HERBICIDE

SIERRA® 3.0 AG Herbicide

SUSPENSION CONCENTRATE HERBICIDE

COMMERCIAL (AGRICULTURAL)

FOR POSTEMERGENCE APPLICATION ON SPRING WHEAT (INCLUDING DURUM WHEAT) AND WINTER WHEAT FOR CONTROL OF WILD OAT, GREEN FOXTAIL, YELLOW FOXTAIL, BARNYARD GRASS, JAPANESE BROME, VOLUNTEER TAME OAT AND CERTAIN BROADLEAF WEEDS.

FOR SALE AND USE IN MANITOBA, SASKATCHEWAN, ALBERTA AND INTERIOR OF BRITISH COLUMBIA (INCLUDING THE PEACE RIVER REGION OF BRITISH COLUMBIA) ONLY

ACTIVE INGREDIENT:

Flucarbazone (present as flucarbazone-sodium).....200 g/L

**READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

**DANGER: CORROSIVE TO EYES
POTENTIAL SKIN SENSITIZER**

STORE IN A COOL, DRY PLACE

REGISTRATION NO.: **33538**
PEST CONTROL PRODUCTS ACT

Syngenta Canada Inc.
140 Research Lane, Research Park
Guelph, Ontario N1G 4Z3
Telephone: 1-877-964-3682

PAMPHLET

TABLE OF CONTENTS

	Section Number
NOTICE TO USER	1.0
FIRST AID	2.0
TOXICOLOGICAL INFORMATION	3.0
PRECAUTIONS.....	4.0
PERSONAL PROTECTIVE EQUIPMENT (PPE)	5.0
ENVIRONMENTAL PRECAUTIONS	6.0
STORAGE	7.0
SPILL CLEAN-UP	8.0
DISPOSAL	9.0
PRODUCT INFORMATION	10.0
DIRECTIONS FOR USE	11.0
General Precautions.....	11.1
Ground Applications	11.2
Mixing Instructions	11.2.1
Spraying Instructions	11.2.2
Equipment Clean-Up.....	11.2.3
Aerial Application	11.3
Aerial Use Precautions	11.3.1
Aerial Operator Precautions.....	11.3.2
Product Specific Precautions	11.3.3
Mixing Instructions	11.3.4
Spraying Instructions	11.3.5
Equipment Clean-Up.....	11.3.6
Spray Buffer Zones	11.4
Rotational Crop Restrictions	11.5
CROP USE DIRECTIONS	12.0
Timing of Application.....	12.1
Postemergence Use Directions in Spring, Durum and Winter Wheat.....	12.2
Tank Mixes.....	12.3
RESISTANCE MANAGEMENT RECOMMENDATIONS	13.0

1.0 NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

2.0 FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Do not give **any** liquid to the person. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

3.0 TOXICOLOGICAL INFORMATION

This product contains petroleum distillate - vomiting may cause aspiration pneumonia.

No specific antidote is available. Treat the patient symptomatically.

4.0 PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

Potential skin sensitizer.

Corrosive to eyes. DO NOT get in eyes.

RESTRICTED ENTRY INTERVAL: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and

sprayer settings.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Syngenta Canada Inc. at 1-87-SYNGENTA/1-877-964-3682.

5.0 PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) and chemical-resistant gloves during mixing and loading. Gloves are not required during application within a closed cab and/or cockpit.

Read and follow handling precautions and protective clothing recommendations on broadleaf herbicide and surfactant tank mix partner labels. Follow manufacturer's instructions for cleaning/maintaining PPE (Personal Protective Equipment). If no such instructions for washables, use detergent and hot water. Keep PPE apart from other laundry and wash separately.

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove personal protective equipment immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

6.0 ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

7.0 STORAGE

Store this product away from food or feed. Store in cool, dry place and in such a manner as to prevent cross contamination with other pesticides and fertilizers. Store in original container and out of reach of children, preferably in a locked storage area.

8.0 SPILL CLEAN-UP

Contact Syngenta Canada Inc. (See EMERGENCY NUMBER) and the provincial regulatory agency in case of a spill and for clean-up of spills. Follow safety precautions as directed for handling the product. If the product is clean, it may be used; otherwise, follow the DISPOSAL instructions.

9.0 DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

For Recyclable Containers

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For Refillable Containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (FASTMED)***

10.0 PRODUCT INFORMATION

SIERRA® 3.0 AG Herbicide is a selective herbicide for post-emergence timings in spring wheat (hard red spring, Canada Prairie spring, soft white spring, and extra strong or utility), durum wheat, and winter wheat

SIERRA 3.0 AG Herbicide applied post-emergence must be tank mixed with a surfactant listed on this label. For broader spectrum activity, SIERRA 3.0 AG Herbicide may be tank mixed with a broadleaf herbicide listed on this label. See TANK MIXES section for recommended products.

SIERRA 3.0 AG Herbicide is absorbed by foliage and roots of grass and broadleaf weeds. These weeds cease growth soon after application, removing the competitive effects of susceptible weeds. However, complete weed control may not be seen for one to two weeks.

SIERRA 3.0 AG Herbicide will control wild oat biotypes which have developed resistance to ACCase (Group 1) and triallate (Group 8) herbicides. SIERRA 3.0 AG Herbicide will also control green foxtail biotypes resistant to ACCase (Group 1) and dinitroaniline (Group 3) herbicides.

11.0 DIRECTIONS FOR USE

11.1. General Precautions

Read the entire DIRECTIONS FOR USE before using SIERRA 3.0 AG Herbicide.

- Do not apply post-emergence treatments if it is raining or if rainfall is expected within one (1) hour after application.
- Do not apply this product directly to freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, and wetlands), or estuarine/marine habitats.
- As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.
- Do not contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of waste.
- Do not mix, load or clean spray equipment within 10 metres of well-heads or aquatic systems, including marshes, ponds, ditches, streams, lakes, etc.
- Do not graze treated fields or use green crop for feed. Wheat grain or straw from harvested treated fields may be fed to livestock.
- Do not allow this chemical to drift on to other crops, especially canola, tame oats, or other non target crops.
- Observe minimum interval to harvest of 80 days after treatment.
- Do not treat wheat underseeded to legumes.
- Do not freeze.

11.2. Ground Application

Apply alone or in tank mixes in a spray volume of 50 – 100 L/ha (5 – 10 gallons/acre) at 207-345 kPa (30-50 PSI) pressure to ensure proper weed coverage.

Flat fan nozzles of 80° or 110° are recommended for optimum coverage. Do not use floodjet nozzle or control droplet application equipment. Nozzles may be oriented 45° forward to enhance crop penetration and to give better weed coverage.

Do not apply during periods of dead calm, when winds are gusty or when wind speed is greater than 15 km/h at 2 metres high above ground at the site of application.

Do not apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. Boom height must be 60 cm or less above the crop or ground.

11.2.1. Mixing Instructions

1. Ensure the spray-tank is clean, including spray tank, lines and filters. In-line strainers and nozzle screens should be clean and 50-mesh or coarser.
2. Fill the spray-tank one-third (1/3) to one-half (1/2) full with clean water and begin agitation. Maintain sufficient agitation during both mixing and application.
3. If the tank-mix partner is a dry or granule formulation, add the appropriate amount of the tank mix partner. Agitate until thoroughly mixed.
4. Add the appropriate SIERRA 3.0 AG Herbicide rate directly to the spray-tank. Agitate until SIERRA 3.0 AG Herbicide is thoroughly mixed.
5. Add the remaining tank mix partners. Agitate until thoroughly mixed.
6. Add the adjuvant, then complete filling tank with balance of water needed. Maintain sufficient agitation during both mixing and application.

11.2.2. Spraying Instructions

Unless otherwise specified in the tank mix partner section of this label, apply in a spray volume of 50 – 100 L/ha (5 – 10 gallons/acre) at 207-345 kPa (30-50 PSI) pressure to ensure proper weed coverage.

Flat fan nozzles of 80° or 110° are recommended for optimum coverage. Do not use floodjet nozzle or control droplet application equipment. Nozzles may be oriented 45° forward to enhance crop penetration and to give better weed coverage.

11.2.3. Equipment Clean-Up

1. Drain the tank and thoroughly rinse spray-tank, boom and hoses with clean water. Pay attention to flushing out any visible deposits.
2. Fill the tank with clean water and 1% v/v (1 L/100 L) household ammonia. Flush the hoses, boom and nozzles with the cleaning solution. Circulate for at least 15 minutes. Flush hoses, boom and nozzles once more, then drain the tank.
3. Clean nozzles and screens in a separate container using the ammonia and water.
4. Repeat #2.
5. Rinse tank, boom and hoses with clean water.
 - Do not clean sprayer near desirable vegetation, wells, or other water sources;
 - Dispose of all rinsings in accordance with provincial regulations;
 - Check tank mix partner label for any additional clean-up procedures.

11.3. Aerial Application

Apply alone or in tank mixes (ONLY with the recommended tank mix partners that are registered for aerial use) in no less than 28 L/ha spray volume at a pressure that delivers coarse spray droplets as classified by the ASAE.

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be

applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

To reduce drift caused by turbulent wingtip vortices, the nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotorspan.

11.3.1. Aerial Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the *National Aerial Pesticide Application Manual*, developed by the Federal/Provincial/Territorial committee on Pest Management and Pesticides.

Do not apply to terrain where there is potential for surface run off to enter aquatic systems.

Do not apply to any body of water. Avoid drifting of spray onto any body of water or other non-target areas. Specific buffer zones should be observed.

Do not apply during periods of dead calm. Avoid application of this product when winds are gusty or when wind velocity and direction pose a risk of spray drift.

Do not apply when wind speed is greater than 16 km/h at a flying height at the site of application.

11.3.2. Aerial Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The pilot (aerial applicator) must wear long-sleeved shirt, long pants, socks and shoes.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

11.3.3. Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-877-964-3682 or obtain technical advice from the distributor or your provincial agricultural representative.

11.3.4. Mixing Instructions

Ensure the spray-tank is clean. In-line strainers and nozzle screens should be clean and 50-mesh or coarser.

1. Fill the spray-tank 1/3 to 1/2 full with clean water and begin agitation or bypass.
2. Add the appropriate SIERRA 3.0 AG Herbicide rate directly to the spray-tank. Vigorous agitation is required to dissolve the SIERRA 3.0 AG Herbicide. Maintain sufficient agitation during both mixing and application.
3. Add the broadleaf weed herbicide.
4. Add the surfactant, then complete filling tank with balance of water needed.

11.3.5. Spraying Instructions

Unless otherwise specified in the tank mix partner section of this label, apply in a spray volume no less than 28 L/ha (3 gallons/acre) at no less than 300 kPa pressure to ensure proper weed coverage.

Do not apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) coarse classification. To reduce drift caused by turbulent wingtip vortices, the nozzles distribution along the spray boom length **MUST NOT** exceed 65% of the wing or rotorspan.

11.3.6. Equipment Clean-Up

1. Drain the tank and thoroughly rinse spray-tank, boom and hoses with clean water. Pay particular attention to flushing out any visible deposits.
2. Fill the tank with clean water and 1% v/v (1 L/100 L) household ammonia. Flush the hoses, boom and nozzles with the cleaning solution. Circulate for at least 15 minutes. Flush hoses, boom and nozzles once more, then drain the tank.
3. Clean nozzles and screens in a separate container using the ammonia and water.
4. Repeat #2.
5. Rinse tank, boom and hoses with clean water.
 - Do not clean sprayer near desirable vegetation, wells, or other water sources;
 - Dispose of all rinsings in accordance with provincial regulations;
 - Check tank-mix partner label for any additional clean-up procedures.

11.4. Spray Buffer Zones

Use of the following spray methods or equipment DO NOT require a buffer zone: hand-held or backpack sprayer and spot treatment.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas, and shrublands) and sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

Method of application	Crop		Spray Buffer Zones (metres) Required for the Protection of:		
			Freshwater Habitat of depths		Terrestrial Habitat
			Less than 1 m	Greater than 1 m	
Field sprayer	Spring and winter wheat		2	1	1
Aerial (ASABE coarse spray quality)	Spring and winter wheat	Fixed wing	5	1	20
		Rotary wing	5	2	25
Aerial (ASABE medium spray quality)	Spring and winter wheat	Fixed wing	15	3	40
		Rotary wing	15	4	35

When tank mixes are permitted, consult the labels of the tankmix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Buffer Zone Calculator on the Pesticides portion of the Canada.ca web site. This tool cannot be used to reduce the vegetative filter strips.

11.5. Rotational Crops

The following crops may be planted 11 months after an application of SIERRA 3.0 AG Herbicide.

Soil Zones and Rotational Crops			
Gray-Wooded	Black	Dark Brown	Brown
Spring wheat Barley Canola Field peas*	Spring wheat Barley Canola Field peas* Field bean Flax Durum wheat Soybean Sunflower	Spring wheat Barley Canola Field peas* Flax Durum wheat Soybean Sunflower	Spring wheat

*Field peas may be grown the year following SIERRA 3.0 AG Herbicide application providing the following are all met:

1. Soil pH must be below 7.5,
2. Organic matter must be above 4%, and
3. Precipitation must be equal to or above 10 year average (minimum 100 mm within 60 days of application in year of application).

Or, fields can be summerfallowed.

As SIERRA 3.0 AG Herbicide is degraded by soil microbes, environmental conditions that decrease microbial activity must be considered when making rotational cropping decisions. These environmental conditions include prolonged drought and/or cold temperatures within the following cropping season, as well as soils with both low OM (less than 2%) and high pH (greater than 7.5). If these conditions exist, a soil bioassay may be necessary to ensure rotational crop safety.

Do not plant crops other than those listed above in the year following application of SIERRA 3.0 AG Herbicide on wheat.

12.0 CROP USE DIRECTIONS

12.1. Timing of Application

SIERRA 3.0 AG Herbicide can be applied post-emergence in all types of spring wheat (including durum wheat) and winter wheat.

Research has demonstrated that optimum wheat yield is obtained by early removal of target weeds prior to tillering of wheat. In post-emergence use, apply SIERRA 3.0 AG Herbicide to wheat that has a minimum of one leaf to a maximum of four leaves on the main stem, plus two tillers. Application beyond this stage may result in yield losses due to weed competition.

12.2. Postemergence Use Directions in Spring, Durum and Winter Wheat

Apply SIERRA 3.0 AG Herbicide plus a surfactant once per growing season when the majority of wheat has a minimum of one leaf to a maximum of four leaves on the main stem, plus two tillers. Use a registered non-ionic surfactant such as AGRAL[®] 90, Ag-Surf[®], etc. at a rate of 0.25% v/v (0.25 L per 100 L total spray solution).

Do not apply more than 144 mL/ha of SIERRA 3.0 AG Herbicide (30 g ai/ha) per growing season.

RECOMMENDED APPLICATIONS

CROP APPLICATION TIMING		
CROP	GROWTH STAGE	REMARKS
Spring Wheat Durum Wheat Winter Wheat	1 leaf to 6 total leaves	1 leaf to 4 leaves on main stem, plus 2 tillers

GRASS WEED CONTROL IN WHEAT			
CROP	WEEDS	RATE	REMARKS
Spring Wheat Including: hard red spring, Canada Prairie spring, soft white spring and extra strong (utility) wheat	Green Foxtail (Wild millet)	72 mL/ha	Green foxtail should be no larger than 4-leaf, 2 tillers.
	Weeds controlled at 72 mL/ha plus Wild Oat Volunteer Tame Oat	95 mL/ha	Wild oat should be no larger than 4-leaf, 2 tillers. Use 95 mL/ha when wild oat infestations are below 100 plants/m ² . Volunteer tame oat should be no larger than 4-leaf, 2 tillers.
Durum Wheat Winter Wheat	Weeds controlled at 95 mL/ha plus Wild Oat Japanese Brome Yellow Foxtail* Barnyard Grass*	119 mL/ha	Wild oat, yellow foxtail and barnyard, grass should be no larger than 4-leaf, 2 tillers. Japanese brome should be no larger than 4-leaf. Use 119 mL/ha when wild oat infestations are greater than 100 plants/m ² . Use 119 mL/ha when wild oat, yellow foxtail, barnyard grass and Japanese brome grasses are actively growing under ideal environmental conditions. For control of yellow foxtail and barnyard grass, see "Tank-mixes" section of the label under Inferno® WDG Herbicide.
	Weeds controlled at 119 mL/ha plus Wild Oat Japanese Brome Yellow Foxtail* Barnyard Grass*	144 mL/ha	Wild oat, yellow foxtail and barnyard grass should be no larger than 4-leaf, 2 tillers. Japanese brome should be no larger than 4-leaf. Use 144 mL/ha when wild oat infestations are greater than 100 plants/m ² . Use 144 mL/ha when wild oats, yellow foxtail, barnyard grass and Japanese brome grasses are not actively growing due to poor environmental conditions or when tank mixing with a product that contains dicamba. For control of yellow foxtail and barnyard grass, see "Tank-mixes" section of the label under Inferno WDG Herbicide

*Suppression. For control of yellow foxtail and barnyard grass, see "Tank mixes" section of the label under Inferno WDG Herbicide.

BROADLEAF WEED CONTROL IN WHEAT		
WEED	RATE	REMARKS
Redroot Pigweed (<i>Amaranthus retroflexus</i>)	95 mL/ha	For control of redroot pigweed, apply SIERRA 3.0 AG Herbicide to weeds at the 2- to 6-leaf stage.
Wild Mustard (<i>Brassica kaber</i>)		For control of wild mustard apply SIERRA 3.0 AG Herbicide to weeds at the 2- to 6-leaf stage.
Stinkweed (<i>Thlaspi arvense</i>)		For control of stinkweed apply SIERRA 3.0 AG Herbicide to weeds at the 2- to 9-leaf stage.
Volunteer Canola (<i>Brassica napus</i>)		For control of volunteer canola apply SIERRA 3.0 AG Herbicide to weeds at the 2- to 6-leaf stage. Will not control imidazolinone-tolerant (Clearfield®) canola varieties.
Green Smartweed (<i>Polygonum scabrum</i> Moench.)		For control of green smartweed apply SIERRA 3.0 AG Herbicide to weeds at the 2- to 6-leaf stage.
Shepherd's Purse (<i>Capsella bursa-pastoris</i>)		For control of shepherd's purse, apply SIERRA 3.0 AG Herbicide to weeds at the 2- to 6-leaf stage.
Wild Buckwheat (<i>Fallopia confolifulus</i>)	119 – 144 mL/ha	For suppression of wild buckwheat, apply SIERRA 3.0 AG Herbicide to weeds at the 1- to 4-leaf stage. Apply at the highest rate when wild buckwheat is not actively growing due to poor environmental conditions.

Wheat exposed to water-logged or saturated soils, or temperature extremes such as heat or freezing weather, or drought, low fertility or plant disease at application time could show unacceptable injury symptoms. Weed control also may be reduced by these same conditions.

12.3. Tank Mixes

For broader spectrum control of both annual grasses and broadleaf weeds, SIERRA 3.0 AG Herbicide may be mixed with one of the broadleaf herbicides listed in the following tables.

When tank-mixes are permitted, read and observe all label directions, including rates and restrictions for each product used in the tank mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.

Refer to previous tables for the appropriate rate and application timing for SIERRA 3.0 AG Herbicide. Use a registered non-ionic surfactant (0.25% v/v or 0.25 L/100 L total spray solution) with SIERRA 3.0 AG Herbicide tank mixes.

APPROVED TANK MIX PARTNERS IN SPRING WHEAT	
BROADLEAF HERBICIDE TANK MIX PARTNER	REMARKS
2,4-D Amine	Use up to 560 g ae/ha 2,4-D Amine Herbicide.
2,4-D Ester	Use up to 560 g ae/ha 2,4-D Ester Herbicide.
Akito® Herbicide	
Akito Herbicide + 2,4-D Ester 700	Use up to 560 g ae/ha 2,4-D Ester 700 Herbicide.
Akito Herbicide + MCPA Ester 600	Use up to 560 g ae/ha MCPA Ester 600 Herbicide.
Ally® Herbicide + 2,4-D Amine	Addition of a second surfactant is not required. Apply tank mix in at least 100 L of water per hectare (10 gallons/acre) if 95 or 144 mL/ha rate of SIERRA 3.0 AG Herbicide is used.
Ally Herbicide + 2,4-D Ester	Addition of a second surfactant is not required. Apply tank mix in at least 100 L of water per hectare (10 gallons/acre) if 95 or 144 mL/ha rate of SIERRA 3.0 AG Herbicide is used.
Attain™ XC A Herbicide	Apply tank mix in at least 100 L of water per hectare (10 gallons/acre) if 95 or 144 mL/ha rate of SIERRA 3.0 AG Herbicide is used.
Buctril® M/Logic® M Liquid Herbicide	
Bromoxynil products	Products containing bromoxynil as the solo active ingredient only.
Bromoxynil + MCPA Ester	Products containing bromoxynil as the solo active ingredient only.
Bromoxynil + 2,4-D Ester	Products containing bromoxynil as the solo active ingredient only.
Curtail® M Herbicide	Apply tank mix in at least 100 L of water per hectare (10 gallons/acre) if 95 or 144 mL/ha rate of SIERRA 3.0 AG Herbicide is used.
DyVel® Herbicide	Reduction in wild oat control may be observed with this tank mix partner. Use 144 mL/ha rate of SIERRA 3.0 AG Herbicide for control of wild oat. Do not use 95 mL/ha rate of SIERRA 3.0 AG Herbicide.
Enforcer™ D Herbicide	
Enforcer M Herbicide	
Evito® 480 SC Fungicide	
Frontline™ 2,4-D XC Herbicide Tank Mix	Frontline 2,4-D XC Herbicide consists of Frontline 2,4-D XC A Herbicide plus Frontline 2,4-D XC B Herbicide
Frontline XL Herbicide	Apply tank mix in at least 100 L of water per hectare (10 gallons/acre).
Inferno WDG Herbicide	For control of yellow foxtail and barnyard grass, tank-mix Inferno WDG Herbicide with 95 - 119 mL/ha of SIERRA 3.0 AG Herbicide. Apply at the higher rate when yellow foxtail and barnyard grass are not actively growing due to poor environmental conditions.
Infinity Herbicide / Infinity FX	
IPCO® Dichlorprop-DX Herbicide	
MCPA Amine	Use up to 560 g ae/ha MCPA Amine Herbicide.
MCPA Ester	Use up to 560 g ae/ha MCPA Ester Herbicide.
Momentum	
OcTTain™ XL Herbicide	Apply tank mix in at least 100 L of water per hectare (10 gallons/acre).
Optica™ Trio	Reduction in wild oat control may be observed with this tank mix partner. For control of high wild oat populations, use 144 mL/ha rate of SIERRA 3.0 AG Herbicide.

APPROVED TANK MIX PARTNERS IN SPRING WHEAT	
BROADLEAF HERBICIDE TANK MIX PARTNER	REMARKS
Prestige™ XC Herbicide Tank Mix	Prestige XC Herbicide Tank Mix consists of Prestige XC A Herbicide plus Prestige SC B Herbicide. Apply tank mix in at least 100 L of water per hectare (10 gallons/acre) if 95 or 144 mL/ha rate of SIERRA 3.0 AG Herbicide is used.
Pixxaro™ A Herbicide	
Pixxaro A Herbicide + MCPA Ester	
Pixxaro A Herbicide + Curtail M Herbicide	
Refine® SG Herbicide / Deploy® WDG Herbicide + 2,4-D Amine	Addition of a second surfactant is not required.
Refine SG Herbicide / Deploy WDG Herbicide + 2,4-D Ester	Addition of a second surfactant is not required.
Refine SG Herbicide / Deploy WDG Herbicide	Addition of a second surfactant is not required.
Roxar™ Fungicide	
Spectrum™ Herbicide Tank Mix	Spectrum Herbicide Tank Mix consists of Spectrum A Herbicide plus Spectrum B Herbicide
Stellar™ XL Herbicide	Apply tank mix in at least 100 L of water per hectare (10 gallons/acre) if 95 or 144 mL/ha rate of SIERRA 3.0 AG Herbicide is used.
Thumper® Emulsifiable Selective Weedkiller / IPCO Leader® Liquid Herbicide	
Triton™ C Herbicide	Addition of a second surfactant is not required.
Ohm™	Apply at 1 L/ha with SIERRA 3.0 AG Herbicide alone or in a tank mixture with Akito Herbicide.
Wave™	Apply at 150 mL/ha with SIERRA 3.0 AG Herbicide alone or in a tank mixture with Akito Herbicide.

APPROVED TANK MIX HERBICIDE PARTNERS IN DURUM WHEAT	
BROADLEAF HERBICIDE TANK MIX PARTNERS	REMARKS
2,4-D Amine	Use 420 g ae/ha 2,4-D Amine Herbicide. Apply tank mix in at least 100 L of water per hectare (10 gallons/acre).
2,4-D Ester	Use 420 g ae/ha 2,4-D Ester Herbicide. Apply tank mix in at least 100 L of water per hectare (10 gallons/acre).
Akito Herbicide	
Akito Herbicide + 2,4-D Ester 700	Use up to 560 g ae/ha 2,4-D Ester 700 Herbicide.
Akito Herbicide + MCPA Ester 600	Use up to 560 g ae/ha MCPA Ester 600 Herbicide.
Enforcer D Herbicide	
Enforcer M Herbicide	
Inferno WDG Herbicide	For control of yellow foxtail and barnyard grass, tank mix Inferno WDG Herbicide with 95 - 119 mL/ha of SIERRA 3.0 AG Herbicide. Apply at the higher rate when yellow foxtail and barnyard grass are not actively

	growing due to poor environmental conditions.
Pixxaro A Herbicide	
Pixxaro A Herbicide + MCPA Ester	
Pixxaro A Herbicide + Curtail M Herbicide	
Spectrum Herbicide Tank Mix	Spectrum Herbicide Tank Mix consists of Spectrum A Herbicide plus Spectrum B Herbicide. Apply tank mix in at least 100 L of water per hectare (10 gallons/acre).
Ohm	Apply at 1 L/ha with SIERRA 3.0 AG Herbicide alone or in a tank mixture with Akito Herbicide.
Wave	Apply at 150 mL/ha with SIERRA 3.0 AG Herbicide alone or in a tank mixture with Akito Herbicide.

APPROVED TANK MIX HERBICIDE PARTNERS IN WINTER WHEAT	
BROADLEAF HERBICIDE TANK MIX PARTNERS	REMARKS
2,4-D Amine	Use up to 560 g ae/ha 2,4-D Amine Herbicide.
2,4-D Ester	Use up to 560 g ae/ha 2,4-D Ester Herbicide.
Akito Herbicide	
Akito Herbicide + 2,4-D Ester 700	Use up to 560 g ae/ha 2,4-D Ester 700 Herbicide.
Akito Herbicide + MCPA Ester 600	Use up to 560 g ae/ha MCPA Ester 600 Herbicide.
Buctril M /IPCO Logic M Liquid Herbicide	
Evito 480 SC Fungicide	
MCPA Amine	Use up to 560 g ae/ha MCPA Amine Herbicide.
MCPA Ester	Use up to 560 g ae/ha MCPA Ester Herbicide.
Refine SG Herbicide / Deploy WDG Herbicide	Addition of a second surfactant is not required.
Roxar Fungicide	
Ohm	Apply at 1 L/ha with SIERRA 3.0 AG Herbicide alone.
Wave	Apply at 150 mL/ha with SIERRA 3.0 AG Herbicide alone.

13.0 RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, SIERRA 3.0 AG Herbicide is a Group 2 Herbicide. Any weed population may contain or develop plants naturally resistant to SIERRA 3.0 AG Herbicide and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

Where possible, rotate the use of SIERRA 3.0 AG Herbicide or other Group 2 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.

Use tank mixtures with herbicides from a different group when such use is permitted. To delay

resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.

Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other chemical control practices.

Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.

Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.

For further information or to report suspected resistance, contact Syngenta Canada Inc. at 1-87-SYNGENTA (1-877-964-3682) or at www.syngenta.ca.

AGRAL[®] and SIERRA[®] are trademarks of a Syngenta Group Company.
Ag-Surf[®], Akito[®], Ally[®], Attain[™], Buctril[®], Clearfield[®], Curtail[®], Deploy[®], DyVel[®], Enforcer[™], Evito[®], Frontline[™], Inferno[®], IPCO[®], Leader[®], Logic[®], OcTTain[™], Ohm[™], Optica[™], Pixxaro[™], Prestige[™], Refine[®], Roxar[™], Spectrum[™], Stellar[™], Thumper[®], Triton[™] and Wave[™] are trademarks of their respective companies.