

2024-6576
2025-05-01

Container

| | | |
|-------|---|-----------|
| GROUP | 2 | HERBICIDE |
|-------|---|-----------|

SOLO® ADV II

HERBICIDE

SOLUTION

COMMERCIAL (AGRICULTURAL)

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND INTERIOR OF BRITISH COLUMBIA
(INCLUDING THE PEACE RIVER REGION) ONLY

ACTIVE INGREDIENT: Imazamox25 g/L

REGISTRATION NO. 33720

PEST CONTROL PRODUCTS ACT

CAUTION – SKIN IRRITANT

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

**IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT
1-800-454-2673**

NET CONTENTS: 1 L - 1000 L, Bulk

BASF Agricultural Solutions Canada Inc.
510 - 28 Quarry Park Blvd. SE
Calgary, Alberta T2C 5P9
1-877-371-2273

SOLO is a registered trade-mark of BASF SE, used under license by BASF Agricultural
Solutions Canada Inc.

2024-6576

2025-05-01

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

DO NOT take internally. Harmful if swallowed.

May irritate skin. Avoid contact with skin.

Avoid breathing spray and mist. Use with adequate ventilation.

Wash thoroughly after handling and before eating, drinking or smoking.

Wear a long-sleeved shirt and long pants during all activities, and wear chemical-resistant gloves during mixing/loading, clean-up and repair activities.

If clothing becomes contaminated, remove immediately and wash separately from household laundry before reuse.

Do not enter or allow workers entry into treated areas during the restricted entry interval (REI) of 12 hours.

Apply only to agricultural crops when the potential for drift to areas of human habitation and areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

2024-6576

2025-05-01

TOXICOLOGICAL INFORMATION

Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS

TOXIC to non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

STORAGE

1. Shake well before use.
2. Store this product away from food or feed. Store in original, tightly-closed container.
3. Do not ship or store the product near food, feed, seed or fertilizers.
4. Store in a cool, dry, locked, well-ventilated area without floor drain.
5. Herbicides should be shipped or stored separately from other pesticides to avoid cross-contamination.
6. Protect from freezing. **SOLO ADV II** will freeze at -5°C . If this product has been stored where freezing temperatures have occurred, thaw the product completely at room temperature, then shake well before use.

DISPOSAL

Recyclable Containers

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

2024-6576

2025-05-01

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

Refillable Containers

For disposal, this empty container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

2024-6576
2025-05-01

Booklet

| | | |
|-------|---|-----------|
| GROUP | 2 | HERBICIDE |
|-------|---|-----------|

SOLO® ADV II

HERBICIDE

SOLUTION

COMMERCIAL (AGRICULTURAL)

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND INTERIOR OF BRITISH COLUMBIA
(INCLUDING THE PEACE RIVER REGION) ONLY

ACTIVE INGREDIENT: Imazamox25 g/L

REGISTRATION NO. 33720 PEST CONTROL PRODUCTS ACT

CAUTION – SKIN IRRITANT
READ THE LABEL AND BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

**IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT
1-800-454-2673**

NET CONTENTS: 1 L - 1000 L, Bulk

BASF Agricultural Solutions Canada Inc.
510 - 28 Quarry Park Blvd. SE
Calgary, Alberta T2C 5P9
1-877-371-2273

SOLO is a registered trade-mark of BASF SE, used under license by BASF Agricultural
Solutions Canada Inc.

2024-6576

2025-05-01

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

DO NOT take internally. Harmful if swallowed.

May irritate skin. Avoid contact with skin.

Avoid breathing spray and mist. Use with adequate ventilation.

Wash thoroughly after handling and before eating, drinking or smoking.

Wear a long-sleeved shirt and long pants during all activities, and wear chemical-resistant gloves during mixing/loading, clean-up and repair activities.

If clothing becomes contaminated, remove immediately and wash separately from household laundry before reuse.

Do not enter or allow workers entry into treated areas during the restricted entry interval (REI) of 12 hours.

Apply only to agricultural crops when the potential for drift to areas of human habitation and areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

2024-6576
2025-05-01

TOXICOLOGICAL INFORMATION

Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS

TOXIC to non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

STORAGE

1. Shake well before use.
2. Store this product away from food or feed. Store in original, tightly-closed container.
3. Do not ship or store the product near food, feed, seed or fertilizers.
4. Store in a cool, dry, locked, well-ventilated area without floor drain.
5. Herbicides should be shipped or stored separately from other pesticides to avoid cross-contamination.
6. Protect from freezing. **SOLO ADV II** will freeze at -5°C . If this product has been stored where freezing temperatures have occurred, thaw the product completely at room temperature, then shake well before use.

DISPOSAL

Recyclable Containers

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

2024-6576

2025-05-01

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

Refillable Containers

For disposal, this empty container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

2024-6576

2025-05-01

GENERAL INFORMATION

SOLO ADV II is a selective herbicide that can be applied as an early post-emergence treatment in **Clearfield®** Lentils¹ (lentil varieties with the **Clearfield** trait), **Clearfield** Sunflowers¹ (sunflower varieties with the **Clearfield** trait) and soybeans.

When **SOLO ADV II** is applied early post-emergence, absorption may occur through both the roots and foliage. Susceptible weeds stop growing and eventually die.

Use of **SOLO ADV II** on sunflower or lentil varieties not designated with the **Clearfield** trait will cause severe injury.

DIRECTIONS FOR USE

DO NOT apply more than once per year.

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Field Sprayer Application

DO NOT apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets finer than the American Society of Agricultural and Biological Engineers (ASABE) S572 (572.1 to 572.3) medium classification. Boom height must be 60 cm or less above the crop or ground.

DO NOT apply by air.

Spray Buffer Zones

A spray buffer zone is **NOT** required for uses with hand-held application equipment permitted on this label.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands).

| Method of Application | Crop | Spray Buffer Zones (metres) Required for the Protection of Terrestrial Habitat |
|-----------------------|-----------|--|
| Field sprayer | All crops | 1 |

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASABE) category indicated on the labels for those tank mix partners.

2024-6576

2025-05-01

The spray buffer zones for this product for conventional application equipment can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Drift Management portion of the Canada.ca website.

Registered Crops and Application Timing

SOLO ADV II may be applied as an early post-emergence treatment to the following crops. Applications should be made at the appropriate crop staging as indicated below and after weeds have emerged. To control broadleaf weeds, apply **SOLO ADV II** from the cotyledon to the 4-leaf stage. To control grasses, apply **SOLO ADV II** from the 1 to 4-true leaf stage up until early tillering.

| Crop | Leaf Stage |
|------------------------------------|---------------------------|
| Clearfield Lentils ¹ | 1-9 node stage |
| Clearfield Sunflowers ¹ | 2 to 8 true leaf stage |
| Soybeans | cotyledon to 4-leaf stage |

Application Rates and Weeds Controlled

Apply **SOLO ADV II** in a water volume of 100 L/ha for control of broadleaf and grass weeds as indicated below. Initial transient crop yellowing may be observed after application but this is usually outgrown and should not affect crop yield.

| SOLO ADV II at 0.8 L/ha will control: | |
|--|--|
| Barnyard grass | Volunteer barley |
| Cleavers* | Volunteer canary seed |
| Cow cockle | Volunteer canola (non- Clearfield tolerant canola only) |
| Downy brome** | Volunteer durum wheat (non- Clearfield tolerant wheat) |
| Green foxtail | Volunteer spring wheat (non- Clearfield tolerant wheat) |
| Green smartweed | Volunteer tame oats |
| Japanese brome grass* | Wild buckwheat* |
| Kochia* | Wild mustard |
| Lamb's-quarters | Wild oats |
| Persian dandelion | Yellow foxtail |
| Redroot pigweed | |
| Round-leaved mallow* | |
| Russian thistle | |
| Shepherd's purse | |
| Stinkweed | |

* suppression only

** including glyphosate-resistant biotypes

Tank Mix Options

To provide enhanced weed control, **SOLO ADV II** may be tank-mixed with the following tank-mix partners.

This product may be tank mixed with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions for Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact BASF at 1-877-371-2273 for information before applying any tank mix that is not specifically recommended on this label.

SOLO ADV II + POAST® ULTRA Liquid Emulsifiable Herbicide

SOLO ADV II may be tank-mixed with POAST ULTRA Liquid Emulsifiable Herbicide and applied to **Clearfield** Lentils¹ (lentil varieties with the **Clearfield** trait), **Clearfield** Sunflowers¹ (sunflower varieties with the **Clearfield** trait) and soybeans.

Apply to broadleaf weeds from the cotyledon to 4-leaf stage and to grassy weeds from 1-6 true leaf or up to 2 tillers.

| | |
|-------------------------|--|
| Rate | 0.8 L/ha SOLO ADV II + POAST ULTRA Liquid Emulsifiable Herbicide at 0.32 to 0.47 L/ha |
| Water Volume | 100 L/ha |
| Weeds Controlled | <p>This tank-mix combination will provide control of the following weeds in addition to those listed in the Application Rates and Weeds Controlled section of this label. Apply POAST ULTRA Liquid Emulsifiable Herbicide at the 0.47 L/ha rate when weed densities are high and overlapping, when staging is late, or when weeds are under stress and not growing as actively due to moisture stress or temperature stress.</p> <ul style="list-style-type: none"> Crabgrass (large) Fall panicum Green foxtail ¹ (including Group 1 and 2 resistant) Proso millet Quack grass (suppression, 2 to 5 leaf) ² Volunteer corn Volunteer wheat (including Clearfield) Wild oats ¹ (including Group 1 and 2 resistant) Witchgrass Yellow foxtail |

¹ The tank mix of **SOLO ADV II** + POAST ULTRA Liquid Emulsifiable Herbicide will not control weed biotypes that are multiple-resistant to both Group 1 and Group 2 Herbicides.

² Apply 0.38 – 0.47 L/ha of POAST ULTRA Liquid Emulsifiable Herbicide.

2024-6576
2025-05-01

SOLO ADV II + Caziva® Ultra Q, Assure® II Herbicide or Yuma® GL Liquid EC Herbicide

SOLO ADV II may be tank-mixed with Caziva Ultra Q, Assure II Herbicide or Yuma GL Liquid EC Herbicide and applied to **Clearfield** Lentils¹ (lentil varieties with the **Clearfield** trait), **Clearfield** Sunflowers¹ (sunflower varieties with the **Clearfield** trait) and soybeans as a post-emergent application. Apply after weeds have emerged and from the 1-9 node stage of **Clearfield** Lentils¹; 2 to 8 true leaf stage of **Clearfield** Sunflowers¹; and cotyledon to 4-leaf stage of soybeans.

Application Rate

Apply in 100 L/ha of water. Use ground equipment only.

| Crops | SOLO ADV II (L/ha) | Caziva Ultra Q, Assure II or Yuma GL Liquid EC Herbicide (L/ha) |
|---|--------------------|---|
| Clearfield Lentils ¹ , Clearfield Sunflowers ¹ , soybeans | 0.8 | 0.38 – 0.75 |

Weeds Controlled

This tank-mix combination will provide control of the grass weeds listed below in addition to the broadleaf weeds listed in the Application Rates and Weeds Controlled section of this label. Refer to the tank mix partner product labels for weed stages of control.

| Caziva Ultra Q ¹ | Grass Weeds Controlled |
|-----------------------------|--|
| 0.38 L/ha | Barnyard grass Downy brome Foxtail barley Green foxtail ² Japanese brome Persian darnel Volunteer barley Volunteer canary seed Volunteer corn Volunteer tame oats Volunteer wheat (spring and durum) ³ Yellow foxtail Wild oats ² |
| 0.50 L/ha | <u>Those controlled above by 0.38 L/ha and:</u> Fall panicum Old witchgrass Proso millet Quackgrass (suppression) |
| 0.75 L/ha | <u>Those controlled above and:</u> Quackgrass |

¹ Assure II or Yuma GL Liquid EC Herbicide may be substituted for Caziva Ultra Q at same product rate.

² including Group 1 or Group 2 resistant biotypes.

³ including Clearfield tolerant varieties.

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than BASF Agricultural Solutions Canada Inc. ("BASF") under the User Requested Minor Use Label Expansion program. For these uses, BASF has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

Imidazolinone-tolerant Chickpeas (chickpea varieties designated as imidazolinone-tolerant)

Apply only to imidazolinone-tolerant chickpeas. **Use of SOLO ADV II on chickpea varieties not designated with the imidazolinone trait will cause severe injury.** Growers should consult the seed guide and supplier at point of purchase for tolerant varieties.

Applications should be made at the appropriate crop staging as indicated below and after weeds have emerged. To control broadleaf weeds, apply **SOLO ADV II** from the cotyledon to the 4-leaf stage. To control grasses, apply **SOLO ADV II** from the 1 to 4-true leaf stage up until early tillering.

| | |
|---------------------------|---|
| Rate | 0.8 L/ha SOLO ADV II |
| Water Volume | 100 L/ha |
| Application Timing | Early post-emergence, 1 – 6 node of chickpea |
| Weeds Controlled | Labelled weeds as listed in the Application Rates and Weeds Controlled section of this label. |
| PHI | 60 days |

PRE-HARVEST AND GRAZING INTERVALS (PHI)

The following pre-harvest and grazing intervals should be observed for respective crops.

| Crop | PHI (days) |
|---|-------------------|
| Clearfield Lentils ¹ | 60 |
| Clearfield Sunflowers ¹ | 70 |
| Soybeans | 60 |

DO NOT graze the treated **Clearfield** Lentils¹ or soybeans or cut for hay within 20 days of application.

DO NOT graze treated sunflower plants or cut for straw; sufficient data are not available to support such use.

RESTRICTIONS AND LIMITATIONS

1. **DO NOT** apply when weather conditions may cause spray drift from treated fields to adjacent crops. Clean sprayer thoroughly after use to avoid damage to the next crop sprayed.
2. **DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.
3. **DO NOT** apply to any crops that have been subjected to stress from conditions such as frost, hail damage, flooding, drought, hot, humid weather, widely fluctuating temperature conditions, prolonged cold weather or injury from prior herbicide applications, as crop injury may result.
4. Apply only to agricultural crops when the potential for drift to areas of human habitation and areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

SPRAYING INSTRUCTIONS

Select proper nozzle to avoid spraying fine mist. For best results, use sprayers equipped with flat fan or similar nozzles to ensure coverage. Apply in a spray volume of 100 L/ha and at a pressure of 275 kPa. Better coverage of the product results in enhanced control of weeds.

MIXING INSTRUCTIONS

1. Shake well before use. When applying **SOLO ADV II**, always start with a clean sprayer. Refer to previously applied product labels for specific cleaning instructions.
2. Fill the clean spray tank with half to three-quarters of the required amount of clean water and start agitation system. Agitation should be running during the entire mixing procedure.
3. Add the correct amount of **SOLO ADV II** and continue to agitate.
4. If required, add the required amount of POAST ULTRA Liquid Emulsifiable Herbicide, Caziva Ultra Q, Assure II Herbicide or Yuma GL Liquid Herbicide while agitating the spray solution.
5. Continue agitation while filling the spray tank with the remaining amount of water.
6. Maintain continuous and constant agitation throughout application until spraying is complete.
7. Upon completion of spraying **SOLO ADV II**, thoroughly flush tank, boom, hoses and in-line and nozzle screens with clean water to avoid possible injury to other crops.

2024-6576

2025-05-01

FOLLOW CROPPING

Winter wheat can be planted 3 months after treatment (3 MAT) as a rotational crop.*

Initial crop injury to non-**Clearfield** canola may be observed. Avoid spray overlap as yield reduction may result. The following crops may be grown safely the year following an application:

- Canary seed*
- Field corn
- Field peas
- Clearfield** Canola ¹
- Non-**Clearfield** canola*
- Lentils
- Spring wheat
- Durum wheat*
- Spring barley
- Soybeans
- Sunflower
- Tame oats*
- Flax*
- Chickpeas
- Clearfield** Sunflowers ¹

The following crop may be grown safely two years following an application:

- Mustard (condiment type only)*

* If drought conditions are experienced between June 1 and September 1 in the year of application, delay planting of winter wheat, durum wheat, canary seed, tame oats, flax and canola (non-**Clearfield**) by an additional year. If drought is received between June 1 and September 1 in the year of application OR between June 1 and September 1 in the year following application, delay planting of mustard by an additional year.

In the grey, black, and dark brown soil zones, drought is defined as less than 125 mm of total precipitation between June 1 and September 1. In the brown soil zone, drought is defined as less than 125 mm of total precipitation between June 1 and September 1, OR less than 15 mm of precipitation in any month between June 1 and September 1.

There are insufficient data for other follow crops. Conduct a field bioassay (a test strip grown to maturity) the year before growing any crop other than those listed above.

WARNING: Certain environmental conditions may delay the breakdown of herbicide residues in soil. These conditions include but are not limited to drought, extremes in soil pH (less than 6.5), low organic matter soils (less than 3%), extremely coarse textured (sandy) soils, and excessive cold. Under these conditions, the level of phytotoxic herbicide residues present in the field the season following an application may result in an increased potential for injury to succeeding crops to occur. This potential for increased residues under these environmental conditions is not unique to any specific herbicide or herbicide group but is a property of those herbicides which persist in the soil and are dependent on soil microbial activity and other non-microbial processes (e.g. hydrolysis) to breakdown.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, **SOLO ADV II** herbicide is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to **SOLO ADV II** herbicide and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of **SOLO ADV II** herbicide or other Group 2 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact BASF at 1-877-371-2273 or at www.agsolutions.ca.

CAZIVA is a registered trade-mark of BASF SE, used under license by BASF Agricultural Solutions Canada Inc.

Clearfield is a registered trade-mark of BASF Agrochemical Products B.V. and POAST is a registered trade-mark of BASF SE, all used under license by BASF Agricultural Solutions Canada Inc.

¹ *Methods of growing plants which are resistant to certain acetohydroxyacid synthase (AHAS) herbicides are protected by Canadian Patent No. 1341465. Other applicable patents include Canadian Patent Application No. 2447445. The use of this product in the practice of patented*

2024-6576
2025-05-01

methods could constitute patent infringement. Further information can be obtained by contacting BASF Agricultural Solutions Canada Inc. at 1-877-371-2273.