

2025-0049  
2025-03-26

This label has been updated according to the re-evaluation decision for Tebuconazole, RVD2024-09. While users are encouraged to follow this updated label immediately, the previously approved label is valid until 2026-11-20 in accordance with the phase out period set out in the re-evaluation decision for Tebuconazole, RVD2024-09. This previously approved label will be provided upon request by emailing [hc.pmra.info-arla.sc@canada.ca](mailto:hc.pmra.info-arla.sc@canada.ca). In your email please include the product name and Registration number of the label you are requesting.

Label

# Toledo 430 SC Foliar Fungicide

## SUSPENSION

**FOR SUPPRESSION OF *FUSARIUM* HEAD BLIGHT (SCAB) AND CONTROL OF FOLIAR DISEASES ON WHEAT (SPRING, WINTER AND DURUM), BARLEY, OATS, SOYBEAN AND CERTAIN MINOR USE CROPS**

## COMMERCIAL

ACTIVE INGREDIENT: Tebuconazole ..... 430 g/L

Contains 1,2-benzisothiazolin-3-one at 0.021% as a preservative.

REGISTRATION NO. 34339 PEST CONTROL PRODUCTS ACT

**READ THIS LABEL AND BOOKLET BEFORE USING**

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION: EYE IRRITANT**

**DANGER: SKIN IRRITANT**

NET CONTENTS: 0.03 L to 200 L (BULK)

Albaugh, LLC  
1525 NE 36th Street  
Ankeny, IA 50021  
1-800-247-8013

For medical or treatment information from exposure to this product, call 1-888-347-6732 (7 days/week, 24-hr).

For 24-hour chemical spill, leak, fire, exposure, or accident response information, call CHEMTREC toll free at 1-800-424-9300.

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

## **PRECAUTIONS**

**KEEP OUT OF REACH OF CHILDREN.** Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist.

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and boots during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab or cockpit.

For application using handheld airblast/mistblower equipment, wear chemical-resistant coveralls with a chemical-resistant hood over long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear and a respirator with a NIOSH-approved organic-vapour-removing cartridge with a prefilter approved for pesticides OR a NIOSH-approved canister approved for pesticides.

DO NOT handle more than 140 ml per person, per day when using a handheld airblast/mistblower. These restrictions are in place to minimize exposure to individual applicators. Application may need to be performed over multiple days or using multiple applicators.

When mixing, loading and applying using a mechanically-pressurized handgun, wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes, and a respirator with a NIOSH-approved organic-vapour removing cartridge with a prefilter approved for pesticides, or a NIOSH-approved canister approved for pesticides during mixing, loading, application, cleanup and repair.

Follow manufacturer's instructions for cleaning/maintaining the Personal Protective Clothing (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

**DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

When tank-mixes are permitted, read and observe all label directions, including rates and restrictions for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.

### **User Safety Recommendations:**

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**FIRST AID**

**In case of poisoning**, call physician or Poison Control Centre immediately. Have patient lie down and keep quiet.

**IF SWALLOWED:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION:**

The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.

**To Physician:** No specific antidote. Treat symptomatically.

**ENVIRONMENTAL PRECAUTIONS**

Toxic to birds, small wild mammals, aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

Tebuconazole is persistent and may carryover. It is recommended that this product not be used in areas treated with any products containing tebuconazole during the previous season.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

**STORAGE:** Store this product away from food or feed. Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

**LEAK OR SPILL PROCEDURE:** Handle and open container in a manner as to prevent spillage. If container is leaking invert to prevent leakage. If the container is leaking or material is spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away. For decontamination procedures or any other assistance that may be necessary, you may contact Albaugh, LLC at the 1-800-247-8013.

**DISPOSAL:**

**Recyclable Container:** Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

**Returnable Container:** Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

**Refillable Container:** For disposal, this empty container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not use this container for any other purpose.

**Disposal of Unused, Unwanted Product:**

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills.

Booklet label

## Toledo 430 SC Foliar Fungicide

Group	<b>3</b>	Fungicide
-------	----------	-----------

### SUSPENSION

**FOR SUPPRESSION OF *FUSARIUM* HEAD BLIGHT (SCAB) AND CONTROL OF FOLIAR DISEASES ON WHEAT (SPRING, WINTER AND DURUM), BARLEY, OATS, SOYBEAN AND CERTAIN MINOR USE CROPS**

### COMMERCIAL

ACTIVE INGREDIENT: Tebuconazole ..... 430 g/L

Contains 1,2-benzisothiazolin-3-one at 0.021% as a preservative.

REGISTRATION NO. 34339      PEST CONTROL PRODUCTS ACT

**READ THE LABEL AND THIS BOOKLET BEFORE USING**

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION: EYE IRRITANT**

**DANGER: SKIN IRRITANT**

NET CONTENTS: 0.03 L to 200 L (BULK)

Albaugh, LLC  
1525 NE 36th Street  
Ankeny, IA 50021  
1-800-247-8013

For medical or treatment information from exposure to this product, call 1-888-347-6732 (7 days/week, 24-hr).

For 24-hour chemical spill, leak, fire, exposure, or accident response information, call CHEMTREC toll free at 1-800-424-9300.

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

## **PRECAUTIONS**

**KEEP OUT OF REACH OF CHILDREN.** Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist.

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and boots during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab or cockpit.

For application using handheld airblast/mistblower equipment, wear chemical-resistant coveralls with a chemical-resistant hood over long-sleeved shirt, long pants, chemical-resistant gloves, socks, chemical-resistant footwear and a respirator with a NIOSH-approved organic-vapour-removing cartridge with a prefilter approved for pesticides OR a NIOSH-approved canister approved for pesticides.

DO NOT handle more than 140 ml per person, per day when using a handheld airblast/mistblower. These restrictions are in place to minimize exposure to individual applicators. Application may need to be performed over multiple days or using multiple applicators.

When mixing, loading and applying using a mechanically-pressurized handgun, wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes, and a respirator with a NIOSH-approved organic-vapour removing cartridge with a prefilter approved for pesticides, or a NIOSH-approved canister approved for pesticides during mixing, loading, application, cleanup and repair. Follow manufacturer's instructions for cleaning/maintaining the Personal Protective Clothing (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

**DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

When tank-mixes are permitted, read and observe all label directions, including rates and restrictions for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.

**User Safety Recommendations:** User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

## **FIRST AID**

**In case of poisoning**, call physician or Poison Control Centre immediately. Have patient lie down and keep quiet.

**IF SWALLOWED:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

### **TOXICOLOGICAL INFORMATION:**

The compound does not cause any definite symptoms that would be diagnostic. Contact with the eyes may cause irritation.

**To Physician:** No specific antidote. Treat symptomatically.

### **ENVIRONMENTAL PRECAUTIONS**

Toxic to birds, small wild mammals, aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

Tebuconazole is persistent and may carryover. It is recommended that this product not be used in areas treated with any products containing tebuconazole during the previous season.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

**STORAGE:** Store this product away from food or feed. Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

### **DISPOSAL:**

**Recyclable Container:** Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

**Returnable Container:** Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

**Refillable Container:** For disposal, this empty container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not use this container for any other purpose.

**Disposal of Unused, Unwanted Product:**

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills.

**LEAK OR SPILL PROCEDURE:** Handle and open container in a manner as to prevent spillage. If container is leaking invert to prevent leakage. If the container is leaking or material is spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away.

**DIRECTIONS FOR USE:**

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

**IMPORTANT:** Read this entire label before using TOLEDO 430 SC Foliar Fungicide. **Ensure no bystanders are present during the application operation.** If you have questions, call the manufacturer at 1-800-247-8013 or obtain technical advice from the distributor or your provincial agricultural representative. Use mechanical flaggers only.

**DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

**Spray Volume:** TOLEDO 430 SC Foliar Fungicide should be applied in a minimum of 100 litres of spray solution per hectare by ground sprayer or 47 litres of spray solution per hectare by aircraft spray equipment. Check equipment calibration frequently.

**Chemigation:** Do not apply this product through any type of irrigation system.

**Mixing:** Add the required amount of TOLEDO 430 SC Foliar Fungicide into the spray tank while filling with water to the desired level. Operate the agitator while mixing. If other materials are added to the spray tank, the TOLEDO 430 SC Foliar Fungicide should be thoroughly dispersed prior to the addition

of other materials. TOLEDO 430 SC Foliar Fungicide is recommended to be used with the registered nonionic surfactant, Agral 90 or Agsurf at 0.125% vol/vol.

## SPRAY DRIFT MANAGEMENT FOR AERIAL AND GROUND APPLICATIONS

For the protection of non-target habitats, overspray or drift to any body of water or other environmentally sensitive habitats must be avoided. Do not apply under conditions where drift to an unprotected person(s), occupied dwelling, or to food, forage, or other plantings can occur.

The interaction of many equipment- and weather-related factors determines the potential for spray drift. In addition to specific restriction related to spray drift, the applicator is responsible for considering all these factors when making application decisions.

**SPRAY HEIGHT:** For aerial applications, spray should be released at the lowest height consistent with efficacy and flight safety. Applications more than 3 metres above the crop canopy should be avoided.

**TEMPERATURE INVERSIONS:** Do not make aerial or ground applications during temperature inversions. Drift potential is high during temperature inversions. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however if fog is not present, inversions can also be identified by the movement of smoke from a ground source. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical mixing.

**HUMIDITY AND TEMPERATURE:** Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift. Avoid spraying during conditions of low humidity and/or high temperature.

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

Airblast application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** direct spray above plants to be treated. Turn off outward pointing nozzles at row ends and outer rows. **DO NOT** apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side.

Aerial application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 km/h at flying height at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotorspan.

**Spray buffer zones:**

A spray buffer zone is NOT required for uses with hand-held application equipment permitted on this label.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop		Spray Buffer Zones (metres) Required for the Protection of:				
			Freshwater Habitat of Depths:		Estuarine/Marine Habitat of Depths:		Terrestrial Habitat:
			Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer	Barley, Wheat (winter, spring, durum, hard red, Canada prairie, soft white), Spring Barley, Oat, Soybean, Asparagus, SRIC poplar and willow		1	0	0	0	1
Airblast	Short rotation intensive culture poplar and willow	Early growth stage	5	1	0	0	2
		Late growth stage	3	1	0	0	1
Aerial	SRIC poplar and willow, Wheat (winter durum and spring), Barley, Oats	Fixed wing	5	0	0	0	15
		Rotary wing	4	0	0	0	15
	Soybean (1 application)	Fixed wing	5	0	0	0	15
		Rotary wing	5	0	0	0	15

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

Cereals (Wheat, Barley and Oats):			
CROP	DISEASES	DOSAGE OF TOLEDO 430 SC FOLIAR FUNGICIDE	REMARKS
Wheat (spring, winter and durum)	For suppression*of <i>Fusarium</i> head blight (scab) ( <i>Gibberella zeae</i> / <i>Fusarium graminearum</i> )  For control of Septoria glume blotch ( <i>Stagonospora nodorum</i> )	292 mL/ha	<p>* TOLEDO 430 SC Foliar Fungicide should only be applied when the risk of <i>Fusarium</i> head blight infection is high. Consult your local extension authority regarding the need for TOLEDO 430 SC Foliar Fungicide use. Head blight is extremely difficult to control. For that reason only suppression of <i>Fusarium</i> head blight (scab) is claimed.</p> <ul style="list-style-type: none"> <li><i>Fusarium</i> head blight (scab) outbreaks occur when the weather is warm and wet at the flowering to soft dough stages.</li> <li>The application of TOLEDO 430 SC Foliar Fungicide for protection against <i>Fusarium</i> head blight (scab) should be considered when these weather conditions are forecasted for this stage of wheat development.</li> <li><b>Timing of application is critical:</b> For optimum suppression of <i>Fusarium</i> Head Blight and control of Septoria glume blotch, apply Toledo 430 SC Foliar Fungicide within the time period from when at least 75% of the wheat heads on the main stem are fully emerged to when 50% of the heads on the main stem are in flower.</li> <li><b>Spray coverage is essential:</b> Ensure thorough coverage of all wheat heads.</li> <li>Spray equipment must be set up to provide good coverage to wheat heads.</li> <li>TOLEDO 430 SC Foliar Fungicide may be applied by ground or air equipment</li> <li><b>GROUND APPLICATION:</b> Apply specified dosage in a minimum of 100 L of water per hectare.</li> <li><b>AERIAL APPLICATION:</b> Apply specified dosage in a minimum of 47 L of water per hectare.</li> <li>TOLEDO 430 SC Foliar Fungicide is recommended to be used with a registered non-ionic surfactant, such as Agral 90 or AgSurf at 0.125% vol/vol.</li> </ul>
	Rusts (Leaf, Stem and Stripe) ( <i>Puccinia triticina</i> , <i>P. graminis</i> , <i>P. striiformis</i> )  Septoria (Leaf Blotch) ( <i>Septoria tritici</i> )  Tan Spot ( <i>Pyrenophora tritici-repentis</i> )	220-292 mL/ha	<ul style="list-style-type: none"> <li>Apply TOLEDO 430 SC Foliar Fungicide to leaf foliage at the first sign or very early stage of disease, especially if weather conditions are conducive to disease development, up to the end of the flowering stage.</li> <li>Where a rate range is specified, use of the highest rate should be considered when weather conditions are conducive to heavy disease development.</li> <li>TOLEDO 430 SC Foliar Fungicide may be applied by ground or air equipment</li> <li><b>GROUND APPLICATION:</b> Apply specified dosage in a minimum of 100 L of water per hectare.</li> <li><b>AERIAL APPLICATION:</b> Apply specified dosage in a minimum of 47 L of water per hectare.</li> <li>TOLEDO 430 SC Foliar Fungicide is recommended to be used with a registered non-ionic surfactant, such as Agral 90 or AgSurf, at 0.125% vol/vol.</li> </ul>
	Powdery Mildew ( <i>Erysiphe graminis</i> )	292 mL/ha	

Barley	Net Blotch ( <i>Pyrenophora teres</i> )  Spot Blotch ( <i>Cochliobolus sativus</i> )  Scald ( <i>Rhynchosporium secalis</i> )  Rusts (Leaf, Stem and Stripe) ( <i>Puccinia hordei</i> , <i>P. graminis</i> , <i>P. striiformis</i> )  Septoria Leaf Blotch ( <i>Septoria passerinii</i> )  Powdery Mildew ( <i>Erysiphe graminis</i> )	220-292 mL/ha	<ul style="list-style-type: none"> <li>• Apply TOLEDO 430 SC Foliar Fungicide at the very early stages of disease development.</li> <li>• Use of the highest rate should be considered when weather conditions are conducive to heavy disease development.</li> <li>• TOLEDO 430 SC Foliar Fungicide may be applied by ground or air equipment</li> <li>• <b>GROUND APPLICATION:</b> Apply specified dosage in a minimum of 100 L of water per hectare.</li> <li>• <b>AERIAL APPLICATION:</b> Apply specified dosage in a minimum of 47 L of water per hectare.</li> <li>• TOLEDO 430 SC Foliar Fungicide is recommended to be used with a registered non-ionic surfactant, such as Agral 90 or AgSurf, at 0.125% vol/vol.</li> </ul>
Oats	Crown Rust ( <i>Puccinia coronata</i> )  Stem Rust ( <i>Puccinia graminis</i> )	220 mL/ha	<ul style="list-style-type: none"> <li>• Apply TOLEDO 430 SC Foliar Fungicide at the very early stages of disease development.</li> <li>• TOLEDO 430 SC Foliar Fungicide may be applied by ground or air equipment</li> <li>• <b>GROUND APPLICATION:</b> Apply specified dosage in a minimum of 100 L of water per hectare.</li> <li>• <b>AERIAL APPLICATION:</b> Apply specified dosage in a minimum of 47 L of water per hectare.</li> <li>• TOLEDO 430 SC Foliar Fungicide is recommended to be used with a registered non-ionic surfactant, such as Agral 90 or AgSurf, at 0.125% vol/vol.</li> </ul>
<p>Restrictions: A maximum of one application of TOLEDO 430 SC Foliar Fungicide may be applied per year to wheat, barley and oats. Applications may not be made within 36 days of harvest. Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with TOLEDO 430 SC Foliar Fungicide. Straw cut after harvest may be fed or used for bedding.</p>			

### Resistance Management Advisory for Cereal Crops:

Repeated application of standalone DMI Fungicides should not be used on the same crop in one season against risky pathogens such as cereal powdery mildew in areas of high disease pressure for that particular pathogen. Mixture products, tank-mixtures or alternation with fungicides having a different mode of action have been shown to protect against the development of resistant forms of disease.

### Tank Mix Combinations with Herbicides:

TOLEDO 430 SC FOLIAR FUNGICIDE can be tank-mixed with the herbicides noted below, for control of early season leaf diseases and weeds on the respective labels. Weeds and crop must be at the correct growth stage as specified in both the TOLEDO 430 SC FOLIAR FUNGICIDE and the tank mix partner label. Always read and follow the label for TOLEDO 430 SC FOLIAR FUNGICIDE and that of the tank mix partner(s) regarding the directions for use, restrictions and precautions. Do not tank mix TOLEDO 430 SC FOLIAR FUNGICIDE with pesticides, fertilizers or any other chemical additives unless recommended on this label.

CROP	DISEASES and WEEDS	TANKMIXES	REMARKS
Wheat	Leaf Diseases on the Toledo 430 SC Foliar Fungicide label and weeds on the respective herbicide labels	Toledo 430 SC Foliar Fungicide at 220 mL/ha + Buctril M at 1L/ha	Apply Toledo 430 SC Foliar Fungicide and herbicide at labeled rates for the respective leaf diseases and weeds in wheat.  Apply once when plants are at the 4-leaf stage up to the early flag leaf stage.
		Toledo 430 SC Foliar Fungicide at 220 mL/ha + Puma <sup>120</sup> Super at 385 or 770 mL/ha + Buctril M at 1L/ha	For use in the prairie provinces and Peace River region of British Columbia only.  Apply Toledo 430 SC Foliar Fungicide and herbicides at labeled rates for the respective leaf diseases and weeds in wheat.
		Toledo 430 SC Foliar Fungicide at 220 mL/ha + Puma <sup>120</sup> Super at 385 or 770 mL/ha + Refine Extra at 20 g/ha	Apply once when plants are at the 4-leaf stage up to the 3 <sup>rd</sup> tiller stage.  DO NOT apply Toledo 430 SC Foliar Fungicide mixed with Puma <sup>120</sup> Super 2-3 days prior to or following cold temperatures (3°C or lower) as crop injury may occur.
Barley	Leaf Diseases on the Toledo 430 SC Foliar Fungicide label and weeds on the respective herbicide labels	Toledo 430 SC Foliar Fungicide at 220 mL/ha + Puma <sup>120</sup> Super at 385 or 770 mL/ha + Refine Extra at 20 g/ha	For use in the prairie provinces and Peace River region of British Columbia only.  Apply Toledo 430 SC Foliar Fungicide and herbicides at labeled rates for the respective leaf diseases and weeds in barley.  Apply once when plants are at the 4-leaf stage to the 5-leaf, 2 <sup>nd</sup> tiller stage.  DO NOT apply Toledo 430 SC Foliar Fungicide mixed with Puma <sup>120</sup> Super 2-3 days prior to or following cold temperatures (3°C or lower) as crop injury may occur.

#### Tank Mix Combinations with Insecticides:

Toledo 430 SC Foliar Fungicide can be tank-mixed with Lorsban 4E Insecticide for control of Orange wheat blossom midge (*Sitodiplosis mosellana*) in wheat at labelled rates. Read carefully and follow use directions and use precautions on both the Toledo 430 SC Foliar Fungicide and Lorsban 4E Insecticide label. When tank mixing Toledo 430 SC Foliar Fungicide and Lorsban 4E Insecticide, always add Toledo 430 SC Foliar Fungicide to the tank first, then add Lorsban 4E Insecticide. Do not add any surfactants, such as Agral 90 or Agsurf.

#### Soybean:

CROP	DISEASES	DOSAGE OF TOLEDO 430 SC FOLIAR FUNGICIDE	REMARKS

Soybean	Asian Soybean Rust ( <i>Phakopsora pachyrhizi</i> )  Frogeye Leaf Spot ( <i>Cercospora sojina</i> )	220-292 mL/ha	<ul style="list-style-type: none"> <li>Apply TOLEDO 430 SC Foliar Fungicide when first symptoms of disease can be found or when the risk of infection is imminent.</li> <li>Use the highest rate when disease pressure is severe. Always use a registered non-ionic surfactant, such as Agral 90 or Agsurf, at 0.125% Vol/Vol.</li> <li>TOLEDO 430 SC Foliar Fungicide may be applied by ground or air equipment</li> <li>GROUND APPLICATION: Apply specified dosage in a minimum of 100 L of water per hectare.</li> <li>AERIAL APPLICATION: Apply specified dosage in a minimum of 47 L of water per hectare.</li> </ul>
---------	---	---------------	--

Restrictions: A maximum of one application of TOLEDO 430 SC Foliar Fungicide may be applied per year. Applications may not be made within 20 days of harvest.

### MINOR USES

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Albaugh, LLC under the User Requested Minor Use Label Expansion program. For these uses, Albaugh, LLC has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

### ASPARAGUS

CROP	DISEASES	DOSAGE OF TOLEDO 430 SC FOLIAR FUNGICIDE	REMARKS
Asparagus	Rust ( <i>Puccinia asparagi</i> )	292 mL/ha	<ul style="list-style-type: none"> <li>Apply at the earliest sign of rust pustules or when weather conditions are conducive for rust development.</li> <li><b>Post-harvest treatment:</b> Apply TOLEDO 430 SC Foliar Fungicide as a foliar spray to the developing ferns after harvest of spears is completed or</li> <li><b>Newly planted, non-harvested asparagus fields:</b> Apply TOLEDO 430 SC Foliar Fungicide as a foliar spray as soon as ferns are 30 cm high.</li> <li>Apply TOLEDO 430 SC Foliar Fungicide in alternation with another effective fungicide with a different mode of action.</li> <li>Do not apply to harvestable spears.</li> <li>Apply specified rate in a minimum of 200L water/ha.</li> <li>For optimum disease control, always use a registered non-ionic surfactant, such as Agral 90 or Agsurf, at 0.125% Vol/Vol.</li> <li>Pre-harvest Interval: 8 months.</li> <li>Apply by ground only.</li> </ul>

### SRIC POPLAR AND WILLOW and their planting stock

#### USE DIRECTIONS

CROP	DISEASES	DOSAGE OF TOLEDO 430 SC FOLIAR FUNGICIDE	REMARKS
Short Rotation Intensive Culture (SRIC) poplar ( <i>Populus</i> spp.) and SRIC willow ( <i>Salix</i> spp.) and their planting stock grown in production facilities (stoolbeds, bareroot beds, seedling beds).	Leaf rust ( <i>Melampsora</i> spp.)	292 mL/ha	<p>A maximum of one application of TOLEDO 430 SC Foliar Fungicide may be applied per year.</p> <p><b>Timing of applications:</b></p> <p><b>First year trees or shoots:</b> apply when trees or shoots are 5-10 cm tall and have at least 3-4 expanded leaves; timing depends on the poplar or willow clone and climatic zone. The first application could be as early as late April to late May (B.C. Coast) and as late as mid June to early July (Prairie Region).</p> <p><b>Second year or older trees or shoots:</b> Two (to three) weeks after new foliage has emerged. The first application could be as early as mid to late April (B.C. coast) and as late as mid to late June (Prairie Region), depending on the poplar or willow clone and climatic zone.</p> <ul style="list-style-type: none"> <li>• <b>GROUND APPLICATION (field sprayer and airblast equipment):</b> Apply specified dosage in a minimum of 100 L of water per hectare. Use higher water volumes to ensure adequate coverage especially for the second application.</li> <li>• <b>AERIAL APPLICATION:</b> Apply specified dosage in a minimum of 47 L of water per hectare.</li> <li>• Check equipment calibration frequently.</li> <li>• TOLEDO 430 SC Foliar Fungicide is recommended to be used with the registered non-ionic surfactant, Agral 90 or AgSurf at 0.125% vol/vol.</li> </ul>

## RESISTANCE-MANAGEMENT RECOMMENDATIONS:

For resistance management, Toledo 430 SC Foliar Fungicide contains a Group 3 fungicide. Any fungal population may contain individuals naturally resistant to Toledo 430 SC Foliar Fungicide and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

### To delay fungicide resistance:

Where possible, rotate the use of Toledo 430 SC Foliar Fungicide or other Group 3 fungicides with different groups that control the same pathogens.

Do not apply more than the indicated maximum number of applications specified for each crop in the DIRECTIONS FOR USE. Avoid using consecutive applications of other fungicides in the same group in a season.

Use tank mixtures with fungicides from a different group that is effective on the target pathogen when such use is permitted.

Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.

Where possible, make use of predictive disease models to effectively time fungicide applications.

Monitor treated fungal populations for resistance development. Notify Albaugh, LLC if reduced sensitivity of the pathogen to Toledo 430 SC Foliar Fungicide is suspected.

If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.

For further information and to report suspected resistance, contact a Albaugh, LLC representative at 1-800-247-8013.

## **AERIAL APPLICATION:**

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices, or equivalent electronic positioning systems (GPS). The use of a spotter plane is recommended. Use mechanical flaggers only.

## **Use Precautions**

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the *National Aerial Pesticide Application Manual*, developed by the Federal/Provincial/Territorial committee on Pest Management and Pesticides.

**Operator Precautions**

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear coveralls, chemical resistant gloves, and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

**Product Specific Precautions**

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-800-247-8013 or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

Volume: Apply the recommended rate in a minimum spray volume of 47 litres per hectare.

**ROTATIONAL CROPS**

Treated areas may be replanted immediately following harvest with any crop listed on this label. For crops not listed on this label, do not plant back within 120 days of last application.