

2024-4416
2025-01-16

This label has been updated according to the re-evaluation decision of Quizalofop-p-ethyl RVD2024-03. While users are encouraged to follow this updated label immediately, the previously approved label is valid until 02-23-2026 in accordance with the phase out period set out in the re-evaluation decision of Quizalofop-p-ethyl RVD2024-03. This previously approved label will be provided upon request by emailing hc.pmra.info-arla.sc@canada.ca. In your email please include the product name and Registration number of the label you are requesting.

2024-4416
2025-01-16

GROUP 1 HERBICIDE

SELECT PLUS HERBICIDE

EMULSIFIABLE CONCENTRATE

**COMMERCIAL (AGRICULTURAL)
FOR POSTEMERGENCE APPLICATION ON REGISTERED CROPS FOR CONTROL
OF GRASSES.**

ACTIVE INGREDIENTS:

Clethodim 240g/L
Quizalofop-p-ethyl 120g/L

WARNING – EYE IRRITANT

REGISTRATION NUMBER 34744 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

Do not freeze.

NET CONTENTS: 250 mL to 1000 L

For Product Use Information Call 1-800-438-6071

**UPL AgroSolutions Canada Inc.
PO Box 12219
Research Triangle Park, NC 27709**

**United States FOR CHEMICAL EMERGENCY: spill, leak, fire, exposure, or accident
call CHEMTREC 1-800-424-9300**

2024-4416
2025-01-16

PRECAUTIONS

- **KEEP OUT OF REACH OF CHILDREN.**
- Causes eye irritation. DO NOT get in eyes
- Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.
- DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.
- When tank mixes are permitted, read and observe all label directions, including rates and restrictions for each product used in the tank mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.

PROTECTIVE CLOTHING

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair. Gloves are not required during application within a closed cab/closed cockpit. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove personal protective equipment immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Remove contaminated clothing immediately if pesticide gets inside.

ENVIRONMENTAL PRECAUTIONS

- TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.
- This product contains active ingredients and aromatic petroleum distillates, which are toxic to aquatic organisms.
- Toxic to certain beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites). Minimize spray drift to reduce harmful effects on beneficial arthropods in habitats next to the application site such as hedgerows and woodland.
- Refer to attached booklet for complete environmental precautions.

FIRST AID

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control centre or doctor for treatment advice.

2024-4416
2025-01-16

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

No specific antidote is available. Treat the patient symptomatically. This product contains petroleum distillate, vomiting may cause aspiration pneumonia.

STORAGE

PESTICIDE STORAGE:

Store this product away from food or feed.

Store in cool, dry place and in such a manner as to prevent cross contamination with other pesticides, and fertilizers. Store in original container and out of reach of children, preferably in a locked storage area. Do not freeze.

SPILL CLEAN-UP: Contact the manufacturer and the provincial regulatory agency in case of a spill and for clean-up of spills. Follow safety precautions as directed for handling the product. If the product is clean, it may be used; otherwise, follow the DISPOSAL instructions.

DISPOSAL

Recyclable Containers

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/ territorial requirements.

Returnable Containers

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

2024-4416
2025-01-16

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/ territorial regulatory agency. Contact the manufacturer and the provincial/ territorial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label.

UPL, the UPL logo SELECT, AMIGO, OHM, WAVE and INTERLINE are trademarks of a UPL Corporation Limited Group Company.

©2024 UPL Corporation Limited Group Company.

2024-4416
2025-01-16

GROUP	1	HERBICIDE
--------------	----------	------------------

SELECT PLUS HERBICIDE

EMULSIFIABLE CONCENTRATE

**COMMERCIAL (AGRICULTURAL)
FOR POSTEMERGENCE APPLICATION ON REGISTERED CROPS FOR CONTROL
OF GRASSES.**

ACTIVE INGREDIENTS:

Clethodim 240g/L
Quizalofop-p-ethyl 120g/L

WARNING – EYE IRRITANT

REGISTRATION NUMBER 34744 PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

Do not freeze.

NET CONTENTS: 250 mL to 1000 L

For Product Use Information Call 1-800-438-6071

**UPL AgroSolutions Canada Inc.
PO Box 12219
Research Triangle Park, NC 27709
United States**

**FOR CHEMICAL EMERGENCY: spill, leak, fire, exposure, or accident call
CHEMTREC 1-800-424-9300**

2024-4416
2025-01-16

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label.

PRECAUTIONS

- **KEEP OUT OF REACH OF CHILDREN.**
- Causes eye irritation. DO NOT get in eyes
- Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.
- DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.
- When tank mixes are permitted, read and observe all label directions, including rates and restrictions for each product used in the tank mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.

PROTECTIVE CLOTHING

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair. Gloves are not required during application within a closed cab/closed cockpit.

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. The pilot is allowed to load premixed chemicals with a closed system. It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Read and follow handling precautions and protective clothing recommendations on broadleaf herbicide and adjuvant tank mix partner labels. Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If no such instructions for washables exist, use detergent and hot water. Keep PPE apart from other laundry and wash separately. Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove personal protective equipment immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Remove contaminated clothing immediately if pesticide gets inside. Wash contaminated clothing separately from household laundry before reuse.

2024-4416
2025-01-16

ENVIRONMENTAL PRECAUTIONS

- TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.
- This product contains active ingredients and aromatic petroleum distillates, which are toxic to aquatic organisms.
- Toxic to certain beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites). Minimize spray drift to reduce harmful effects on beneficial arthropods in habitats next to the application site such as hedgerows and woodland.
- This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.
- To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.
- Avoid application when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

FIRST AID

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control centre or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

No specific antidote is available. Treat the patient symptomatically. This product contains petroleum distillate, vomiting may cause aspiration pneumonia.

STORAGE

2024-4416
2025-01-16

PESTICIDE STORAGE:

Store this product away from food or feed.

Store in cool, dry place and in such a manner as to prevent cross contamination with other pesticides, and fertilizers. Store in original container and out of reach of children, preferably in a locked storage area. Do not freeze.

SPILL CLEAN-UP: Contact the manufacturer and the provincial regulatory agency in case of a spill and for clean-up of spills. Follow safety precautions as directed for handling the product. If the product is clean, it may be used; otherwise, follow the DISPOSAL instructions.

DISPOSAL

Recyclable Containers

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/ territorial requirements.

Returnable Containers

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/ territorial regulatory agency. Contact the manufacturer and the provincial/ territorial regulatory agency in case of a spill, and for clean-up of spills.

PRODUCT INFORMATION

Select Plus is a selective herbicide for post-emergence control of a broad range of annual and perennial grass weeds in *Brassica carinata*, *Brassica juncea* (Oriental / Brown), canola, flax, crop group 6 (legume vegetables), seed alfalfa, soybean, and yellow mustard.

Select Plus applied post-emergence must be tank mixed with AMIGO Adjuvant. For broader spectrum weed control, **Select Plus** may be tank mixed with a broadleaf herbicide listed on this label. See TANK MIXES for recommended products.

Select Plus is a systemic herbicide which is translocated from the treated foliage to the

2024-4416
2025-01-16

growing points of the leaves and shoots. Uptake into the plant is through the foliage, therefore thorough coverage of the foliage is important for consistent and effective grass control. Susceptible weeds cease growth soon after application, removing the competitive effects of susceptible weeds. Treated plants show a reduction in growth with yellowing and browning of the tissue followed by plant death. However, complete weed control may not be seen for one to three weeks depending on plant species, growing conditions and crop competition.

Weed control may be reduced when applications are made to weeds stressed by drought, low fertility, saline soils, water logged soils, and cool weather. These stresses can also reduce crop tolerance.

DIRECTIONS FOR USE

General Precautions

Read the entire DIRECTIONS FOR USE before using **Select Plus**.

- Do not apply post-emergence treatments if it is raining or if rainfall is expected within one (1) hour after application.
- As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.
- Do not contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of waste.
- Do not graze treated fields, cut treated crops for forage or use as a green feed until after the appropriate interval has been observed as outlined in the Crop Recommendations table.
- Do not allow this chemical to drift onto other crops.
- A 30-day plant back interval should be observed for all unlabeled crops.
- Do not freeze.
- DO NOT graze the treated crops or cut for hay.
- Not for use in greenhouses.

Ground Application

- Apply alone or in tank mixes in a spray volume of 50 - 100 L/ha (22.5 - 45 L/acre or 5 - 10 gallons/acre) at 207 - 345 kPa (30 - 50 PSI) pressure to ensure proper weed coverage.
- Flat fan nozzles of 80° or 110° are recommended for optimum coverage. Do not use floodjet nozzles or control droplet application equipment. Nozzles may be oriented 45° forward to enhance crop penetration and to give better weed coverage.
- Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

Aerial Application

- DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotorspan.
- Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.
- Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.
- Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.
- To reduce drift caused by turbulent wingtip vortices, the nozzle distribution along the spray boom length **MUST NOT** exceed 65 % of the wing or rotorspan.

Aerial Use Precautions

- Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the *National Aerial Pesticide Application Manual*, developed by the Federal/Provincial/Territorial committee on Pest management and Pesticides.

Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-866-761-9397 or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

- Apply alone or in tank mixes (ONLY with the recommended tank mix partners that are registered for aerial use) in no less than 28 L/ha spray volume at a pressure that delivers coarse spray droplets as classified by the ASAE.

SPRAY BUFFER ZONES

A spray buffer zone is NOT required for:

- uses with hand-held application equipment permitted on this label,
- low-clearance hooded or shielded sprayers that prevent spray contact with crop, fruit or foliage.

2024-4416
2025-01-16

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands) and sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

Method of Application	Crop	Spray Buffer Zones (metres) Required for the Protection of:			
		Freshwater Habitat of Depths:		Terrestrial Habitat:	
		Less than 1 m	Greater than 1 m		
Field sprayer	<i>Brassica carinata</i> , <i>Brassica juncea</i> , brown mustard, canola, dry pea, flax (including low linolenic acid varieties), Crop subgroups 6A, 6B and 6C (except dry pea and soybean), seed alfalfa, soybean, yellow mustard	1	1	2	
Aerial	<i>Brassica carinata</i> , <i>Brassica juncea</i> , brown mustard, canola, dry pea, flax (including low linolenic acid varieties), Crop subgroup 6C (except dry pea and soybean), soybean, yellow mustard	Fixed wing	3	1	40
		Rotary wing	1	1	40

When tank mixes are permitted, consult the labels of the tank mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

TIMING OF APPLICATION

Apply **Select Plus** when annual and perennial grasses are in the leaf stages specified under RECOMMENDED APPLICATIONS of **Select Plus**. Most effective control is achieved when application is made to grasses actively growing in good growing conditions in the absence of stress, there is a good crop stand, and adequate moisture and fertility.

Select Plus will be less effective when plants are stressed by the lack of moisture, excessive moisture, low-temperature and / or very low relative humidity. Regrowth by tillering may occur if application is made under any of the above stress conditions. In

2024-4416
2025-01-16

wide row crops, where the canopy may be slow to close, cultivation may be necessary to control grasses that emerge after treatment.

Apply **Select Plus** plus **Amigo Adjuvant** once per growing season. Rates of Amigo Adjuvant are summarized within this label.

RECOMMENDED APPLICATIONS

Weeds	Rate	Remarks
Barnyard grass, Green foxtail, Fall panicum, Foxtail barley, Japanese brome, Proso millet, Volunteer barley, Volunteer canary grass, Volunteer corn, Volunteer oat, Volunteer wheat, Wild oat, Witchgrass, Yellow foxtail.	125 mL/ha	2 to 6 leaf stage for grasses (prior to tillering) Apply with Amigo Adjuvant 0.5 % v/v .
Barnyard grass, Downy brome*, Green foxtail, Fall panicum, Foxtail barley*, Japanese brome*, Large crabgrass, Persian darnel, Proso millet, Smooth crabgrass, Volunteer barley, Volunteer canary grass, Volunteer corn, Volunteer oat, Volunteer wheat, Wild oat, Witchgrass, Yellow foxtail.	188 mL/ha	2 to 6 leaf stage for annual grasses. *2 to 6 leaf plus 3 tillers. Apply with Amigo Adjuvant 0.5 % v/v.

2024-4416
2025-01-16

Crop Recommendations and Maximum Rate

Crop	Maximum Rate	Remarks
<i>Brassica carinata</i>	188 mL/ha	Ground and aerial application. Preharvest interval 64 days.
<i>Brassica juncea</i> (Oriental / Brown mustard (condiment and oilseed types))		Ground and aerial application. Preharvest interval 64 days.
Canola		Ground and aerial application. Preharvest interval 64 days.
Dry pea		Ground and aerial application. Preharvest interval 75 days.
Flax (including low linolenic acid varieties)		Ground and aerial application. Preharvest interval 82 days.
Lentil		Ground and aerial application. Preharvest interval 65 days.

2024-4416
2025-01-16

Edamame and Edible-podded legume vegetables (Crop subgroup 6A): Bean (*Phaseolus* spp.) (includes runner bean, snap bean, wax bean), Bean (*Vigna* spp.) (includes asparagus bean, Chinese longbean, moth bean, yardlong bean), Jackbean (*Canavalia ensiformis*), Pea (*Pisum* spp.) (includes dwarf pea, edible-pod pea, snow pea, sugar snap pea), Pigeon pea (*Cajanus cajan*), Soybean (immature seed) (*Glycine max*), Sword bean (*Canavalia gladiata*).

Succulent shelled pea and bean (Crop subgroup 6B): Bean (*Phaseolus* spp.) (includes lima bean), Bean (*Vigna* spp.) (includes blackeyed pea, cowpea, southern pea), Broad bean (fava bean) (*Vicia faba*), Pea (*Pisum* spp.) (includes English pea, garden pea, green pea), Pigeon pea (*Cajanus cajan*).

Dried shelled pea and bean (Crop subgroup 6C) (except dry pea, soybean): Bean (*Lupinus* spp.) (includes grain lupin, sweet lupin, white lupin, white sweet lupin), Bean (*Phaseolus* spp.) (includes field bean, kidney bean, lima bean, navy bean, pinto bean, tepary bean), Bean (*Vigna* spp.) (includes adzuki bean, blackeyed pea, catjang, cowpea, moth bean, mung bean, rice bean, southern pea, urd bean), Broad bean (fava bean) (*Vicia faba*), Chickpea (garbanzo bean) (*Cicer arietinum*), Guar (*Cyamopsis tetragonoloba*), Lablab bean

Subgroup 6A and 6B - ground application only; preharvest interval 65 days.

Subgroup 6C - ground and aerial application; preharvest interval 65 days.

Chickpea – ground and aerial application; preharvest interval 85 days

2024-4416
2025-01-16

(hyacinth bean) (<i>Lablab purpureus</i>), Lentil (<i>Lens esculenta</i>), Pea (<i>Pisum</i> spp.) (includes field pea), Pigeon pea (<i>Cajanus cajan</i>).	
Seed Alfalfa	Ground application
Soybean	Ground and aerial application. Preharvest interval 80 days.
Yellow Mustard	Ground and aerial application. Preharvest interval 64 days.

MIXING INSTRUCTIONS

GENERAL MIXING INSTRUCTIONS

Ensure the spray tank is clean. In-line strainers and nozzle screens should be clean and 50-mesh or coarser.

1. Thoroughly clean the sprayer by flushing the system with water and detergent.
2. Fill the spray tank 1/3 to 1/2 full with clean water and begin agitation or bypass.
3. If the tank mix partner herbicide is a dry or a granule, add the appropriate amount of the tank mix partner. Agitate until thoroughly mixed.
4. Add the appropriate amount of **Select Plus** to the spray tank. Agitate until **Select Plus** is thoroughly mixed. Maintain sufficient agitation during both mixing and application.
5. Add the remaining tank mix partner herbicides. Agitate until thoroughly mixed.
6. Add the adjuvant, agitate until thoroughly mixed.
7. Add Ohm™ or Wave™ then complete filling the tank with balance of water needed.

MIXING INSTRUCTIONS FOR GLUFOSINATE AMMONIUM PRODUCT INTERLINE HERBICIDE or INTERLINE 280 HERBICIDE TANK MIX

1. Thoroughly clean the sprayer by flushing the system with water and detergent.
2. Fill the spray tank 1/3 to 1/2 full with clean water and begin agitation or bypass.
3. Add the appropriate amount of adjuvant directly to the spray tank and continue to agitate until thoroughly mixed.
4. Add the appropriate amount of glufosinate ammonium herbicide to the spray tank and continue to agitate until thoroughly mixed.
5. Add the appropriate amount of **Select Plus**. Agitate until thoroughly mixed.
6. If Facet L is to be added, add the appropriate amount of Facet L. Agitate until thoroughly mixed.
7. Add Ohm or Wave then complete filling tank with balance of water needed.
8. Continue to agitate or run the by-pass system and spray out immediately.
9. After any break in spraying, agitate thoroughly before spraying again. Check inside the tank to ensure agitation is sufficient to remix the spray solution components. Do not allow the mixture to sit overnight.
10. If an oil film starts to build up in the tank, drain and clean tank with a strong

2024-4416
2025-01-16

detergent.

11. Immediately after use, thoroughly clean the sprayer by flushing the system with clean water and detergent.

TANK MIXES

When applied as a tank mix combination, read and observe all label directions, including rates, restrictions, and grazing limitations for each product used in the tank mix. Follow the more stringent label precautionary measures for mixing, loading, and application stated on product labels.

Crop	Tank mix Partner	Rate	Remarks
Glufosinate Ammonium Tolerant / Liberty Link Canola	Interline Herbicide or Interline 280 Herbicide	2.67 to 4 L/ha (150 g ai/L) ¹ or 1.43 to 2.14 L/ha (280 g ai/L) ¹	Glufosinate ammonium tolerant canola only: 2 to 6 leaf stage, prior to bolting. Amigo Adjuvant 0.5 % v/v. See tank mix instructions. Ground application: apply in a minimum of 100 L/ha. When aerial application is permitted: apply in a minimum of 50 L/ha.
	Interline Herbicide + Facet L	3.33 L/ha + 0.28 L/ha	Glufosinate ammonium tolerant canola only: 2 to 6 leaf stage, prior to bolting. Amigo Adjuvant 0.5 % v/v. Ground application: apply in a minimum of 100 L/ha.
Canola	Lontrel 360 Herbicide, Lontrel XC Herbicide, or Lontrel 72 Herbicide	0.42 – 0.83 L/ha 0.25 to 0.5 L/ha, or 210 to 415 g/ha	Canola: 2 to 6 leaf stage, prior to bolting. Amigo Adjuvant 0.5 % v/v. Ground application: apply in a minimum of 100 L/ha.
Flax	A herbicide containing ONLY bromoxynil + MCPA (e.g., Buctril M Herbicide or IPCO Logic M Liquid Herbicide)	280 g ae/ha + 280 g ae/ha (e.g., Buctril M Herbicide at 1 L / ha or IPCO Logic M Liquid Herbicide at 1.25 L/ha	Flax: 5 cm to 10 cm in height. Early crop injury may be observed. Amigo Adjuvant 0.5 % v/v.

2024-4416
2025-01-16

	Curtail M Herbicide	1.5 to 2 L/ha	<p>Flax: 5 cm to 15 cm in height. Early crop injury may be observed.</p> <p>Amigo Adjuvant 0.5 % v/v. Ground application: apply in a minimum of 100 L/ha.</p> <p>Do not cut treated crops for feed or graze until 60 days after application.</p>
	Lontrel 360 Herbicide, Lontrel XC Herbicide, or Lontrel 72 Herbicide	0.56 – 0.83 L/ha 0.34 to 0.5 L/ha, or 280 to 415 g/ha	<p>Flax: 5 cm to 10 cm in height.</p> <p>Amigo Adjuvant 0.5 % v/v. Ground application: apply in a minimum of 100 L/ha.</p>
	MCPA Ester Herbicide (600 Ester)	700 mL/ha	<p>Flax (excluding low linolenic acid varieties): 5 cm to prior to bud stage (pre-bud) while following the preharvest interval.</p> <p>Early crop injury may be observed.</p> <p>Maximum rate of MCPA 600 Ester Herbicide is 700 mL/ha</p> <p>Amigo Adjuvant 0.5 % v/v. Ground application: apply in a minimum of 100 L/ha Aerial application: apply in a minimum of 30 L/ha.</p>
Dry Pea	Odyssey WDG Herbicide or Odyssey NXT II	29 to 43 g/ha	<p>Dry pea: 1 to 6 leaf stage</p> <p>Amigo Adjuvant 0.5 % v/v Ground application: apply in a minimum of 50 L/ha.</p>

¹ Use the appropriate application rate according to the glufosinate-ammonium product guarantee.

This product can be tank mixed with Ohm or Wave. Consult the Ohm and Wave labels and observe all use directions including crops, timings, rates and restrictions.

2024-4416
2025-01-16

SPRAYING INSTRUCTIONS

Ground Application

Unless otherwise specified in the tank mix partner section of this label, apply in a spray volume of 50 - 100 L/ha (22.5 - 45 L/acre or 5 - 10 gallons/acre) at 207 - 345 kPa (30 - 50 PSI) pressure to ensure proper weed coverage.

Flat fan nozzles of 80° or 110° are recommended for optimum coverage. Do not use floodjet nozzle or control droplet application equipment. Nozzles may be oriented 45° forward to enhance crop penetration and to give better weed coverage.

Aerial Application

Unless otherwise specified in the tank mix partner section of this label, apply in a spray volume no less than 28 L/ha (11 L/acre or 3 gallons/acre) at no less than 300 kPa pressure to ensure proper weed coverage.

SPRAYER CLEAN-UP

1. Drain the tank and thoroughly rinse spray tank, boom and hoses with clean water. Pay particular attention to flushing out any visible deposits.
2. Fill the tank with clean water and detergent according to label directions. Flush the hoses, boom and nozzles with the cleaning solution. Circulate for at least 15 minutes. Flush hoses, boom and nozzles once more, then drain the tank.
3. Clean nozzles and screens in a separate container using detergent and water.
4. Repeat #2.
5. Rinse tank, boom and hoses with clean water.
 - Do not clean sprayer near desirable vegetation, wells, or other water sources.
 - Dispose of all rinsings in accordance with provincial regulations.
 - Check tank-mix partner label for any additional clean-up procedures.

ROTATIONAL CROPS

Crops not listed on the label can be planted 30 days after **Select Plus** application.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, **Select Plus** is a Group 1 Herbicide. Any weed population may contain or develop plants naturally resistant to **Select Plus** and other Group 1 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of **Select Plus** or other Group 1 herbicides within a growing season or among growing seasons with different herbicide groups that control the same weeds in a field.

2024-4416
2025-01-16

- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seedling rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other chemical control practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance management and/or integrated weed management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact UPL AgroSolutions Canada Inc. at 1-800-438-6071.

UPL, the UPL logo, SELECT, AMIGO, OHM, WAVE and INTERLINE are trademarks of a UPL Corporation Limited Group Company.

©2024 UPL Corporation Limited Group Company.

BUCTRIL is a registered trademark of Bayer.

LONTREL is a trademark and CURTAIL is a registered trademark of Corteva AgriScience.

PURSUIT, LIBERTY and are registered trademarks of BASF.

FACET is a trademark of BASF SE, used with permission by BASF Canada Inc.

ODYSSEY is a trademark of BASF Agrochemical Products B.V., used with permission by BASF Canada Inc.