

(Container)

GROUP	2	HERBICIDE
-------	---	-----------

Start 25 WDG

**FOR SALE FOR USE IN THE PRAIRIE PROVINCES, PEACE RIVER REGION AND INTERIOR OF
BRITISH COLUMBIA**

Start 25 WDG is a selective herbicide for post emergent control of annual broadleaf weeds including cleavers, wild buckwheat and chickweed in pre-seed application for spring wheat (including durum), spring barley, winter wheat and oats.

HERBICIDE
COMMERCIAL

READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: florasulam 25%
Water dispersible granules

REGISTRATION NO. 34969 PEST CONTROL PRODUCTS ACT

WARNING EYE AND SKIN IRRITANT

NET CONTENTS: 0.02 kg - bulk

SHARDA Cropchem Limited
2nd Floor, Prime Business Park
Dashrathlal Joshi Road
Vile Parle (West)
Mumbai - 400056, India

Canadian Agent:
SHARDA Cropchem Limited
63 Kingsview Blvd
Etobicoke, Ontario, CA
M9R 1V1
1-844-810-5720
1-416-840-5639

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

DO NOT APPLY BY AIR

Avoid contact with eyes, skin and clothing. Causes skin irritation. DO NOT get on skin. Avoid breathing dust or spray mist. Causes eye irritation. DO NOT get in eyes.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and chemical-resistant footwear during mixing, loading, application, clean-up and repair. In addition, wear protective eye wear (goggles or face shield) during mixing, loading, clean-up and repair. Rinse gloves before removal.

At completion of spraying or end of the day: Take a shower immediately. Wash thoroughly with soap and water before eating, drinking or smoking. Remove contaminated clothing at the end of the work session and store and wash separately from household laundry using detergents and hot water before reuse.

Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

When tank-mixes are permitted, read and observe all label directions, including rates and restrictions for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.”

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

No specific antidote. Employ supportive care. Treatment should be based on judgment of the physician in response to reactions of the patient.

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

STORAGE

Store this product away from food or feed.

Store in original containers in a secure, dry storage. Do not allow contamination of seeds, plants, fertilizers or other pesticides. Do not contaminate food, feedstuffs or domestic water supplies. If containers are damaged or spill occurs, use the product immediately or contain the spill with absorbent materials and dispose of waste.

DISPOSAL

Recyclable Containers:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

Returnable Containers:

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable Containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

Disposal of unused, unwanted product:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

(Booklet)

GROUP

2

HERBICIDE

Start 25 WDG

**FOR SALE FOR USE IN THE PRAIRIE PROVINCES, PEACE RIVER REGION AND INTERIOR OF
BRITISH COLUMBIA**

Start 25 WDG is a selective herbicide for post emergent control of annual broadleaf weeds including cleavers, wild buckwheat and chickweed in pre-seed application for spring wheat (including durum), spring barley, winter wheat and oats.

HERBICIDE
COMMERCIAL

READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENT: florasulam 25%
Water dispersible granules

REGISTRATION NO. 34969 PEST CONTROL PRODUCTS ACT

WARNING EYE AND SKIN IRRITANT

NET CONTENTS: 0.02 kg - bulk

SHARDA Cropchem Limited
2nd Floor, Prime Business Park
Dashrathlal Joshi Road
Vile Parle (West)
Mumbai - 400056, India

Canadian Agent:
SHARDA Cropchem Limited
63 Kingsview Blvd
Etobicoke, Ontario, CA
M9R 1V1
1-844-810-5720
1-416-840-5639

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

DO NOT APPLY BY AIR

Avoid contact with eyes, skin and clothing. Causes skin irritation. DO NOT get on skin. Avoid breathing dust or spray mist. Causes eye irritation. DO NOT get in eyes.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and chemical-resistant footwear during mixing, loading, application, clean-up and repair. In addition, wear protective eye wear (goggles or face shield) during mixing, loading, clean-up and repair. Rinse gloves before removal.

At completion of spraying or end of the day: Take a shower immediately. Wash thoroughly with soap and water before eating, drinking or smoking. Remove contaminated clothing at the end of the work session and store and wash separately from household laundry using detergents and hot water before reuse.

Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

When tank-mixes are permitted, read and observe all label directions, including rates and restrictions for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

No specific antidote. Employ supportive care. Treatment should be based on judgment of the physician in response to reactions of the patient.

AGRICULTURAL CHEMICAL

Do not ship or store with food, feeds, drugs or clothing.

ENVIRONMENTAL PRECAUTIONS

TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

STORAGE

Store this product away from food or feed.

Store in original containers in a secure, dry storage. Do not allow contamination of seeds, plants, fertilizers or other pesticides. Do not contaminate food, feedstuffs or domestic water supplies. If containers are damaged or spill occurs, use the product immediately or contain the spill with absorbent materials and dispose of waste.

DISPOSAL

Recyclable Containers:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

Returnable Containers:

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable Containers:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

Disposal of unused, unwanted product:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

GENERAL INFORMATION

Start 25 WDG, alone or in tank-mix with glyphosate herbicides, controls weeds prior to seeding spring wheat (including durum), spring barley, winter wheat and oats. Start 25 WDG can be applied in the fall or spring prior to planting or as an initial treatment in summerfallow.

Start 25 WDG is a dispersible granule that is mixed with water and applied as a uniform broadcast spray. It is non-corrosive, non-flammable, and non-volatile.

Start 25 WDG must be applied early post emergence to the main flush of actively growing broadleaf weeds. Warm, moist growing conditions promote active weed growth and enhance the activity of Start 25 WDG by allowing maximum foliar uptake and contact activity. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and re-growth may occur. For best results, ensure thorough spray coverage of target weeds. See "DIRECTIONS FOR USE" section of this label for complete use details.

Start 25 WDG stops growth of susceptible weeds rapidly. However, typical symptoms (discolouration) of dying weeds may not be noticeable for 1 to 2 weeks after application, depending upon growing conditions and weed susceptibility. Degree of control and duration of effect are dependent on weed sensitivity, weed size, crop competition, growing conditions at and following treatment, and spray coverage.

MODE OF ACTION

Start 25 WDG inhibits the production of the ALS enzyme in plants. This enzyme is essential for the production of certain amino acids which are essential for plant growth. Start 25 WDG is a Group 2 mode of action herbicide.

GENERAL USE PRECAUTIONS

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

DO NOT APPLY BY AIR.

Do not use in successive years at the same site.

This product has potential to leach. Do not apply excessive irrigation.

Do not apply through any type of irrigation system.

Sensitive Plants

Do not apply Start 25 WDG directly to, or otherwise permit it to come in direct contact with susceptible crops or desirable plants including alfalfa, edible beans, canola, flowers and ornamentals, lentils, lettuce, peas, potatoes, radishes, soybeans, sugar beets, sunflowers, tomatoes or tobacco.

Non-Target Sites

Do not apply where proximity of susceptible crops (e.g. canola and legumes) or other desirable plants is likely to result in exposure to spray or spray drift. See Environmental Hazards section of the label.

Crop Rotation

Fields previously treated with Start 25 WDG can be seeded the following year to barley, canola, chickpeas, field beans, flax, Juncea canola, lentils, mustard (brown, oriental and/or yellow), oats, peas, soybeans, sunflower, wheat, or fields can be summerfallowed. Do not use in successive years at the same site.

Preharvest/Grazing Intervals

Livestock may be grazed on treated crops 7 days following application. Do not harvest the treated crop within 60 days after application.

Tank Mixtures

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact SHARDA CropChem Limited at 1-844-810-5720 for information before mixing any pesticide or fertilizer that is not specifically recommended on this label.

To Reduce Spray Drift

1. Use nozzles delivering higher volumes and coarser droplets.
2. Use low pressures (200 to 275 kPa).
3. Use 100 L/ha of spray solution.
4. Spray when the wind velocity is 15 km/hr or less.
5. Spot treatments should only be applied with a calibrated boom to prevent over-application.

Sprayer Clean-Out Instructions

To avoid injury to desirable plants, thoroughly clean equipment used to apply this product before re-use or using it to apply other chemicals.

1. Immediately after spraying, completely drain the sprayer tank. Any contamination on the outside of the spraying equipment should be removed by washing with clean water.
2. First rinse:
 - Spray the inside of tank with clean water and fill the sprayer with at least one tenth of the spray tank volume.
 - Agitate and circulate for 15 minutes, and flush through booms and hoses.
 - Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
 - Drain tank completely.
3. Second rinse:
 - Fill the tank with clean water.
 - Add All Clear Spray Tank Decontaminator plus 1 L of household ammonia (containing a minimum 3% ammonia) per 100L of water as per manufacturer's recommendations while filling the tank with clear water.
 - Agitate and then flush the boom and hoses with the cleaning solution. Top up with water making sure the tank is completely full. Allow to stand for 15 minutes with agitation. Flush the solution out of the spray tank through the spray booms. Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
 - If possible, let the solution stand in the sprayer tank and booms for an extended period of time, overnight if possible.
 - After flushing the boom and hoses, drain tank completely.
 - Remove nozzles and screens and clean separately with a cleaning agent or an ammonia solution (100 mL in 10 L water).
- 4.. Third rinse:
 - Rinse the tank with clean water and flush through the boom and hoses using at least one tenth of the spray tank volume.
 - Remove end caps or open ball valves on the ends of each boom section, and flush solution through the boom ends to ensure there is no spray solution trapped between the boom end and the nozzles.
 - Drain tank completely.

Do not use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty odour which may cause eye, nose, throat and lung irritation. Do not clean equipment in an enclosed area.

SPRAY BUFFER ZONES

Spot treatments using hand-held equipment do not require a spray buffer zone.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Crop	Method of application	Spray Buffer Zones (metres) Required for the Protection of:		
		Freshwater Habitat of Depths:		Terrestrial Habitat:
		Less than 1 m	Greater than 1 m	
All labelled crops	Field sprayer	1	0	3

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

DIRECTIONS FOR USE

READ THE ENTIRE LABEL BEFORE USE. FAILURE TO FOLLOW LABEL INSTRUCTIONS MAY RESULT IN ERRATIC WEED CONTROL OR CROP DAMAGE. DO NOT APPLY TO CROPS UNDERSEEDED WITH LEGUMES.

PRE-SEED AND SUMMERFALLOW USE - START 25 WDG ALONE

Crops Registered

spring wheat (including durum), winter wheat, spring barley and oats

Application Directions

Apply 20 grams of Start 25 WDG per hectare in a minimum of 100 L per hectare of water. Add Agral 90 at 0.2% v/v. See weeds species controlled under "Weeds Controlled or Suppressed by Start 25 WDG Alone at 20 g/ha". Apply prior to seeding spring wheat (including durum), winter wheat, spring barley and oats. Apply when weeds are actively growing. Only weeds emerged at the time of treatment will be controlled. Best results are obtained from applications made to seedling weeds.

Application Timing

Apply to actively growing weeds in the 2-4 leaf stage. Extreme growing conditions such as drought or near freezing temperature prior to, at or following time of application may reduce weed control and increase the risk of crop injury. Only weeds which are emerged at the time of application will be affected. If foliage is wet at the time of application control may be decreased.

Weeds Controlled or Suppressed by Start 25 WDG Alone at 20 g/ha

Weeds Controlled:

- | | |
|--------------------|------------------|
| buckwheat, wild | shepherd's-purse |
| canola, volunteer* | smartweed |
| chickweed, common | stinkweed |
| cleavers | mustard, wild |
| cow cockle | |

Weeds Suppressed:

hempsnettle	sowthistle, annual
narrow-leaved hawks beard	sowthistle, perennial**I
pigweed, redroot	

*Including herbicide-tolerant canola varieties except CLEARFIELD canola.

**Applications made at advanced leaf stages will reduce product effectiveness.

Mixing Instructions

1. Fill sprayer tank 1/2 full of water.
2. Start sprayer tank agitation.
3. Add the required amount of Start 25 WDG.
4. Fill the sprayer tank with sufficient water to spray 100 L of spray mixture per hectare.
5. Add Agral 90 as the last ingredient at the rate of 0.2% v/v (2 L per 1000 L of spray volume).
6. Follow sprayer directions and precautions as outlined above, especially when applying next to sensitive crops (e.g. canola and legumes).
7. Follow sprayer clean-up directions.

TANK-MIX COMBINATIONS OF START 25 WDG + GLYPHOSATE HERBICIDES

Start 25 WDG + glyphosate herbicides (PRESENT AS ISOPROPYLAMINE SALT, DIAMMONIUM SALT, POTASSIUM SALT OR DIMETHYLAMINE SALT) will control annual broadleaved weeds and grasses when applied in the fall or spring prior to planting spring wheat (including durum), winter wheat, barley and oats, or as an initial treatment in summerfallow.

Start 25 WDG + glyphosate must be applied to emerged actively growing weeds. Warm, moist growing conditions promote active weed growth and enhance the activity of Start 25 WDG + glyphosate herbicides by allowing maximum foliar uptake and contact activity. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and regrowth may occur. For best results, ensure thorough spray coverage of target weeds.

Start 25 WDG + glyphosate stops growth of susceptible weeds rapidly. However, typical symptoms (discolouration) of dying weeds may not be noticeable for 1 to 2 weeks after application, depending upon growing conditions and weed susceptibility. Degree of control and duration of effect are dependent on weed sensitivity, weed size, crop competition, growing conditions at and following treatment, and spray coverage.

Delay application until weeds have emerged to the stages described (see „Weeds Controlled or Suppressed by Start 25 WDG + glyphosate,“) to provide adequate leaf surface to receive the spray. Unemerged weeds or vegetation arising from underground rhizomes or root stocks of perennials will not be affected by the spray and will continue to grow. For this reason, best control of most perennial weeds is obtained when treatment is made at late growth stages approaching maturity.

Do not treat weeds under poor growing conditions such a drought stress, disease or insect damage, as reduced weed control may result. Reduced results may also occur when treating weeds heavily covered with dust.

This product does not provide long-term residual weed control. For subsequent residual weed control, follow a label approved herbicide program. Read and carefully observe the cautionary statement and all other information appearing on the labels of all herbicides used.

Heavy rainfall immediately after application may wash the chemical off the foliage and a repeat treatment may be required. Do not apply if rainfall is forecast for the time of application.

Do not mix with any surfactant, pesticide, herbicide oil or any other material other than water unless specified in this booklet. For best results, spray coverage should be uniform and complete. Do not spray weed foliage to the point of runoff.

Mode of Action

This herbicide tank-mix moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days but on most perennial weeds may not occur until 7 to 10 days. Extremely cool or cloudy weather at treatment time may slow down activity of this product and delay visual effects of control. Visible effects are a gradual wilting and yellowing of the plant which advances to complete browning of above ground growth and deterioration of underground plant parts.

ATTENTION: AVOID CONTACT WITH FOLIAGE, GREEN STEMS, OR FRUIT OF CROPS, DESIRABLE PLANTS AND TREES SINCE SEVERE INJURY OR DESTRUCTION MAY RESULT.

APPLY THESE SPRAY SOLUTIONS IN PROPERLY MAINTAINED AND CALIBRATED EQUIPMENT CAPABLE OF DELIVERING DESIRED VOLUMES.

AVOID DRIFT - EXTREME CARE MUST BE USED WHEN APPLYING THIS PRODUCT TO PREVENT INJURING DESIRABLE PLANTS AND CROPS. Do not allow spray mist to drift since even minute quantities of spray can cause severe damage or destruction to nearby crops, plants or other areas on which treatment is not intended, or may cause other unintended consequences. Do not apply when winds are gusty or in excess of 8 km/h or when other conditions, including lesser wind velocities, will allow drift to occur. When spraying, avoid combinations of pressure and nozzle type that will result in fine particles (mist) which are more likely to drift.

DO NOT USE IN GREENHOUSES. REDUCED RESULTS MAY OCCUR IF WATER CONTAINING SOIL IS USED, SUCH AS WATER FROM PONDS AND UNLINED DITCHES.

Application Directions

Start 25 WDG can be tank mixed with glyphosate herbicides (present as isopropylamine salt, diammonium salt, potassium salt or dimethylamine salt) to broaden the spectrum of broadleaf weeds. These tank mixes will provide control of most grass and broadleaf species.

Start 25 WDG + Glyphosate Herbicides (present as isopropylamine salt, diammonium salt, potassium salt or dimethylamine salt)

Apply 450 - 2500 grams a.e. per hectare of glyphosate herbicide (present as isopropylamine salt, diammonium salt, potassium salt or dimethylamine salt) tank mixed with 20 g of Start 25 WDG per hectare.

Always refer to the product label for the tank-mix partner for further information on weeds controlled, directions for use, restrictions and precautionary label statements. When applied as a tank-mix, consult the labels of the tank-mix partners and observe the most restrictive buffer zone of the products involved in the tank mixture.

Weeds Controlled or Suppressed with Start 25 WDG + Glyphosate Herbicides (present as isopropylamine salt, diammonium salt, potassium salt or dimethylamine salt)

Rate of Glyphosate†	Weeds Controlled or Suppressed																																	
<p><u>For an application rate of 450 g ae/ha apply:</u></p> <p>0.90 L/ha (500 g ae/L)</p> <p>0.94 L/ha (480 g ae/L)</p> <p>1.0 L/ha (450 g ae/L)</p> <p>1.25 L/ha (360 g ae/L)</p> <p>Use water volumes of 50 to 100 L/ha</p>	<p>Annual Broadleaved Weeds Controlled or Suppressed:</p> <table border="0"> <tr> <td>buckwheat, wild</td> <td>Narrow-leaved hawk's</td> <td>ragweed, common**</td> </tr> <tr> <td>canola, volunteer*</td> <td>Beard**</td> <td>scentless chamomile</td> </tr> <tr> <td>chickweed, common</td> <td>hempsnettle</td> <td>shepherd's purse</td> </tr> <tr> <td>cleavers</td> <td>kochia</td> <td>smartweed</td> </tr> <tr> <td>cow cockle</td> <td>lady's thumb</td> <td>sowthistle, annual</td> </tr> <tr> <td>flax, volunteer</td> <td>lamb's-quarters</td> <td>(suppression only)</td> </tr> <tr> <td>fleabane, Canada**</td> <td>mustard, wild</td> <td>stinkweed</td> </tr> <tr> <td>flixweed</td> <td>pigweed, redroot</td> <td>thistle, Russian</td> </tr> </table> <p>Annual Grasses Controlled:</p> <table border="0"> <tr> <td>barley, volunteer</td> <td>foxtail, green</td> <td>wheat, volunteer</td> </tr> <tr> <td>brome, downy</td> <td>oats, wild</td> <td></td> </tr> <tr> <td>foxtail, giant</td> <td>Persian darnel</td> <td></td> </tr> </table> <p>Perennial Weeds Controlled: dandelion (seedling, overwintered rosettes, mature plants up to 30 cm. in diameter)</p> <p>Perennial Weeds Suppressed: sow-thistle, perennial***</p>	buckwheat, wild	Narrow-leaved hawk's	ragweed, common**	canola, volunteer*	Beard**	scentless chamomile	chickweed, common	hempsnettle	shepherd's purse	cleavers	kochia	smartweed	cow cockle	lady's thumb	sowthistle, annual	flax, volunteer	lamb's-quarters	(suppression only)	fleabane, Canada**	mustard, wild	stinkweed	flixweed	pigweed, redroot	thistle, Russian	barley, volunteer	foxtail, green	wheat, volunteer	brome, downy	oats, wild		foxtail, giant	Persian darnel	
buckwheat, wild	Narrow-leaved hawk's	ragweed, common**																																
canola, volunteer*	Beard**	scentless chamomile																																
chickweed, common	hempsnettle	shepherd's purse																																
cleavers	kochia	smartweed																																
cow cockle	lady's thumb	sowthistle, annual																																
flax, volunteer	lamb's-quarters	(suppression only)																																
fleabane, Canada**	mustard, wild	stinkweed																																
flixweed	pigweed, redroot	thistle, Russian																																
barley, volunteer	foxtail, green	wheat, volunteer																																
brome, downy	oats, wild																																	
foxtail, giant	Persian darnel																																	
<p><u>For an application rate of 900-2500 g ae/ha apply:</u></p> <p>1.8 - 5.0 L/ha (500 g ae/L)</p> <p>1.9 - 5.2 L/ha (480 g ae/L)</p> <p>2.0 - 5.6 L/ha (450 g ae/L)</p> <p>2.5 - 6.9 L/ha (360 g ae/L)</p> <p>Use water volume of 100 L/ha</p>	<p>Weeds Controlled: Weed claims above plus control of annual sow-thistle</p> <p>Perennial Weeds Controlled: Canada thistle (rosette stage) quack grass</p>																																	

<u>For an application rate of 1700-2500 g ae/ha apply:</u>	Weeds Controlled: Weed claims above plus control of Canada thistle (bud stage or beyond)
3.4 - 5.0 L/ha (500 g ae/L)	
3.6 - 5.2 L/ha (480 g ae/L)	
3.8 - 5.6 L/ha (450 g ae/L)	
4.7 - 6.9 L/ha (360 g ae/L)	
Use water volume of 100 L/ha	

†The product application rate is dependent upon the guarantee of the product. Refer to glyphosate product label for further information on weeds controlled, directions for use, restrictions and precautionary label statements.

* Including all herbicide tolerant canola varieties

** Less than 8 cm in height

*** Applications made at advanced stages will reduce effectiveness

Mixing Instructions

1. Fill sprayer tank 1/2 full of water.
2. Start sprayer tank agitation.
3. Add the required amount of Start 25 WDG, continue agitation.
4. Add the required amount of glyphosate, continue agitation.
5. Fill the sprayer tank with sufficient water to spray 50 - 100 L of spray mixture per hectare.

Application Timing

Apply to actively growing weeds in the 2-4 leaf stage, except where noted above. Extreme growing conditions such as drought or near freezing temperature prior to, at or following time of application may reduce weed control. Only weeds which are emerged at the time of application will be affected. If foliage is wet at the time of application, control may be decreased. Under conditions of high weed density, control may be reduced.

Pre-Seed (spring or fall)

Start 25 WDG + glyphosate may be applied prior to seeding and no longer than 48 hours after seeding prior to any crop emergence. Fields treated with Start 25 WDG + glyphosate may be planted to barley, oats, spring wheat (including durum), winter wheat or summerfallowed.

Chem-Fallow

May 1 to July 31: Start 25 WDG may be applied to summerfallow fields and seeded in the fall to winter wheat and in the following spring to barley, canola, oats, peas or wheat (including durum) or summerfallowed.

Fall Application

Start 25 WDG plus glyphosate may be applied to stubble or summerfallow fields after August 1st and prior to freeze-up and may be seeded in the fall to winter wheat and in the following spring to barley, oats or spring wheat (including durum) or summerfallowed.

Do not use in successive years at the same site.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, Start 25 WDG is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to Start 25 WDG and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of Start 25 WDG or other Group 2 herbicide within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact SHARDA Cropchem Limited at 1-844-810-5720

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

All products are registered trademarks of their respective companies.