

**\*\*Container Label\*\***

GROUP

28

INSECTICIDE



## VIKING™ Rorvik Insecticide

SUSPENSION

COMMERCIAL (AGRICULTURAL)

FOR SALE FOR USE ON ROOT AND TUBER VEGETABLES GROUP, FRUITING VEGETABLES GROUP, BRASSICA VEGETABLES GROUP, LEAFY VEGETABLES GROUP, LEGUME VEGETABLES GROUP, CUCURBIT VEGETABLES GROUP, CORN (FIELD, POP, SWEET, AND SEED), GRASS FORAGE, FODDER, AND HAY GROUP, NON-GRASS ANIMAL FEEDS GROUP, MINT, OKRA, GREEN ONION SUB GROUP, GLOBE ARTICHOKE, HOPS, GREENHOUSE CUCUMBERS, GREENHOUSE TOMATOES, GREENHOUSE EGGPLANT, GREENHOUSE PEPPERS, OILSEEDS GROUP, AND CEREALS

**REGISTRATION NO.** 35421 PEST CONTROL PRODUCTS ACT

**ACTIVE INGREDIENT :** Chlorantraniliprole 200 g/L  
Contains sodium benzoate at 0.2% as a preservative.

**NET CONTENTS:** 0.5 L to 1020 L

**READ THE LABEL AND ATTACHED BOOKLET BEFORE USING**

**In case of emergency call the 24 hour Emergency Response Number 1-800-424-9300  
or 1-703-527-3887**

**Viking Crop Production Partners Inc.**

Box 18033  
RPO Shawnessy  
Calgary, AB. T2Y 0K3  
1-587-743-1029

Viking (TM) is a trademark of Viking Crop Production Partners Inc

**PRECAUTIONS:**

• **KEEP OUT OF REACH OF CHILDREN.**

- Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair.
- Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.
- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing
- DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.
- See the Aerial Application label instructions section of this label for additional precautions specific to aerial application.

**FIRST AID:**

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**IF SWALLOWED:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

For medical emergencies call **CHEMTREC 1-800-424-9300 / +1 703-527-3887**

**TOXICOLOGICAL INFORMATION:** Treat symptomatically.

**ENVIRONMENTAL PRECAUTIONS:**

- Toxic to aquatic organisms. Observe spray buffer zones specified under **DIRECTIONS FOR USE.**
- Toxic to certain beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites). Minimize spray drift to reduce harmful effects on beneficial arthropods in habitats next to the application site such as hedgerows and woodland. For greenhouse uses: may harm certain beneficial arthropods, including those used in greenhouse production. Avoid application when beneficial arthropods are in the treatment area.
- Chlorantraniliprole is persistent and may carry over. It is recommended that this product not be used in areas treated with any products containing chlorantraniliprole during the previous season.

- This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.
- To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.
- Avoid application when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

**STORAGE:**

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed. Store this product away from food or feed.

**DISPOSAL:**

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

**NOTICE TO USER:**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

**\*\*Booklet Label\*\***

**GROUP**

**28**

**INSECTICIDE**



**VIKING™ Rorvik Insecticide**

SUSPENSION

COMMERCIAL (AGRICULTURAL)

FOR SALE FOR USE ON ROOT AND TUBER VEGETABLES GROUP, FRUITING VEGETABLES GROUP, BRASSICA VEGETABLES GROUP, LEAFY VEGETABLES GROUP, LEGUME VEGETABLES GROUP, CUCURBIT VEGETABLES GROUP, CORN (FIELD, POP, SWEET, AND SEED), GRASS FORAGE, FODDER, AND HAY GROUP, NON-GRASS ANIMAL FEEDS GROUP, MINT, OKRA, GREEN ONION SUB GROUP, GLOBE ARTICHOKE, HOPS, GREENHOUSE CUCUMBERS, GREENHOUSE TOMATOES, GREENHOUSE EGGPLANT, GREENHOUSE PEPPERS, OILSEEDS GROUP, AND CEREALS

**ACTIVE INGREDIENT:** Chlorantraniliprole 200 g/L  
Contains sodium benzoate at 0.2% as a preservative.

**REGISTRATION NO.** 35421 PEST CONTROL PRODUCTS ACT

**NET CONTENTS:** 0.5 L to 1020 L

**READ THE LABEL AND BOOKLET BEFORE USING**

**In case of emergency call the 24 hour Emergency Response Number 1-800-424-9300 or 1-703-527-3887**

**Viking Crop Production Partners Inc.**

Box 18033  
RPO Shawnessy  
Calgary, AB. T2Y 0K3  
1-587-743-1029

Viking (TM) is a trademark of Viking Crop Production Partners Inc

**PRECAUTIONS:****• KEEP OUT OF REACH OF CHILDREN.**

- Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables are available, use detergent and hot water. Keep and wash PPE separately from other laundry.
- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.
- See the Aerial Application label instructions section of this label for additional precautions specific to aerial application.

**FIRST AID:**

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**IF SWALLOWED:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

For medical emergencies call **CHEMTREC 1-800-424-9300 / +1 703-527-3887**

**TOXICOLOGICAL INFORMATION:** Treat symptomatically.

**ENVIRONMENTAL PRECAUTIONS:**

- Toxic to aquatic organisms. Observe spray buffer zones specified under **DIRECTIONS FOR USE.**
- Toxic to certain beneficial arthropods (which may include predatory and parasitic insects, spiders, and mites). Minimize spray drift to reduce harmful effects on beneficial arthropods in habitats next to the application site such as hedgerows and woodland. For greenhouse uses: may harm certain beneficial arthropods, including those in greenhouse production. Avoid application when beneficial arthropods are in the treatment area.
- Chlorantraniliprole is persistent and may carry over. It is recommended that this product not be used in areas treated with any products containing chlorantraniliprole during the previous season.
- This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

- To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.
- Avoid application when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

**STORAGE:**

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed. Store this product away from food or feed.

**DISPOSAL:**

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

**GENERAL INFORMATION**

**VIKING Rorvik Insecticide** is a liquid formulation that can be applied as a foliar spray, using ground application or aerial application (potatoes, sugar beets, garden beets, legume vegetables, corn, oilseeds and cereals only) to control many important insect pests. **VIKING Rorvik Insecticide** is mixed with water for application.

Apply at the recommended rates when insect populations reach locally determined economic thresholds. Consult your local extension specialist or certified crop advisor to determine appropriate threshold levels for treatment in your area.

Use sufficient water to obtain thorough, uniform coverage.

**DIRECTIONS FOR USE**

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

**DO NOT** contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

**DO NOT** allow releases, effluent or runoff from greenhouses containing this product to enter lakes, streams, ponds or other waters.

**ROOT AND TUBER VEGETABLES – Crop Group 1**

**Spray Volume for Root and Tuber Vegetables:** Apply in a minimum finished spray volume of 100 L/ha by ground. For potatoes only, apply in a minimum finished spray volume of 50 L/ha by air.

SITE	PEST	RATE VIKING Rorvik Insecticide /HA	SPECIFIC DIRECTIONS
POTATOES	Colorado potato beetle	250 mL to 500 mL	For potatoes, apply by either ground or aerial application equipment.
	European corn borer	250 mL to 375 mL	
ROOT AND TUBER VEGETABLES  Arracacha; Arrowroot; Artichoke, Chinese; Artichoke, Jerusalem; Beet, garden; Beet, sugar; Burdock, edible; Canna, edible; Carrot;	Diamondback moth	250 mL	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.
Cabbage looper	For Colorado potato beetle, use the higher application rate under heavy pest pressure and/or when larger larvae are present.		
Black cutworm	Colorado potato beetle resistance management: DO NOT apply <b>VIKING Rorvik Insecticide</b> for Colorado potato beetle control if any Group 28 Insecticide was used at planting as an in-furrow, soil or seed-piece treatment.		
Imported cabbageworm			
Swede midge			

Cassava, bitter and sweet; Celeriac (celery root); Chayote (root); Chervil, turnip-rooted; Chicory; Chufa; Dasheen (taro); Ginger; Ginseng; Horseradish; Leren; Parsley, turnip-rooted; Parsnip; Potato;	Corn earworm  European corn borer  Tobacco hornworm  Tomato hornworm  Armyworm  Variegated cutworm	250 mL to 375 mL	For control of the European corn borer, time the application to coincide with peak egg hatch. Scout for European corn borer by monitoring egg laying and egg hatch to determine application timing.
Radish; Radish, oriental (daikon); Rutabaga; Salsify; Salsify, black; Salsify, Spanish; Skirret; Sweet potato; Tanier (cocoyam); Turmeric; Turnip; Yam bean; Yam, true	Fall armyworm  Beet armyworm  Leafminers: <i>Liriomyza sativae</i> <i>Liriomyza trifolii</i>		

**RESTRICTIONS AND PRECAUTIONS: Root and Tuber Vegetables**

- For pests with a range of application rates, use the high rate under heavy pest pressure.
- Do not make more than 4 applications per season.
- Do not apply more than once every 3 days.
- Do not apply less than 1 day before harvest. Observe a 1-day PHI.
- Do not exceed a total of 1.125 litres **VIKING Rorvik Insecticide** per ha per season.
- **Do not make a foliar application of VIKING Rorvik Insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

## LEAFY VEGETABLES – Crop Group 4

**Spray Volume for Leafy Vegetables:** Apply in a minimum finished spray volume of 100 L/ha by ground.

SITE	PEST	RATE VIKING Rorvik Insecticide PER HECTARE	SPECIFIC DIRECTIONS
LEAFY VEGETABLES Amaranth leafy, Arugula, Cardoon, Celery, Celery (Chinese), Celtuce, Chevрил, Chrysanthemum (edible leaved), Chrysanthemum (garland), Corn Salad, Cress (garland), Cress (upland), Dandelion leaves, Dock, Endive, Florence Fennel, Lettuce (head & leaf), Orach, Parsley leaves, Purslane (garden), Purslane (winter), Radicchio, Rhubarb, Spinach, Spinach (vine), Spinach (New Zealand), Swiss Chard	Cabbage looper  Black cutworm	250 mL	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.
	Armyworm  Fall armyworm  Corn earworm  Beet armyworm  Leafminers: <i>Liriomyza sativae</i> <i>Liriomyza trifolii</i>	250 mL to 375 mL	

### RESTRICTIONS AND PRECAUTIONS: Leafy Vegetables

- Do not make more than 4 applications per season.
- Do not apply more than once every 3 days.
- Do not apply less than 1 day before harvest. Observe a 1-day PHI.
- Do not exceed a total of 1 litre of **VIKING Rorvik Insecticide** per ha per season.
- **Do not make a foliar application of VIKING Rorvik Insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

**BRASSICA VEGETABLES - Crop Group 5**

**Spray Volume for Brassica Vegetables:** Apply in a minimum finished spray volume of 100 L/ha by ground.

SITE	PEST	RATE VIKING Rorvik Insecticide PER HECTARE	SPECIFIC DIRECTIONS
BRASSICA VEGETABLES Broccoli, Broccoli (Chinese), Broccoli raab, Brussels sprouts, Cabbage, Cabbage (Chinese, bok choy), Cabbage (Chinese, napa), Cabbage (Chinese mustard, choy), Cauliflower, Cavalo broccolo, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens	Imported cabbage worm	250 mL	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.  For optimum control, apply MSO® Concentrate with LeciTech® at an application rate of 0.5% v/v.
	Diamondback moth  Cabbage looper  Black cutworm  Armyworm  Fall armyworm  Beet armyworm  Corn earworm  Leafminers: <i>Liriomyza sativae</i> <i>Liriomyza trifolii</i>	250 mL to 375 mL	

**RESTRICTIONS AND PRECAUTIONS: Brassica Vegetables**

- Do not make more than 4 applications per season.
- Do not apply more than once every 3 days.
- Do not apply less than 3 days before harvest. Observe a 3-day PHI.
- Do not exceed a total of 1 litre of **VIKING Rorvik Insecticide** per ha per season.
- **Do not make a foliar application of VIKING Rorvik Insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

**LEGUME VEGETABLES - Crop Group 6**

**Spray Volume for Legume Vegetables:** Apply in a minimum finished spray volume of 100 L/ha by ground. Apply in a minimum finished spray volume of 50 L/ha by air.

SITE	PEST	RATE VIKING Rorvik Insecticide PER HECTARE	SPECIFIC DIRECTIONS
LEGUME VEGETABLES	Grasshoppers	125 mL to 250 mL	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.
<p>Bean (<i>Lupinus</i>) (includes grain lupin, sweet lupin, white lupin, and white sweet lupin);</p> <p>Bean (<i>Phaseolus</i>) (includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean);</p> <p>Bean (<i>Vigna</i>) (includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpea, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean);</p>	<p>Cabbage looper</p> <p>Cutworms</p>	250 mL	

<p>Broad bean (fava);</p> <p>Chickpea (garbanzo);</p> <p>Guar;</p> <p>Jackbean;</p> <p>Lablab bean;</p> <p>Lentil;</p> <p>Pea (<i>Pisum</i>) (includes dwarf pea, edible podded pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea);</p> <p>Pigeon pea;</p> <p>Sword bean;</p> <p>Soybean;</p>	<p>Armyworm</p> <p>Fall armyworm</p> <p>Beet armyworm</p> <p>Corn earworm</p> <p>European corn borer</p> <p>Western bean cutworm</p>	<p>250 mL to 375 mL</p>	
--	--	-------------------------	--

**RESTRICTIONS AND PRECAUTIONS: Legume Vegetables**

- Do not make more than 4 applications per season.
- Do not apply more than once every 3 days.
- Do not apply less than 1 day before harvest. Observe a 1-day PHI.
- Do not exceed a total of 1.125 litres of **VIKING Rorvik Insecticide** per ha per season.
- **Do not make a foliar application of VIKING Rorvik Insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

**FRUITING VEGETABLES (EXCEPT CUCURBITS) – Crop Group 8**

**Spray Volume for Fruiting Vegetables:** Apply in a minimum finished spray volume of 100 L/ha by ground.

SITE	PEST	RATE VIKING Rorvik Insecticide PER HECTARE	SPECIFIC DIRECTIONS
FRUITING VEGETABLES Eggplant, Groundcherry, Pepino, Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper) Tomatillo, Tomato	Colorado potato beetle	250 mL to 375 mL	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.  Colorado potato beetle resistance management: DO NOT apply <b>VIKING Rorvik Insecticide</b> for Colorado potato beetle control if any Group 28 Insecticide was used at planting as an in-furrow, soil or seed treatment.
	Armyworm		
	Fall armyworm		
	Beet armyworm		
	Variegated cutworm		
	Tobacco hornworm		
	Tomato hornworm		
	Tomato fruitworm (Corn earworm)		
Cabbage looper	250 mL		
			Black cutworm

**RESTRICTIONS AND PRECAUTIONS: Fruiting Vegetables**

- For pests with a range of application rates, use the high rate under heavy pest pressure.

- Do not make more than 4 applications per season.
- Do not apply more than once every 5 days.
- Do not apply less than 1 day before harvest. Observe a 1-day PHI.
- Do not exceed a total of 1.125 litres **VIKING Rorvik Insecticide** per ha per season.
- **Do not make a foliar application of VIKING Rorvik Insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

#### CUCURBIT VEGETABLES – Crop Group 9

**Spray Volume for Cucurbit Vegetables:** Apply in a minimum finished spray volume of 100 L/ha by ground.

SITE	PEST	RATE VIKING Rorvik Insecticide PER HECTARE	SPECIFIC DIRECTIONS
CUCURBIT VEGETABLES Chayote (fruit). Chinese waxgourd (Chinese preserving melon); Citron melon; Cucumber; Gherkin; Gourd, edible (includes hyotan, cucuzza, hechima, Chinese okra); Momordica spp (includes balsam apple, balsam pear, bittermelon, Chinese cucumber); Muskmelon (includes cantaloupe); Pumpkin; Squash, summer; Squash, winter (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); Watermelon	Cabbage looper  Black cutworm	250 mL	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.
	Armyworm  Fall armyworm  Corn earworm  Leafminers: <i>Liriomyza sativae</i> <i>Liriomyza trifolii</i>	250 mL to 375 mL	

#### RESTRICTIONS AND PRECAUTIONS: Cucurbit Vegetables

- For pests with a range of application rates, use the high rate under heavy pest pressure.
- Do not make more than 4 applications per season.
- Do not apply more than once every 5 days.
- Do not apply less than 1 day before harvest. Observe a 1-day PHI.
- Do not exceed a total of 1.125 litres **VIKING Rorvik Insecticide** per ha per season.
- **Do not make a foliar application of VIKING Rorvik Insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

**CORN – FIELD, POP, SEED AND SWEET**

**Spray Volume for Corn:** Apply in a minimum finished spray volume of 100 L/ha by ground. Apply in a minimum finished spray volume of 50 L/ha by air.

SITE	PEST	RATE VIKING Rorvik Insecticide PER HECTARE	SPECIFIC DIRECTIONS
CORN  Field corn, Popcorn, Seed corn, Sweet corn	Black cutworm	250 mL  Apply either by ground or aerial application equipment	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.
	Armyworm  Fall armyworm  Beet armyworm  Variegated cutworm  Corn earworm/ Tomato fruitworm  European corn borer  Western bean cutworm	250 mL to 375 mL  Apply either by ground or aerial application equipment	For corn earworm, European corn borer or western bean cutworm control, time the applications to coincide with peak egg hatch. Scout by monitoring egg laying and egg hatch to determine application timing. Reapply if monitoring indicates it is necessary.  Refer to the section of the label titled AERIAL APPLICATION LABEL INSTRUCTIONS for use information specific to aerial application.

**RESTRICTIONS AND PRECAUTIONS: Corn**

- For pests with a range of application rates, use the high rate under heavy pest pressure.
- Do not make more than 4 applications per season.
- Do not apply more than once every 3 days on seed corn or sweet corn.
- Do not apply more than once every 7 days on field corn or popcorn.
- Do not apply less than 1 day before harvest of seed corn or sweet corn. Observe a 1-day PHI.
- Do not apply less than 14 days before harvest of field corn or popcorn. Observe a 14-day PHI.
- Do not exceed a total of 1.125 litres **VIKING Rorvik Insecticide** per ha per season.
- **Do not make a foliar application of VIKING Rorvik Insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

**CEREAL GRAINS – Crop Group 15 (EXCEPT CORN AND WILD RICE) and FORAGE, FODDER, and STRAW OF CEREALS– Crop Group 16 (EXCEPT CORN AND WILD RICE)**

**Spray Volume for Cereal Grains and Forage, Fodder and Straw of Cereals:** Apply in a minimum finished spray volume of 100 L/ha by ground. Apply in a minimum finished spray volume of 50 L/ha by air.

SITE	PEST	RATE VIKING Rorvik Insecticide PER HECTARE	SPECIFIC DIRECTIONS
CEREAL GRAINS  Barley, Buckwheat, Millet, pearl; Millet, proso; Oats, Rye, Sorghum, Teosinte, Triticale, Wheat	Grasshoppers	125 mL to 250 mL	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.
	Cutworms	250 mL	
	Armyworm	250 mL to 375 mL	
	Fall armyworm		
	Beet armyworm		
Corn earworm			
European corn borer			

**RESTRICTIONS AND PRECAUTIONS: Cereal Grains and Forage, Fodder and Straw of Cereals –**

- Forage, fodder and hay may be fed to livestock.
- Do not make more than 3 applications per season.
- Do not apply more than once every 7 days.
- Do not apply less than 1 day before harvest. Observe a 1-day PHI.
- Do not exceed a total of 1.125 litres **VIKING Rorvik Insecticide** per ha per season.
- For pests with a range of application rates, use the high rate under heavy pest pressure.
- **Do not make a foliar application of VIKING Rorvik Insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

**GRASS FORAGE, FODDER, AND HAY GROUP – Crop Group 17**

**Spray Volume for Grass Forage, Fodder, and Hay Group:** Apply in a minimum finished spray volume of 100 L/ha by ground.

SITE	PEST	RATE VIKING Rorvik Insecticide PER HECTARE	SPECIFIC DIRECTIONS
GRASS FORAGE FODDER AND HAY GROUP	Grasshoppers	125 mL to 250 mL	Begin applications when treatment thresholds have been

Any grass, Gramineae family (either green or cured) except sugarcane and those included in the cereal grains group, that will be fed to or grazed by livestock, all pasture and range grasses and grasses grown for hay or silage	Armyworm	250 mL to 375 mL	reached. Thorough coverage is important to obtain optimum control.
	Fall armyworm		
	Beet armyworm		
	Corn earworm		

**RESTRICTIONS AND PRECAUTIONS: Grass Forage, Fodder, and Hay Group**

- For any of the pests listed above, use the high rate under heavy pest pressure.
- Do not make more than 4 applications per season.
- Do not apply more than once every 7 days.
- Observe a 0-day PHI.
- Do not exceed a total of 1.125 litres **VIKING Rorvik Insecticide** per ha per season.
- **Do not make a foliar application of VIKING Rorvik Insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

**NON-GRASS ANIMAL FEEDS GROUP – Crop Group 18**

**Spray Volume for Non-Grass Animal Feeds Group:** Apply in a minimum finished spray volume of 100 L/ha by ground.

SITE	PEST	RATE VIKING Rorvik Insecticide PER HECTARE	SPECIFIC DIRECTIONS
NON-GRASS ANIMAL FEEDS GROUP Alfalfa; Bean, velvet; Clover ( <i>Trifolium</i> , <i>Melilotus</i> ); Kudzu; Lespedeza; Lupin; Sainfoin; Trefoil; Vetch; Vetch, crown; Vetch, milk	Grasshoppers	125 mL to 250 mL	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.
	Beet armyworm	250 mL to 375 mL	
	Alfalfa weevil (suppression only)	375 mL to 500 mL	

**RESTRICTIONS AND PRECAUTIONS: Non-Grass Animal Feeds Group**

- For either of the pests listed above, use the high rate under heavy pest pressure.
- Do not make more than 1 application per cutting.
- Observe a 0-day PHI.

- Do not make a foliar application of VIKING Rorvik Insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.

#### OILSEEDS GROUP – Crop Group 20

**Spray volume for Oilseeds Group:** Apply a minimum finished spray volume of 100 L/ha by ground. Apply a minimum finished spray volume of 50 L/ha by air.

SITE	PEST	RATE VIKING Rorvik Insecticide PER HECTARE	SPECIFIC DIRECTIONS
OILSEEDS GROUP  Canola Rape seed; Mustard seed; Flax; Linseed; Sunflower seed; Safflower; Borage; Calendula; Crambe; Cuphea; Echium; Euphorbia; Evening Primrose; Gold of Pleasure; Hare's ear mustard; Lesquerella; Lunaria; Meadowfoam; Milkweed; Niger seed; Oil radish; Poppy seed; Rose hip; Stokes aster; Sweet rocket; Tea oil plant;	Diamondback moth	125 mL	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.
	Grasshoppers	125 mL to 250 mL	
	Bertha armyworm	125 mL to 375 mL	
	Cabbage looper	250 mL	
	Imported cabbage worm		
	Swede midge		
	Cutworms		
	Sunflower head moth	250 mL to 375 mL	
	Reduces damage caused by banded sunflower moth.		

#### RESTRICTIONS AND PRECAUTIONS: Oilseeds

- Do not make more than 3 applications per season.
- Do not apply more than once every 5 days.
- Do not apply less than 1 day before harvest. Observe a 1-day PHI
- Do not exceed a total of 1.125 litres VIKING Rorvik Insecticide per ha per season.
- Do not make a foliar application of VIKING Rorvik Insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.

## MINT

**Spray Volume for Mint:** Apply in a minimum finished spray volume of 100 L/ha by ground.

SITE	PEST	RATE VIKING Rorvik Insecticide PER HECTARE	SPECIFIC DIRECTIONS
Mint	Cabbage looper	250 mL	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.

### RESTRICTIONS AND PRECAUTIONS: Mint

- Do not make more than 4 applications per season.
- Do not apply more than once every 14 days.
- Do not apply less than 3 days before harvest. Observe a 3-day PHI.
- Do not exceed a total of 1 litre **VIKING Rorvik Insecticide** per ha per season.
- **Do not make a foliar application of VIKING Rorvik Insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

## OKRA

**Spray Volume for Okra:** Apply in a minimum finished spray volume of 100 L/ha by ground.

SITE	PEST	RATE VIKING Rorvik Insecticide PER HECTARE	SPECIFIC DIRECTIONS
Okra	Beet armyworm  Corn earworm	250 mL to 375 mL	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.

### RESTRICTIONS AND PRECAUTIONS: Okra

- For either of the pests listed above, use the high rate under heavy pest pressure.
- Do not make more than 4 applications per season.
- Do not apply more than once every 5 days.
- Do not apply less than 1 day before harvest. Observe a 1-day PHI.
- Do not exceed a total of 1.125 litres **VIKING Rorvik Insecticide** per ha per season.
- **Do not make a foliar application of VIKING Rorvik Insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

## HOPS

**Spray Volume for Hops:** Apply in a minimum finished spray volume of 100 L/ha by ground.

SITE	PEST	RATE VIKING Rorvik Insecticide PER HECTARE	SPECIFIC DIRECTIONS
Hops	Cutworms  Armyworm	250 to 375 mL	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.
	Cabbage looper	250 mL	

**RESTRICTIONS AND PRECAUTIONS: Hops**

- Do not make more than 4 applications per season.
- Do not apply more than once every 7 days.
- Do not apply less than 0 day before harvest. Observe a 0-day PHI.
- Do not exceed a total of 1.125 litres of **VIKING Rorvik Insecticide** per ha per season.
- Do not apply by air.

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Viking Crop Protection Partners Inc. under the User Requested Minor Use Label Expansion program. For these uses, Viking Crop Protection Partners Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

**FOR USE ON BRASSICA VEGETABLES - Crop Group 5**

**Spray Volume for Brassica Vegetables:** Apply in a minimum finished spray volume of 100 L/ha by ground.

SITE	PEST	RATE VIKING Rorvik Insecticide PER HECTARE	SPECIFIC DIRECTIONS
BRASSICA VEGETABLES Broccoli, Broccoli (Chinese), Broccoli raab, Brussels sprouts, Cabbage, Cabbage (Chinese, bok choy), Cabbage (Chinese, napa), Cabbage (Chinese mustard, choy), Cauliflower, Cavalo broccolo, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard spinach, Rape greens	Swede midge	250 mL  For optimum control, MSO Concentrate with LeciTech at an application rate of 0.5% v/v.	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.  Monitor Swede midge activity every 2-4 days using Jackson traps baited with pheromone lures.  Follow local threshold. Apply <b>VIKING Rorvik Insecticide</b> as soon as possible once the Swede midge threshold is reached. Applications on cabbage, Chinese cabbage (napa), cauliflower and Chinese broccoli (gai lon) must be made early in the growing season, before heads start to form.  Given that there are multiple generations of Swede midge, several applications of insecticide (to a maximum of 4 per season) will likely be required.

**RESTRICTIONS AND PRECAUTIONS: Brassica Vegetables**

- Do not make more than 4 applications per season.
- Do not apply more than once every 3 days. For swede midge, make additional applications when monitoring indicates they are necessary, according to local thresholds.
- Do not apply less than 3 days before harvest. Observe a 3-day PHI.
- Do not exceed a total of 1 litre of **VIKING Rorvik Insecticide** per ha per season.
- Do not apply by air.
- **Do not make a foliar application of VIKING Rorvik Insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

**FOR USE ON Globe Artichokes**

**Spray Volume for Globe Artichoke:** Apply in a minimum finished spray volume of 100 L/ha by ground.

SITE	PEST	RATE VIKING Rorvik Insecticide PER HECTARE	SPECIFIC DIRECTIONS
Globe Artichoke	Cutworms	250 mL to 375 mL	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.
	Armyworm		
	Cabbage looper	250 mL	

**RESTRICTIONS AND PRECAUTIONS: Artichokes**

- Do not make more than 4 applications per season.
- Do not apply more than once every 7 days.
- Do not apply less than 3 days before harvest. Observe a 3-day PHI.
- Do not exceed a total of 1.125 litres of **VIKING Rorvik Insecticide** per ha per season.
- Do not apply by air.
- **Do not make a foliar application of VIKING Rorvik Insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

**FOR USE ON GREENHOUSE CUCUMBERS, GREENHOUSE PEPPERS, GREENHOUSE EGGPLANTS, AND GREENHOUSE TOMATOES**

**Spray Volume for Greenhouse Vegetables:** For greenhouse tomatoes, the maximum finished spray volume is 1400 L/ha. For greenhouse cucumbers, greenhouse eggplants, and greenhouse peppers, the maximum finished spray volume is 2000 L/ha.

SITE	PEST	RATE VIKING Rorvik Insecticide /1000 L finished spray volume	SPECIFIC DIRECTIONS
Greenhouse cucumbers,  Greenhouse peppers,  Greenhouse eggplant,  Greenhouse tomatoes	Cabbage loopers	125 mL	Begin applications when treatment thresholds have been reached. Thorough coverage is required to obtain optimum control. Repeat applications if monitoring indicates it is necessary.

**RESTRICTIONS AND PRECAUTIONS: Greenhouse Vegetables**

- Do not make more than 3 applications per crop cycle.
- Do not apply more than once every 7 days.
- Do not apply less than 1 day before harvest. Observe a 1-day PHI.
- Do not exceed a total of 750 mL of **VIKING Rorvik Insecticide** per ha per crop cycle.
- DO NOT allow releases, effluent or runoff from greenhouses containing this product to enter lakes, streams, ponds or other waters.
- **Do not make a foliar application of VIKING Rorvik Insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

PMRA Approved Label – Jeff Ballentine  
 Sub. No. 2024-6783, 2025-01-XX  
 Compared to precedent Sub. No. 2024-5808, 2024-12-12

**FOR USE ON GREENHOUSE PEPPERS, GREENHOUSE EGGPLANTS, AND GREENHOUSE TOMATOES**

**Spray Volume for Greenhouse Vegetables:** Apply in a maximum finished spray volume of 1250 L/ha. Please note that the use of this product for control of tomato leafminer is restricted to plant growth stages for which thorough coverage can be achieved by application volumes of 1250 L/ha or less.

	SITE	PEST	RATE VIKING Rorvik Insecticide /1000 L finished spray volume	SPECIFIC DIRECTIONS
	Greenhouse peppers, Greenhouse eggplant, Greenhouse tomatoes	Lepidopteran leafminer	200 mL	Apply at egg hatch. Reapply if monitoring indicates it is necessary. Thorough coverage is important to obtain optimum control.

**RESTRICTIONS AND PRECAUTIONS: Greenhouse Vegetables**

- Do not make more than 3 applications per crop cycle.
- Do not apply more than once every 7 days.
- Do not apply less than 1 day before harvest. Observe a 1-day PHI.
- Do not exceed a total of 750 mL of **VIKING Rorvik Insecticide** per ha per crop cycle.
- DO NOT allow releases, effluent or runoff from greenhouses containing this product to enter lakes, streams, ponds or other waters.
- **Do not make a foliar application of VIKING Rorvik Insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

**FOR USE ON GREEN ONIONS SUB GROUP – Crop group 3-07B**

**Spray Volume for Green Onion Subgroup:** Apply in a minimum finished spray volume of 100 L/ha by ground.

SITE	PEST	RATE VIKING Rorvik Insecticide PER HECTARE	SPECIFIC DIRECTIONS
Green onions sub group  Leek, Onion, green; Onion, Welsh, tops; Shallot, fresh leaves; Chive, fresh leaves; Chive, Chinese, fresh leaves; Elegans hosta, Fritillaria, Kurrat, Lady's leek, Onion, Beltsville bunching; Onion, fresh; Onion, macrostem; Onion, tree, tops; Wild leek	Cutworms  Leek moth (suppression)  Leafminers: <i>Liriomyza sativae</i> , <i>Liriomyza trifolii</i>	250 mL to 375 mL	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.

**RESTRICTIONS AND PRECAUTIONS: Green onions sub group**

- Do not make more than 4 applications per season.
- Do not apply more than once every 5 days.
- Do not apply less than 1 day before harvest. Observe a 1-day PHI.
- Do not exceed a total of 1.125 litres of **VIKING Rorvik Insecticide** per ha per season.
- Do not apply by air.
- **Do not make a foliar application of VIKING Rorvik Insecticide for a minimum of 60 days following an in-furrow or soil application or planting of seed or seed pieces treated with any Group 28 insecticide.**

**FOR USE ON GARDEN AND SUGAR BEETS**

**Spray Volume for Garden Beets and Sugar beets:** Apply in a minimum finished spray volume of 100 L/ha by ground or 50 L/ha by air.

SITE	PEST	RATE VIKING Rorvik Insecticide PER HECTARE	SPECIFIC DIRECTIONS

Sugar beets Garden beets	Beet webworm	250 mL to 375 mL	Begin applications when treatment thresholds have been reached. Thorough coverage is important to obtain optimum control.
<p><b>RESTRICTIONS AND PRECAUTIONS: Garden and Sugar Beet</b></p> <ul style="list-style-type: none"> <li>• Do not make more than 4 applications per season.</li> <li>• Do not apply more than once every 3 days.</li> <li>• Do not apply less than 1 day before harvest. Observe a 1-day PHI.</li> <li>• Do not exceed a total of 1.125 litres of <b>VIKING Rorvik Insecticide</b> per ha per season. Use the high rate under heavy pest pressure.</li> <li>• Thorough coverage is important to obtain optimum control.</li> </ul>			

### Tank Mix Options:

To provide a broader spectrum of pest control, **VIKING Rorvik Insecticide** may be tank-mixed with POUNCE® 384EC Insecticide.

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) fine classification. Boom height must be 60 cm or less above the crop or ground.

Aerial application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 km/h at flying height at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) fine classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotor span.

Apply only by fixed-wing or rotary aircraft equipment, which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

#### Use Precautions for Aerial Application

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the *National Aerial Pesticide Application Manual*, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

#### Operator Precautions for Aerial Application

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label. All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions for Aerial Application

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-587-743-1029 or obtain technical advice from the distributor or your provincial/territorial agricultural representative. Application of this specific product must meet and/or conform to the following:

Volume: Apply the recommended rate in a minimum spray volume of 50 litres per hectare.

**Spray Buffer Zones:**

A spray buffer zone is NOT required for:

- uses with hand-held application equipment permitted on this label

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of Application	Crop	Spray Buffer zones (meters) required for the protection of				
		Freshwater habitat of depths:		Estuarine/Marine Habitat of depths:		
		Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer	Root and tuber vegetables, Fruiting vegetables, Leafy vegetables, Brassica vegetables, Legume vegetables, Cucurbit vegetables, Corn, Grass forage, fodder, and hay, Nongrass animal feeds, Mint, Oilseeds, Okra, Hops, Globe artichokes, Green onions subgroup, sugar and garden beets, Cereal grains, Forage, fodder and straw of cereals	1	1	1	0	
Aerial	Potato	Fixed Wing	15	1	1	0

Aerial	Corn (field, pop, seed and sweet), Legume vegetables, Oilseeds, Cereal grains, Sugar beets and garden beets, Forage, fodder and straw of cereals	Rotary Wing	10	1	1	0
--------	--	-------------	----	---	---	---

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

**TANK MIXES:**

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Viking Crop Production Partners Inc. at 1-587-743-1029 for information before applying any tank mix that is not specifically recommended on this label.

**MIXING INSTRUCTIONS:**

Spray equipment must be clean and free of previous pesticide deposits before applying **VIKING Rorvik Insecticide**. Fill spray tank 1/4 to 1/2 full of water. Add **VIKING Rorvik Insecticide** directly to spray tank. Mix thoroughly to fully disperse the insecticide; once dispersed continued agitation is required. Once the product has been well mixed, add an adjuvant, where applicable, while continuously mixing. Recommendations regarding the use of adjuvants are provided under the Directions for Use Section of this label. Use mechanical or hydraulic means; do not use air agitation. Spray mix should not be stored overnight in spray tank.

**SPRAYER CLEANUP:**

Prior to application, start with clean, well maintained application equipment. Immediately following application, thoroughly clean all spray equipment to reduce the risk of forming hardened deposits which might become difficult to remove.

Drain spray equipment. Thoroughly rinse sprayer and flush hoses, boom and nozzles with clean water. Clean all other associated application equipment. Take all necessary safety precautions when cleaning equipment. Do not clean near wells, water sources or desirable vegetation. Dispose of waste rinse water in accordance with local regulations.

**RESISTANCE-MANAGEMENT RECOMMENDATIONS:**

For resistance management, **VIKING Rorvik Insecticide** is a Group 28 insecticide. Any insect population may contain individuals naturally resistant to **VIKING Rorvik Insecticide** and other Group 28 insecticides. The resistant individuals may dominate the insect population if this group of insecticides is used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance:

- Where possible, rotate the use of **VIKING Rorvik Insecticide** or other Group 28 insecticides with different groups that control the same pests.
- Make no more than 2 successive applications per generation to the same insect species on a crop. The following application to the target pest(s) must be with an effective product with a different mode of action.
- Use tank mixtures with insecticides from a different group that is effective on the target pest when such use is permitted.
- Insecticide use should be based on an IPM program that includes scouting and record keeping, and considers cultural, biological and other chemical control practices.
- Monitor treated pest populations for resistance development.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact Viking Crop Production Partners Inc. at 1-587-743-1029.

**NOTICE TO USER:**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

Viking (TM) is a trademark of Viking Crop Production Partners Inc.

All other products mentioned are trademarks of their respective holders.