

2022-2925  
2025-06-05

GROUP 7 3 4 FUNGICIDES

**\*Container label\***

# Raxil<sup>®</sup> Rise



## Fungicide

A Seed Treatment Fungicide for seed and seedling protection against certain seed-borne and soil-borne diseases on labeled crops.

SUSPENSION  
COMMERCIAL (AGRICULTURAL)

ACTIVE INGREDIENTS:	Tebuconazole	3.0 g/L
	Prothioconazole	15.4 g/L
	Penflufen	15.4 g/L
	Metalaxyl	6.2 g/L

Contains 1,2 benzisothiazolin-3-one at 0.035% and 5-chloro-2-methyl-4-isothiazolin-3-one at 0.0009% and 2-methyl-4-isothiazolin-3-one at 0.0003% as preservatives.

REGISTRATION NO. 35475 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

PROTECT FROM FREEZING

NET CONTENTS: 0.947 L – 1000 L

ALL SEED TREATED WITH THIS PRODUCT MUST BE CONSPICUOUSLY COLOURED

For wheat, barley, rye, triticale, and oats: On-farm or commercial seed treatment (facilities and mobile treaters) with open or closed transfer treatment equipment is permitted. Apply only in a way that this product will not contact workers or other persons, either directly or through drift. Only workers wearing personal protective equipment may be in the area when seed is being treated or bagged.

Bayer CropScience Inc.  
Suite 200, 160 Quarry Park Blvd. SE  
Calgary, AB T2C 3G3  
[www.cropscience.bayer.ca](http://www.cropscience.bayer.ca)

For product information, call: 1-888-283-6847  
In case of spills, poisoning, fire or other emergencies, call: 1-800-334-7577 (24 hours a day)

**GENERAL INFORMATION**

Raxil Rise is a seed treatment containing the fungicides penflufen, prothionconazole, tebuconazole and metalaxyl. Raxil Rise provides protection from seed, seedling and soil-borne diseases caused by certain *Ustilago* spp. *Tilletia* spp. *Fusarium* spp., *Pythium* spp., *Cochliobolus sativus*, *Pyrenophora* spp, *Penicillium* spp. and *Rhizoctonia solani* on labeled crops

#### PRECAUTIONS:

- KEEP OUT OF REACH OF CHILDREN.
- Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.
- Do not apply in a way that this product will contact workers or other persons. Only workers wearing personal protective equipment may be in the area when seed is being treated or bagged.
- Follow the personal protective equipment (PPE) and restriction requirements for the appropriate task and scenario:

Seed Type	Tasks	PPE/Engineering Controls
<b>For On-Farm Treatment (including handling and planting of seed treated on-farm)</b>		
Barley, oats, wheat, rye, triticale	Mixing, loading, treatment, and calibration	Open or closed transfer system is permitted.  Wear long pants, a long-sleeved shirt, chemical-resistant gloves, protective eyewear (goggles or face shield), socks and shoes.
	Cleaning, repairing, and activities involving handling and planting seeds treated on-farm	Wear a long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves.  Use a closed-cab tractor when planting. Gloves are not required with a closed cab.
<b>For Commercial Seed Treatment (facilities and mobile treaters)</b>		
Barley, oats, wheat, rye, triticale	Mixing, loading, treating, and calibrating	Open or closed transfer system is permitted.  Wear long pants, a long-sleeved shirt, chemical-resistant gloves, protective eyewear (goggles or face shield), socks and shoes.
	Cleaning and repairing	Wear chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes.
	Bagging, sewing, stacking, transferring seed to a storage bin, and any other	Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks, shoes and a suitable NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested.

	activities involving handling treated seed	
<b>For Planting and Handling Commercially Treated Seeds</b>		
Barley, oats, wheat, rye, triticale	Handling and planting	<p>Wear coveralls over a long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves.</p> <p>During handling of treated seeds, wear a NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested.</p> <p>Use a closed-cab tractor when planting. Gloves and respiratory protection are not required with a closed cab.</p>

- If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact Bayer CropScience Canada Inc. at 1-888-283-6847 or [www.cropscience.bayer.ca](http://www.cropscience.bayer.ca).

**FIRST AID:**

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION:**

There is no specific antidote if this product is ingested. Treat symptomatically.

**ENVIRONMENTAL PRECAUTIONS:**

Refer to the ENVIRONMENTAL PRECAUTIONS section of the booklet label for all environmental precautions and directions for use information.

**STORAGE:**

Store product in original container only, away from other pesticides, fertilizer, food or feed.

Store in a cool, dry place and avoid excessive heat. Keep container closed. Store this product away from food or feed.

## **DISPOSAL:**

### FOR NON-RETURNABLE CONTAINERS:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial or territorial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty, rinsed container unsuitable for further use.
4. Dispose of the container in accordance with provincial or territorial requirements.

### FOR RECYCLABLE CONTAINERS:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial **or territorial** requirements.

### FOR RETURNABLE CONTAINERS:

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

### FOR REFILLABLE CONTAINERS:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT use this container for any other purpose.

**TRIPLE RINSE INSTRUCTIONS (FOR RECYCLABLE AND DISPOSABLE CONTAINERS ONLY):** This container should be triple-rinsed prior to its disposal. The rinsate can be added to the seed treatment. Follow these steps for rinsing of containers:

1. Transfer Raxil Rise from the container to the seed treating equipment applicator tank.
2. Add rinse water to the container and agitate the contents.
3. Transfer the rinsate to the applicator tank.
4. Rinse as above 2 more times.
5. Thoroughly mix the Raxil Rise and rinse water in the applicator tank prior to use.

**CAUTION:** Be sure to adjust the undiluted application rate by the same volume of water added to compensate for the dilution of the product caused by the addition of rinsate.

**DISPOSAL OF UNUSED, UNWANTED PRODUCT:**

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for clean-up of spills.

**DISPOSAL OF TREATED SEED:**

Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. Dispose of seed packaging in accordance with local requirements. DO NOT re-use bags from treated seed to handle food or feed products.

**NOTICE TO USER:**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

**\*Booklet\***

# Raxil Rise



## Fungicide

A Seed Treatment Fungicide for seed and seedling protection against certain seed-borne and soil-borne diseases on labeled crops.

SUSPENSION

COMMERCIAL (AGRICULTURAL)

ACTIVE INGREDIENTS:	Tebuconazole	3.0 g/L
	Prothioconazole	15.4 g/L
	Penflufen	15.4 g/L
	Metalaxyl	6.2 g/L

Contains 1,2 benzisothiazolin-3-one at 0.035% and 5-chloro-2-methyl-4-isothiazolin-3-one at 0.0009% and 2-methyl-4-isothiazolin-3-one at 0.0003% as preservatives.

REGISTRATION NO. 35475 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

PROTECT FROM FREEZING

NET CONTENTS: 0.947 L – 1000 L

ALL SEED TREATED WITH THIS PRODUCT MUST BE CONSPICUOUSLY COLOURED

For wheat, barley, rye, triticale, and oats: On-farm or commercial seed treatment (facilities and mobile treaters) with open or closed transfer treatment equipment is permitted. Apply only in a way that this product will not contact workers or other persons, either directly or through drift. Only workers wearing personal protective equipment may be in the area when seed is being treated or bagged.

Bayer CropScience Inc.  
Suite 200, 160 Quarry Park Blvd. SE  
Calgary, AB T2C 3G3  
[www.cropscience.bayer.ca](http://www.cropscience.bayer.ca)

For product information, call: 1-888-283-6847

In case of spills, poisoning, fire or other emergencies,  
call: 1-800-334-7577 (24 hours a day)

<b>Table of Contents:</b>	<b>Section :</b>	<b>Page Number:</b>
<b>GENERAL INFORMATION</b>	1. The Product .....	3
<b>SAFETY AND HANDLING</b>	2. Precautions, Protective Clothing and Equipment .....	3
	3. First Aid and Toxicological Information .....	4
	4. Environmental Precautions and Information .....	4
	5. Storage .....	4
	6. Disposal .....	5
	7. Notices .....	6
<b>DIRECTIONS FOR USE</b>	8. Pests Controlled .....	6
	9. Recommended Uses .....	7
	10. Application Cautions .....	9
	11. Use Limitations .....	9
	12. Resistance Management Recommendations .....	10

## GENERAL INFORMATION

### Section 1: The Product

Raxil Rise is a seed treatment containing the fungicides penflufen, prothioconazole, tebuconazole and metalaxyl. Raxil Rise provides protection from seed, seedling and soil-borne diseases caused by certain *Ustilago* spp., *Tilletia* spp., *Fusarium* spp., *Pythium* spp., *Cochliobolus sativus*, *Pyrenophora* spp., *Penicillium* spp., and *Rhizoctonia solani* on labeled crops

## SAFETY AND HANDLING

### Section 2: Precautions, Protective Clothing and Equipment

#### PRECAUTIONS:

- KEEP OUT OF REACH OF CHILDREN.
- Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.
- Do not apply in a way that this product will contact workers or other persons. Only workers wearing personal protective equipment may be in the area when seed is being treated or bagged.
- Follow the personal protective equipment (PPE) and restriction requirements for the appropriate task and scenario:

Seed Type	Tasks	PPE/Engineering Controls
<b>For On-Farm Treatment (including handling and planting of seed treated on-farm)</b>		
Barley, oats, wheat, rye, triticale	Mixing, loading, treatment, and calibration	Open or closed transfer system is permitted.  Wear long pants, a long-sleeved shirt, chemical-resistant gloves, protective eyewear (goggles or face shield), socks and shoes.
	Cleaning, repairing, and activities involving handling and planting seeds treated on-farm	Wear a long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves.  Use a closed-cab tractor when planting. Gloves are not required with a closed cab.
<b>For Commercial Seed Treatment (facilities and mobile treaters)</b>		
Barley, oats, wheat, rye, triticale	Mixing, loading, treating, and calibrating	Open or closed transfer system is permitted.  Wear long pants, a long-sleeved shirt, chemical-resistant gloves, protective eyewear (goggles or face shield),

		socks and shoes.
	Cleaning and repairing	Wear chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes.
	Bagging, sewing, stacking, transferring seed to a storage bin, and any other activities involving handling treated seed	Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks, shoes and a suitable NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested.
<b>For Planting and Handling Commercially Treated Seeds</b>		
Barley, oats, wheat, rye, triticale	Handling and planting	Wear coveralls over a long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves.  During handling of treated seeds, wear a NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested.  Use a closed-cab tractor when planting. Gloves and respiratory protection are not required with a closed cab.

- If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact Bayer CropScience Canada Inc. at 1-888-283-6847 or [www.cropscience.bayer.ca](http://www.cropscience.bayer.ca).

### Section 3: First Aid and Toxicological Information

#### FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

#### TOXICOLOGICAL INFORMATION:

There is no specific antidote if this product is ingested. Treat symptomatically.

#### Section 4: Environmental Precautions

- TOXIC to non-target terrestrial plants and aquatic organisms.
- This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned up from the soil surface.

#### Section 5: Storage

Store product in original container only, away from food or feed. Store in a cool, dry place and avoid excessive heat. Keep container closed.

#### Section 6: Disposal

FOR NON-RETURNABLE CONTAINERS:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial or territorial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty, rinsed container unsuitable for further use.
4. Dispose of the container in accordance with provincial or territorial requirements.

FOR RECYCLABLE CONTAINERS:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with

provincial or territorial requirements.

#### FOR RETURNABLE CONTAINERS:

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

#### FOR REFILLABLE CONTAINERS:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT use this container for any other purpose.

**TRIPLE RINSE INSTRUCTIONS (FOR RECYCLABLE AND DISPOSABLE CONTAINERS ONLY):** This container should be triple-rinsed prior to its disposal. The rinsate can be added to the seed treatment. Follow these steps for rinsing of containers:

1. Transfer Raxil Rise from the container to the seed treating equipment applicator tank.
2. Add rinse water to the container and agitate the contents.
3. Transfer the rinsate to the applicator tank.
4. Rinse as above 2 more times.
5. Thoroughly mix the Raxil Rise and rinse water in the applicator tank prior to use.

**CAUTION:** Be sure to adjust the undiluted application rate by the same volume of water added to compensate for the dilution of the product caused by the addition of rinsate.

#### DISPOSAL OF UNUSED, UNWANTED PRODUCT:

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for clean-up of spills.

#### DISPOSAL OF TREATED SEED:

Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. Dispose of seed packaging in accordance with local requirements. DO NOT re-use bags from treated seed to

handle food or feed products.

**NOTICE TO USER:**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

## Section 7: Notices

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

### DIRECTIONS FOR USE

Raxil Rise is a ready-to-use seed treatment formulation for use in commercial seed treatment operations (facilities and mobile treaters) and for on-farm treatment with conventional seed treating equipment which can accurately meter, mix and apply flowable seed treatment formulations. Allow seeds to dry before bagging, storing or seeding. Uniform application to seed is necessary to ensure best disease protection and seed safety.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

## Section 8: Pests Controlled

Raxil Rise is a seed treatment containing the fungicides penflufen, prothioconazole, tebuconazole and metalaxyl. Raxil Rise provides protection from seed, seedling and soil-borne diseases caused by certain *Ustilago* spp., *Tilletia* spp., *Fusarium* spp., *Pythium* spp., *Cochliobolus sativus*, *Pyrenophora* spp., *Penicillium* spp., and *Rhizoctonia solani* on labeled crops.

## Section 9: Recommended Uses

CROP	PEST	APPLICATION RATE	DIRECTIONS
Barley	<p><b>Control:</b></p> <p>Seed rot/pre-emergent damping-off caused by seed- and soil-borne <i>Fusarium</i> spp., <i>Cochliobolus sativus</i> and soil-borne <i>Pythium</i> spp.</p> <p>Seedling blight caused by seed-borne <i>Fusarium</i> spp., <i>Cochliobolus sativus</i> and soil-borne, <i>Fusarium</i> spp. and <i>Pythium</i> spp.</p> <p>Post-emergent damping-off caused by seed- and soil-borne <i>Fusarium</i> spp. and <i>Cochliobolus sativus</i></p> <p>True loose smut (<i>Ustilago nuda</i>)</p> <p>Covered smut (<i>Ustilago hordei</i>)</p> <p>False loose smut (<i>Ustilago avenae</i>)</p> <p>Leaf stripe (<i>Pyrenophora graminea</i>)</p> <p>Seed rot/pre-emergent damping off and root rot caused by <i>Rhizoctonia solani</i></p> <p><b>Suppression:</b></p> <p>Root and crown rot caused by seed- and soil-borne <i>Fusarium</i> spp.</p> <p>Common root rot caused by seed- and soil-borne <i>Cochliobolus sativus</i></p> <p>Seedling blight caused by seed-borne <i>Penicillium</i> spp.</p>	Apply 325 mL per 100 kg of seed	<p>On-farm or commercial seed treatment (facilities and mobile treaters) with open or closed transfer treatment equipment is permitted.</p> <p>Use only in treating equipment which can accurately control application rates and provide good distribution of the chemical onto the seed in the mixing chamber.</p> <p>Uniform application to seed is necessary to ensure best disease protection and seed safety.</p>

CROP	PEST	APPLICATION RATE	DIRECTIONS
<b>Wheat, Oats, Rye, Triticale</b>	<p><b>Control:</b></p> <p>Seed rot/pre-emergent damping-off caused by seed- and soil-borne <i>Fusarium</i> spp., <i>Cochliobolus sativus</i> and soil-borne <i>Pythium</i> spp.</p> <p>Seedling blight caused by seed-borne <i>Fusarium</i> spp., <i>Cochliobolus sativus</i> and soil-borne, <i>Fusarium</i> spp. and <i>Pythium</i> spp.</p> <p>Post-emergent damping-off caused by seed- and soil-borne <i>Fusarium</i> spp. and <i>Cochliobolus sativus</i></p> <p>Loose smut (<i>Ustilago tritici</i>, <i>U. avenae</i>)</p> <p>Covered smut (<i>Ustilago kollerii</i>) on oats</p> <p>Common bunt (<i>Tilletia caries</i>, <i>T. laevis</i>) on wheat, rye, triticale</p> <p>Seed rot/pre-emergent damping off and root rot caused by <i>Rhizoctonia solani</i></p> <hr/> <p><b>Suppression:</b></p> <p>Root and crown rot caused by seed- and soil-borne <i>Fusarium</i> spp.</p> <p>Common root rot caused by seed- and soil-borne <i>Cochliobolus sativus</i></p> <p>Seedling blight caused by seed-borne <i>Penicillium</i> spp.</p>	<p>Apply 325 mL per 100 kg of seed</p>	<p>On-farm or commercial seed treatment (facilities and mobile treaters) with open or closed transfer treatment equipment is permitted.</p> <p>Use only in treating equipment which can accurately control application rates and provide good distribution of the chemical onto the seed in the mixing chamber.</p> <p>Uniform application to seed is necessary to ensure best disease protection and seed safety.</p>

## TANK-MIXTURES

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions and Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

For control of certain insect pests, Raxil Rise may be combined with certain seed treatment insecticides.

For recommended tank-mixes, read and observe all label directions, including rates, personal protective equipment, restrictions and precautions for each product used in the tank-mix. Always use in accordance with the most restrictive label restrictions and precautions.

<b>CROP</b>	<b>TANK-MIX PARTNER</b>
Wheat, Barley, Oats	Stress Shield 600

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Bayer CropScience Inc. at 1-888-283-6847 or [www.cropscience.bayer.ca](http://www.cropscience.bayer.ca) for information before applying any tank mix that is not specifically recommended on this label.

Bayer CropScience Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) of all potential tank mixes under all environmental conditions or for all crop varieties. Conduct a germination test on a small portion of seed before committing the total seed lot to tank mixes that are not specifically listed on this label.

To determine the physical compatibility of this product with other products, use a jar test.

### Section 10: Application Cautions

**SEED QUALITY:** Lab and field studies have shown that Raxil Rise treated seed can be stored without loss in germination or disease performance. Many seed types are prone to mechanical damage during seed handling, and a reduction in germination within a seed lot can occur should handling cause seed damage irrespective of seed treatment application. Raxil Rise-treated or untreated seed may drop in germination with age and germination drops are accelerated under adverse storage conditions or in seed with a high level of mechanical damage. Use handling

equipment designed to minimize mechanical damage when moving seed. Due to seed quality and seed storage conditions beyond the control of Bayer CropScience, no claims are made to guarantee the germination of carry-over seed or propagating materials for all crop seed. Treatment of highly mechanically scarred or damaged seed, or seed known to be of low vigour and poor quality, may result in reduced germination and/or reduction of seed and seedling vigour. Treat a small quantity of seed using equipment similar to that planned for treating the total seed lot. Conduct germination tests on a small portion of seed before committing the total seed lot to a selected seed treatment.

## Section 11: Use Limitations

### USE RESTRICTIONS:

- Treated seed must not be used for food, feed or oil processing.
- Barley, wheat, oats, rye, and triticale may be replanted at any time. All other crops may be planted after 30 days.

### LABELLING TREATED SEED:

All containers or packages containing treated wheat, barley, oat, rye or triticale seed for sale or use in Canada must be labelled with the following information:

- This seed has been treated with Raxil Rise, containing prothioconazole, tebuconazole, penflufen and metalaxyl.
- **KEEP TREATED SEED OUT OF REACH OF CHILDREN AND ANIMALS.**
- **DO NOT** use for food, feed and oil processing.
- Store away from feed and other foodstuffs.
- During handling and planting treated seed, wear coveralls over a long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves. During handling of treated seeds, wear a NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested.
- Use a closed-cab tractor when planting. Gloves and respiratory protection are not required with a closed cab.
- Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil.

## Section 12: Resistance Management Recommendations

For resistance management, please note that Raxil Rise contains Groups 3, 4 and 7 fungicides. Any fungal population may contain individuals naturally resistant to Raxil Rise and other Group 3, 4 and 7 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

- Where possible, rotate the use of Raxil Rise or other Group 3, 4 and 7 fungicides with different

groups that control the same pathogens.

- Use tank mixtures with fungicides from a different group that is effective on the target pathogen when such use is permitted.
- Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications.
- Monitor fungal populations for resistance development. Notify Bayer CropScience if reduced sensitivity of the pathogen to Raxil Rise is suspected.
- If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different site of action, if available.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information and to report suspected resistance contact Bayer CropScience Inc. at 1-888-283-6847 or at [www.cropscience.bayer.ca](http://www.cropscience.bayer.ca).