

2025-1687
2025-04-25

GROUP	27	HERBICIDE
-------	----	-----------

NCS COMETA 480 Herbicide

COMMERCIAL
AGRICULTURAL

SUSPENSION

A herbicide for use in field, seed, and sweet corn, flax, cranberry, asparagus (established), blueberry, rhubarb, pearl millet, sorghum and turf for control of annual broadleaf weeds.

ACTIVE INGREDIENT:

Mesotrione 480 g/L

Contains 1,2-benzisothiazolin-3-one at 0.0097 % as a preservative.

**READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

CAUTION: CORROSIVE

POTENTIAL SKIN SENSITIZER

REGISTRATION NO.: 35492 PEST CONTROL PRODUCTS ACT

NET CONTENTS: **1 L - Bulk**

Northern CropScience Inc.
Suite 300, 336 Sheppard Avenue East
North York, Ontario
M2N 3B4
Tel: 437-880-8283

FOR CHEMICAL EMERGENCY: spill, leak, fire, exposure or accident, call CHEMTREC 1-800-424-9300.

For product and use information, call Northern CropScience Inc.: 1-437-880-8283.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

Treat symptomatically if ingested or in contact with eyes or skin. If gastric lavage is considered necessary, prevent aspiration of gastric material. Administration of activated charcoal may also be considered.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

Avoid working in spray mist. May irritate eyes. Avoid contact with eyes. Potential skin sensitizer.

DO NOT apply by air.

DO NOT apply with airblast equipment.

DO NOT use or store product near heat or open flame. **DO NOT** apply beyond the field boundary.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Northern CropScience Inc. at 1-437-880-8283.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair. Gloves are not required during application within a closed cab.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Wash splashes from skin IMMEDIATELY with plenty of water. Remove PPE immediately after handling this product. Wash the outside of gloves with soap and water before removing. As soon as possible, wash thoroughly and change into clean clothing. After spraying, wash hands and shower thoroughly with soap and water.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Wear freshly laundered clothes daily.

While using product, do not eat, drink or smoke. Wash hands and exposed skin with soap and water thoroughly before eating, drinking, smoking, applying cosmetics or using the toilet.

ENVIRONMENTAL PRECAUTIONS

Toxic to small wild mammals, aquatic organisms and to non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

STORAGE

Store this product away from food or feed

Always store product in original container with top closed in a cool, dry place. NCS COMETA 480 Herbicide can be stored in freezing conditions. If frozen, allow to thaw and agitate thoroughly prior to use.

DISPOSAL

Disposal of unused, unwanted product

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean up of spills.

For recyclable containers

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

For returnable containers

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For refillable containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. **DO NOT** reuse this container for any other purpose.

NCS COMETA 480 Herbicide

COMMERCIAL
AGRICULTURAL

SUSPENSION

A herbicide for use in field, seed, and sweet corn, flax, cranberry, asparagus (established), blueberry, rhubarb, pearl millet, sorghum and turf for control of annual broadleaf weeds.

ACTIVE INGREDIENT:

Mesotrione 480 g/L

Contains 1,2-benzisothiazolin-3-one at 0.0097 % as a preservative.

**READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

**CAUTION: CORROSIVE
POTENTIAL SKIN SENSITIZER**

REGISTRATION NO.: 35492 PEST CONTROL PRODUCTS ACT

Northern CropScience Inc.
Suite 300, 336 Sheppard Avenue East
North York, Ontario
M2N 3B4
Tel: 437-880-8283.

FOR CHEMICAL EMERGENCY: spill, leak, fire, exposure or accident, call CHEMTREC 1-800-424-9300.

For product and use information, call Northern CropScience Inc.: 1-437-880-8283.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

Treat symptomatically if ingested or in contact with eyes or skin. If gastric lavage is considered necessary, prevent aspiration of gastric material. Administration of activated charcoal may also be considered.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

Avoid working in spray mist. May irritate eyes. Avoid contact with eyes. Potential skin sensitizer.

DO NOT apply by air.

DO NOT apply with airblast equipment.

DO NOT use or store product near heat or open flame. **DO NOT** apply beyond the field boundary.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Northern CropScience Inc. at 1-437-880-8283.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair. Gloves are not required during application within a closed cab.

Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Wash splashes from skin IMMEDIATELY with plenty of water. Remove PPE immediately after handling this product. Wash the outside of gloves with soap and water before removing. As soon as possible, wash thoroughly and change into clean clothing. After spraying, wash hands and shower thoroughly with soap and water.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Wear freshly laundered clothes daily.

While using product, do not eat, drink or smoke. Wash hands and exposed skin with soap and water thoroughly before eating, drinking, smoking, applying cosmetics or using the toilet.

ENVIRONMENTAL PRECAUTIONS

Toxic to small wild mammals, aquatic organisms and to non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

STORAGE

Store this product away from food or feed

Always store product in original container with top closed in a cool, dry place. NCS COMETA 480 Herbicide can be stored in freezing conditions. If frozen, allow to thaw and agitate thoroughly prior to use.

DISPOSAL

Disposal of unused, unwanted product

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean up of spills.

For recyclable containers

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.

2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

For returnable containers

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For refillable containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. **DO NOT** reuse this container for any other purpose.

PRODUCT INFORMATION

NCS COMETA 480 Herbicide is a suspension concentrate herbicide that has soil residual activity and provides selective control or suppression of annual broadleaf weeds in corn, and other listed crops. NCS COMETA 480 Herbicide has a high level of crop safety to allow for flexibility in application timing in corn, and cranberries.

In post-emergent applications, NCS COMETA 480 Herbicide is absorbed through the leaves and is rapidly translocated to the plant's growing points.

NCS COMETA 480 Herbicide may also be tank mixed with a post-emergent grass herbicide for broader spectrum post-emergent weed control in corn.

Use Precautions

NCS COMETA 480 Herbicide may be used pre-emergent on field corn, production seed corn and sweet corn. Post-emergent applications may be made on field corn, and on seed and sweet corn (see Minor Use section). Not all seed corn inbreds and sweet corn varieties have been tested; the use of NCS COMETA 480 Herbicide must be approved by the seed company and comply with the directions given by the seed company (corn).

Although NCS COMETA 480 Herbicide has a flexible recropping profile, certain crops may be sensitive to low concentrations in the soil. Therefore, careful consideration should be given to crop rotation plans prior to using NCS COMETA 480 Herbicide (refer to "*RECROPPING GUIDELINES*").

Dry conditions following application may reduce the pre-emergent activity of NCS COMETA 480 Herbicide. If an activating rain (12 mm) is not received within 7-10 days after a pre-emergent application, rotary hoeing is suggested to activate the herbicide.

Do not apply to soils that contain less than 1% or more than 10% organic matter.

When applied post emergent, thorough coverage of emerged weeds is essential for effective control with NCS COMETA 480 Herbicide. Although weed competition is quickly halted, visual symptoms of dying weeds (discolouration) may take up to 2 weeks to appear depending on the weed species and growing conditions.

Optimum control of emerged weeds will be obtained when weeds are actively growing. Under unfavourable conditions, such as drought, heat, flooding, prolonged cool temperatures or insufficient fertility, adequate control may not be achieved and re-growth may occur.

Temporary crop injury (bleaching) may occur under extreme weather conditions or when the crop is under stress due to inadequate or abnormally high moisture levels or extreme temperatures. The symptoms are most visible where excessive rates have been applied, such as sprayer overlaps. Corn quickly outgrows these effects and continues to grow normally.

Do not make a foliar post-emergent application of any organophosphate or carbamate insecticide within 7 days before or 7 days after a NCS COMETA 480 Herbicide application or severe injury may occur to corn. Northern CropScience Inc. will not be held responsible for losses or damage resulting from such use.

NCS COMETA 480 Herbicide may be applied prior to or following a pyrethroid-type insecticide such as FORCE® EVO Insecticide or MATADOR® 120EC Insecticide.

Do not cultivate corn within 7 days before or after a post-emergent NCS COMETA 480 Herbicide application.

Make only one application of NCS COMETA 480 Herbicide per year in all crops except in cranberries for control of cinquefoil, goldenrod, rushes and sedges where a second application is required.

DIRECTIONS FOR USE

General Restrictions

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Northern CropScience Inc. at 1-437-880-8283 for information before applying any tank mix that is not specifically recommended on this label.

Northern CropScience Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) of all potential tank mixes under all environmental conditions or for all crop varieties. Tank mixes that are not specifically listed on this label should be tested on a small area first, under local conditions and using standard practices, to confirm the tank mix is suitable for widespread application.

To determine the physical compatibility of this product with other products, use a jar test.

Always read and follow label directions including WALES mixing order.

Ground Application

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

Mixing Instructions

1. Ensure that the sprayer is totally clean.
2. Fill the sprayer half full with water. Engage gentle agitation.
3. Ensure that the agitation system is working properly and that it creates a rippling or rolling action on the water surface.
4. If required, add Ultim[®] Herbicide or Accent[®] Herbicide slowly and agitate until completely dispersed.
5. Add NCS COMETA 480 Herbicide and PRIMEXTRA[®] II MAGNUM Herbicide or AATREX[®] Liquid 480 herbicide (if an applicable tank-mix) slowly and agitate until completely dispersed. To ensure complete dispersion, wait 2 or 3 minutes after the last of the herbicide has been added to the tank.
6. Fill the tank to three quarters full with water.
7. Add DUAL II MAGNUM[®] Herbicide if required.
8. Add the non-ionic surfactant if applicable (e.g., AGRAL[®] 90 Non-ionic Wetting & Spreading Agent Liquid) at 0.2% v/v.
9. Add glyphosate-containing product if required. Refer to the label of the tank mix partner and the surfactant for the specific mixing instructions.
10. Continue agitation while completing the filling of the sprayer with water.
11. Always ensure that agitation is maintained until spraying is completed, even if stopped for brief periods of time.
12. If agitation is stopped for more than 5 minutes, re-suspend the spray solution by running on full agitation prior to spraying. **NCS COMETA 480 Herbicide must be sprayed the same day as mixing.**
13. Do not mix, load or clean spray equipment where there is a potential to contaminate wells or aquatic systems.

Spraying Instructions

1. **Water Volume:** Apply in 100 – 200 litres per hectare.
2. **Spray Pressure:** 206 – 300 kPa
3. **Spray Nozzles:** Flat Fan with 50 mesh or larger screens. For uniform coverage and increased penetration of the crop canopy, set the nozzles at a downward angle of 90 degrees (90°).
4. **Surfactant:** Post-emergent applications of NCS COMETA 480 Herbicide must be applied with the non-ionic surfactant such as AGRAL 90 Non-ionic Wetting & Spreading Agent Liquid at 0.2% v/v (2.0 L for each 1000 L of spray mixture).

Equipment Clean-Up

Before Spraying:

- Prior to using NCS COMETA 480 Herbicide, ensure that the spray tank, lines and filter are thoroughly clean.

After Spraying:

- Thoroughly clean application equipment immediately after spraying. **DO NOT** allow NCS COMETA 480 Herbicide residue to dry within the spray tank
- When using tank mixes, consult the tank-mix partner label for additional clean-up instructions.
- The following recommendations are provided:

1. Drain and flush tank walls, boom and all hoses for ten minutes with a clean water/detergent mixture. Rinse with clean water. **DO NOT** clean the sprayer near desirable vegetation, wells or other water sources.
2. Remove all nozzles and screens and wash separately.
3. Dispose of all rinsate in accordance with provincial regulations.

Application in Liquid Fertilizers

Nitrogen solutions (such as 28-0-0 UAN), excluding suspension fertilizers, may replace water as a carrier for pre-emergence application in corn. Do not use nitrogen solutions as a carrier to corn that has emerged. Constant agitation is necessary to maintain a uniform mixture during application.

Spray Buffer zones

A spray buffer zone is NOT required for uses with hand-held application equipment permitted on this label.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop	Spray Buffer zone (metres) required for the protection of:	
		Aquatic habitat	Terrestrial habitat
Field sprayer	Field corn, seed and sweet corn, flax, asparagus (established), rhubarb, pearl millet, sorghum, highbush and lowbush blueberry, and turf	1	4
	Cranberry	1	10

When tank mixes are permitted, consult the labels of the tankmix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

Pre-Harvest Intervals (PHI)

Crops	PHI (days)
Field corn grain/stover	100
Field corn forage	90
Sweet corn	50

CROP USE DIRECTIONS

Corn

Pre-emergent Application: (Field Corn, Production Seed Corn and Sweet Corn) – Eastern Canada Only

Apply NCS COMETA 480 Herbicide at 0.30 L per hectare in a spray volume of 200 L and a minimum pressure of 206 kPa (refer to the “*MIXING AND SPRAYING INSTRUCTIONS*” section of this label).

NCS COMETA 480 Herbicide will control:

lamb's-quarters	wild mustard
redroot pigweed	common ragweed (suppression)
velvetleaf	

NCS COMETA 480 Herbicide tank-mixed with DUAL II MAGNUM Herbicide, PRIMEXTRA II MAGNUM Herbicide, DUAL II MAGNUM Herbicide plus AATREX LIQUID 480 Herbicide will provide broad-spectrum control of annual broadleaf and grass weeds. Consult the tank mix partner label for precautions, use rates and weeds controlled.

Use Rates for Pre-emergent Application

Field Corn, Production Seed Corn and Sweet Corn – Pre-emergent	
APPLICATION TIMING	Pre-emergent
PRODUCT RATE	0.30 L/ha
WEEDS CONTROLLED	Broadleaf Weeds
APPLICATION METHOD	Ground application only. Apply in a minimum of 100 L of water per hectare.
APPLICATION TIMING	Pre-emergent
PRODUCT RATE	0.30 L/ha
TANK MIX PARTNER RATE	1.25 – 1.75 L/ha DUAL II MAGNUM Herbicide
WEEDS CONTROLLED	Broadleaf Weeds and Annual Grasses
APPLICATION METHOD	Ground application only. Apply in a minimum of 100 L of water per hectare.
APPLICATION TIMING	Pre-emergent
PRODUCT RATE	0.30 L/ha
TANK MIX PARTNER RATE	3.0 – 4.0 L/ha PRIMEXTRA II MAGNUM Herbicide
WEEDS CONTROLLED	Broadleaf Weeds and Annual Grasses
APPLICATION METHOD	Ground application only. Apply in a minimum of 100 L of water per hectare.
APPLICATION TIMING	Pre-emergent
PRODUCT RATE	0.30 L/ha
TANK MIX PARTNER RATE	1.25 – 1.75 L/ha DUAL II MAGNUM Herbicide + 2.1 – 3.1 L/ha AATREX Liquid 480 Herbicide
WEEDS CONTROLLED	Broadleaf Weeds and Annual Grasses
APPLICATION METHOD	Ground application only. Apply in a minimum of 100 L of water per hectare.
NOTES	Consult the label of the product(s) tank mixed with NCS COMETA 480 Herbicide for specific instructions regarding grass and broadleaf weeds, soil-type variations, and rotational crop, grazing or other restrictions.

Early Post-emergent Application: (Field Corn only) - Eastern Canada only– For Seed and Sweet corn uses and crop stages see MU section

Apply NCS COMETA 480 Herbicide at 0.30 L per hectare in a spray volume of 200 L and a minimum pressure of 206 kPa (refer to the “*MIXING AND SPRAYING INSTRUCTIONS*” section of this label).

NCS COMETA 480 Herbicide will control:

lamb’s-quarters	wild mustard
redroot pigweed	common ragweed (suppression)
velvetleaf	

NCS COMETA 480 Herbicide tank-mixed with DUAL II MAGNUM Herbicide, PRIMEXTRA II MAGNUM Herbicide, DUAL II MAGNUM Herbicide plus AATREX Liquid 480 Herbicide will provide broad-spectrum control of annual broadleaf and grass weeds. Consult the tank mix partner labels for precautions, use rates and weeds controlled.

**Use Rates for Early Post-emergent Application
(Spike to 2 leaf stage of corn and cotyledon to 2 leaf stage of weeds)**

Field Corn Only – Early Post-emergent Application	
APPLICATION TIMING	Early Post-emergent
PRODUCT RATE	0.30 L/ha
WEEDS CONTROLLED	Broadleaf Weeds
APPLICATION METHOD	Ground application only. Apply in a minimum of 100 L of water per hectare.
APPLICATION TIMING	Early Post-emergent
PRODUCT RATE	0.30 L/ha
TANK MIX PARTNER RATE	1.25 – 1.75 L/ha DUAL II MAGNUM Herbicide
WEEDS CONTROLLED	Broadleaf Weeds and Annual Grasses
APPLICATION METHOD	Ground application only. Apply in a minimum of 100 L of water per hectare.
APPLICATION TIMING	Early Post-emergent
PRODUCT RATE	0.30 L/ha
TANK MIX PARTNER RATE	3.0 – 4.0 L/ha PRIMEXTRA II MAGNUM Herbicide
WEEDS CONTROLLED	Broadleaf Weeds and Annual Grasses
APPLICATION METHOD	Ground application only. Apply in a minimum of 100 L of water per hectare.
APPLICATION TIMING	Early Post-emergent
PRODUCT RATE	0.30 L/ha
TANK MIX PARTNER RATE	1.25 – 1.75 L/ha DUAL II MAGNUM Herbicide + 2.1 – 3.1 L/ha AATREX Liquid 480 Herbicide
WEEDS CONTROLLED	Broadleaf Weeds and Annual Grasses

APPLICATION METHOD	Ground application only. Apply in a minimum of 100 L of water per hectare.
NOTES	Consult the label of the product(s) tank mixed with NCS COMETA 480 Herbicide for specific instructions regarding grass and broadleaf weeds, soil type variations, and rotational crop, grazing or other restrictions.

Late Post-emergent Application: (Field Corn only) - Eastern Canada only – For Seed and Sweet corn uses and crop stages see MU section.

Apply NCS COMETA 480 Herbicide at 0.21 L per hectare in a spray volume of 100-200 L of water and a minimum pressure of 206 kPa (refer to the “*MIXING AND SPRAYING INSTRUCTIONS*” section of this label).

Always add a spray surfactant. Use the non-ionic surfactant AGRAL 90 Non-ionic Wetting & Spreading Agent Liquid at 0.20% v/v.

For best results, apply NCS COMETA 480 Herbicide to actively growing weeds. Weeds that emerge after an application of NCS COMETA 480 Herbicide will be controlled after they absorb the herbicide from the soil provided there is sufficient moisture for uptake.

NCS COMETA 480 Herbicide will control:

eastern black nightshade	redroot pigweed
velvetleaf	common ragweed (suppression only)

up to the 8 leaf weed stage.

NCS COMETA 480 Herbicide tank-mixed with AATREX Liquid 480 Herbicide will provide control of all four weeds listed above. NCS COMETA 480 Herbicide tank-mixed with AATREX Liquid 480 Herbicide and a glyphosate containing product will provide control of all four weeds listed above plus emerged annual and perennial weeds for use ONLY on glyphosate tolerant corn. NCS COMETA 480 Herbicide tank-mixed with Ultim Herbicide or Accent Herbicide will provide broad-spectrum control of annual broadleaf and grass weeds. Consult tank mix partner labels for precautions. Consult Ultim Herbicide and Accent Herbicide labels for use rates and weeds controlled.

**Use Rates for Post Emergent Application
(3 to 8 leaf stage of corn and up to 8 leaf stage of broadleaf weeds)**

Field Corn Only – Post-emergent Application	
APPLICATION TIMING	Post-emergent
PRODUCT RATE	0.21 L/ha
TANK MIX PARTNER RATE	0.2% v/v AGRAL 90 Non-ionic Wetting & Spreading Agent Liquid
WEEDS CONTROLLED	Eastern black nightshade, redroot pigweed, velvetleaf
WEEDS SUPPRESSED	Common ragweed
APPLICATION METHOD	Ground application only. Apply in a minimum of 100 L of water per hectare.
APPLICATION TIMING	Post-emergent
PRODUCT RATE	0.21 L/ha
TANK MIX PARTNER RATE	33.3 g/ha Ultim Herbicide + 0.2% v/v AGRAL 90 Non-ionic Wetting & Spreading Agent Liquid

WEEDS CONTROLLED	Weeds controlled by NCS COMETA 480 Herbicide plus weeds controlled by Ultim Herbicide (up to the 6-leaf stage of corn only)
APPLICATION METHOD	Ground application only. Apply in a minimum of 100 L of water per hectare.

APPLICATION TIMING	Post-emergent
PRODUCT RATE	0.21 L/ha
TANK MIX PARTNER RATE	33.4 g/ha Accent Herbicide + 0.2% v/v AGRAL 90 Non-ionic Wetting & Spreading Agent Liquid
WEEDS CONTROLLED	Weeds controlled by NCS COMETA 480 Herbicide plus weeds controlled by Accent Herbicide
APPLICATION METHOD	Ground application only. Apply in a minimum of 100 L of water per hectare.
APPLICATION TIMING	Post-emergent
PRODUCT RATE	0.21 L/ha
TANK MIX PARTNER RATE	1.67 L/ha glyphosate (of a 540 g ae/L formulation) + 0.2% v/v AGRAL 90 Non-ionic Wetting & Spreading Agent Liquid Some perennial weeds may not be controlled with this rate of glyphosate. Consult the glyphosate product label regarding perennial weeds controlled. For glyphosate formulations that are not 540 g ae/L, refer to the glyphosate product label to determine the appropriate rate for the target weeds.
WEEDS CONTROLLED	Weeds controlled by NCS COMETA 480 Herbicide plus emerged annual and perennial weeds. For use ONLY on glyphosate tolerant corn.
APPLICATION METHOD	Ground application only. Apply in a minimum of 100 L of water per hectare.

MINOR USES

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Northern CropScience Inc. under the User Requested Minor Use Label Expansion program. For these uses, Northern CropScience Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

Field Corn (Eastern Canada Only)

CROP	Field Corn (Eastern Canada Only) – apply at 3 – 8 leaf stage of corn
WEEDS CONTROLLED	Tall Waterhemp
TIMING	Pre-emergent
PRODUCT RATE	0.30 L/ha
TANK MIX PARTNER RATE	3 – 4 L/ha PRIMEXTRA II MAGNUM Herbicide in 200 L carrier/ha Use higher rate of PRIMEXTRA II MAGNUM Herbicide for heavier weed problems.
REMARKS	Make only one application per year. Apply by ground equipment only.
TIMING	Post-emergent Apply at up to the 8-leaf stage of tall waterhemp.
NOTES 1. Consult tank mix partner label for precautions.	

Seed and Sweet Corn (Eastern Canada and British Columbia Only)

CROP	Seed and Sweet Corn (Eastern Canada and British Columbia Only)
TIMING	Post-emergent Late post-emergence (3-6 leaf stage)
PRODUCT RATE	0.21 L/ha
TANK MIX PARTNER RATE	0.2% v/v AGRAL 90 Non-ionic Wetting & Spreading Agent Liquid
REMARKS	Make only one application per year. Apply in 100-200 L water/ha at 205-300 kPa. Do not harvest sweet corn treated with NCS COMETA 480 Herbicide within 50 days of treatment. Harvest seed corn at the normal interval. Seed corn inbreds and sweet corn varieties may vary in their tolerance to herbicides, including NCS COMETA 480 Herbicide. Since not all seed corn inbreds or sweet corn varieties have been tested for tolerance to NCS COMETA 480 Herbicide, first use of NCS COMETA 480 Herbicide should be limited to a small area of each variety to confirm tolerance prior to adoption as a general field practice. Additionally, consult your seed supplier for information on the tolerance of specific seed corn inbreds to NCS COMETA 480 Herbicide. DO NOT apply on Delmonte 2038 as unacceptable crop injury will occur.
NOTES 1. Consult tank mix partner label for precautions.	

Established Cranberry Beds

CROP	Established Cranberry Beds
WEEDS CONTROLLED	Lamb's-quarters, redroot pigweed, velvetleaf, and wild mustard
WEEDS SUPPRESSED	Common ragweed, vetch and creeping buttercup
TIMING	Pre-emergent (up to 2 leaf stage) and early post-emergent to the weeds
PRODUCT RATE	0.30 L/ha
REMARKS	<p>Make only one application per year except for control of cinquefoil, goldenrod, rushes and sedges where a second application is required.</p> <p>Allow for a minimum of 14 days between applications.</p> <p>Apply in 200 L water/ha for pre-emergence application and in 100 - 200 L water/ha for post-emergence application at 206 - 300 kPa.</p> <p>DO NOT apply NCS COMETA 480 Herbicide through any type of irrigation equipment.</p> <p>Bearing Beds – Apply between bud break and fruit set. Do not harvest within 60 days of treatment.</p> <p>Non-Bearing Beds – Apply between bud break and 60 days prior to fall or winter flooding. DO NOT harvest within 365 days of treatment. After fall application of NCS COMETA 480 Herbicide, do not apply again the following spring.</p> <p>DO NOT flood within 60 days of treatment.</p> <p>DO NOT apply directly to water or areas where surface water is present.</p> <p>DO NOT spray to runoff.</p> <p>DO NOT apply if rainfall or sprinkler irrigation for frost protection is expected within 48 hours.</p> <p>Wait at least 18 months after the last NCS COMETA 480 Herbicide application in cranberries before replanting cranberries.</p>

OR	
WEEDS CONTROLLED	Eastern black nightshade, redroot pigweed, and velvetleaf
WEEDS SUPPRESSED	Common ragweed, vetch and creeping buttercup
TIMING	Late post emergent (3 to 8 leaf weed stage) to the weeds
PRODUCT RATE	0.21 L/ha
TANK MIX PARTNER RATE	0.2% v/v AGRAL 90 Non-ionic Wetting & Spreading Agent Liquid
REMARKS	<p>Make only one application per year except for control of cinquefoil, goldenrod, rushes and sedges where a second application is required.</p> <p>Allow for a minimum of 14 days between applications.</p> <p>Apply in 200 L water/ha for pre-emergence application and in 100 - 200 L water/ha for post-emergence application at 206 - 300 kPa.</p> <p>DO NOT apply NCS COMETA 480 Herbicide through any type of irrigation equipment.</p> <p>Bearing Beds – Apply between bud break and fruit set. Do not harvest within 60 days of treatment.</p> <p>Non-Bearing Beds – Apply between bud break and 60 days prior to fall or winter flooding. DO NOT harvest within 365 days of treatment. After fall application of NCS COMETA 480 Herbicide, do not apply again the following spring.</p> <p>DO NOT flood within 60 days of treatment.</p> <p>DO NOT apply directly to water or areas where surface water is present.</p> <p>DO NOT spray to runoff.</p> <p>DO NOT apply if rainfall or sprinkler irrigation for frost protection is expected within 48 hours.</p> <p>Wait at least 18 months after the last NCS COMETA 480 Herbicide application in cranberries before replanting cranberries.</p>

OR	
WEEDS CONTROLLED	Eastern black nightshade, redroot pigweed, velvetleaf. goldenrod
WEEDS SUPPRESSED	Common ragweed, vetch, creeping buttercup, cinquefoil (<i>Potentilla</i> spp.), rushes (<i>Juncus Canadensis</i> , <i>J. effuses</i> , <i>J. bufonlus</i> , <i>J. tenuis</i>), and sedges (<i>Carex</i> spp.)
TIMING	Post emergent (3 to 8 leaf weed stage) to the weeds
PRODUCT RATE	2 applications of 0.21 L/ha
TANK MIX PARTNER RATE	0.2% v/v AGRAL 90 Non-ionic Wetting & Spreading Agent Liquid

REMARKS	<p>Make only one application per year except for control of cinquefoil, goldenrod, rushes and sedges where a second application is required.</p> <p>Allow for a minimum of 14 days between applications.</p> <p>Apply in 200 L water/ha for pre-emergence application and in 100 - 200 L water/ha for post-emergence application at 206 - 300 kPa.</p> <p>DO NOT apply NCS COMETA 480 Herbicide through any type of irrigation equipment.</p> <p>Bearing Beds – Apply between bud break and fruit set. Do not harvest within 60 days of treatment.</p> <p>Non-Bearing Beds – Apply between bud break and 60 days prior to fall or winter flooding.</p> <p>DO NOT harvest within 365 days of treatment.</p> <p>DO NOT flood within 60 days of treatment.</p> <p>DO NOT apply directly to water or areas where surface water is present.</p> <p>DO NOT spray to runoff.</p> <p>DO NOT apply if rainfall or sprinkler irrigation for frost protection is expected within 48 hours.</p> <p>Wait at least 18 months after the last NCS COMETA 480 Herbicide application in cranberries before replanting cranberries.</p>
NOTES	1. Consult tank mix partner label for precautions.

Asparagus (Established)	
CROP	Asparagus (Established)
WEEDS CONTROLLED	Lamb's-quarters, redroot pigweed, velvetleaf, wild mustard
WEEDS SUPPRESSED	Common ragweed
TIMING	Pre-emergent to asparagus (established)
PRODUCT RATE	0.30 L/ha
REMARKS	<p>Can be applied by broadcast or banded application. For banded applications, the application must be made to account for band width (i.e. to deliver desired rate per hectare).</p> <p>Apply after fern mowing, disking, or other tillage operation but prior to spear emergence.</p> <p>Apply in 100 – 200 L/ha of water at 206 – 300 kPa.</p> <p>Make only one application per season.</p>

	<p>If weeds are emerged at the time of NCS COMETA 480 Herbicide application, the addition of AGRAL 90 Non-ionic Wetting & Spreading Agent Liquid at the rate of 0.2% v/v is recommended. If weeds have not emerged, no adjuvant is recommended.</p> <p>Use of NCS COMETA 480 Herbicide on asparagus (established) may cause bleaching and may even lead to yield reduction.</p> <p>Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.</p>
--	--

Highbush Blueberry (Bearing and Non-Bearing Years)

CROP	Highbush Blueberry (Bearing and Non-Bearing Years)
WEEDS CONTROLLED	Lamb's-quarters, redroot pigweed, velvetleaf, wild mustard
WEEDS SUPPRESSED	Common ragweed
TIMING	Pre-emergent (up to 2-leaf weed stage) to weeds
PRODUCT RATE	0.30 L/ha
REMARKS	<p>Make only one application per year.</p> <p>Apply NCS COMETA 480 Herbicide up to 8 leaf stage to weeds and prebloom to crop.</p> <p>Apply broadcast (directed spray to the base of the plant) ground application.</p> <p>Apply in 100-200 L/ha water at 206-300 kPa.</p>

OR

WEEDS CONTROLLED	Eastern black nightshade, velvetleaf, redroot pigweed
WEEDS SUPPRESSED	Common ragweed
TIMING	Post-emergent (3 to 8 leaf weed stage) to weeds
PRODUCT RATE	0.30 L/ha
TANK MIX PARTNER RATE	0.2% v/v AGRAL 90 Non-ionic Wetting & Spreading Agent Liquid
REMARKS	<p>Make only one application per year.</p> <p>Apply NCS COMETA 480 Herbicide up to 8 leaf stage to weeds and prebloom to crop.</p> <p>Apply broadcast (directed spray to the base of the plant) ground application.</p> <p>Apply in 100-200 L/ha water at 206-300 kPa.</p>

Lowbush Blueberry (Bearing and Non-Bearing Years)

CROP	Lowbush Blueberry (Bearing and Non-Bearing Years)
WEEDS CONTROLLED	Lamb's-quarters, redroot pigweed, velvetleaf, wild mustard
WEEDS SUPPRESSED	Common ragweed
TIMING	Pre-emergent (up to 2-leaf weed stage) to weeds
PRODUCT RATE	0.30 L/ha
REMARKS	<p>Make only one application per year.</p> <p>Apply NCS COMETA 480 Herbicide up to 8 leaf stage to weeds.</p>

	<p>Do not harvest crop within 60 days of application.</p> <p>Apply broadcast (over the top of the blueberries) ground application.</p> <p>Apply in 100-200 L/ha water at 206-300 kPa.</p>
OR	
WEEDS CONTROLLED	Eastern black nightshade, velvetleaf, redroot pigweed
WEEDS SUPPRESSED	Common ragweed
TIMING	Post-emergent (3 to 8 leaf weed stage) to weeds
PRODUCT RATE	0.30 L/ha
TANK MIX PARTNER RATE	0.2% v/v AGRAL 90 Non-ionic Wetting & Spreading Agent Liquid
REMARKS	<p>Make only one application per year.</p> <p>Apply NCS COMETA 480 Herbicide up to 8 leaf stage to weeds.</p> <p>Do not harvest crop within 60 days of application.</p> <p>Apply broadcast (over the top of the blueberries) ground application.</p> <p>Apply in 100-200 L/ha water at 206-300 kPa.</p>

Sod (Production Only)

CROP	Sod (Production Only)
WEEDS CONTROLLED	Lamb's-quarters, redroot pigweed, velvetleaf, wild mustard
WEEDS SUPPRESSED	Common ragweed
TIMING	Pre-emergent (up to 2-leaf weed stage) to weeds
PRODUCT RATE	0.30 L/ha
REMARKS	<p>Make only one application per year.</p> <p>Apply NCS COMETA 480 Herbicide pre-emergence to weeds in established sod OR prior to, or post-seeding and prior to emergence of weeds OR postemergence to weeds (regardless of whether prior to, or post-seeding)</p> <p>Cool-season turf species and varieties may vary in their tolerance to herbicides, including NCS COMETA 480 Herbicide. Since not all cool season turf varieties have been tested for tolerance to NCS COMETA 480 Herbicide, first use of NCS COMETA 480 Herbicide should be limited to a small area of each variety or mixture to confirm tolerance prior to adoption as a general field practice. Additionally, consult your seed supplier for information on the tolerance of specific cool-season turf varieties to NCS COMETA 480 Herbicide. Do not apply NCS COMETA 480 Herbicide to creeping bentgrass.</p> <p>Apply in 250 L/ha of water.</p> <p>Make application close to anticipated weed seed germination. Apply at grass seeding or close to seeding for best performance. Weed control is most effective on young, actively growing weeds.</p>

	<p>Avoid spraying on newly germinated turfgrass plants. Efficacy will be reduced under moisture stress or from applications to mature weeds.</p> <p>NCS COMETA 480 Herbicide may reduce density of fine fescue seedings. NCS COMETA 480 Herbicide can be used on grass seed blends that contain less than 20% by weight of hard or fine fescue.</p> <p>Wait until the newly germinated turf has been mowed two times or four weeks after emergence (whichever is longer) before making a postemergence application.</p> <p>Use only ground boom equipment when applying this product.</p> <p>Do not apply NCS COMETA 480 Herbicide through any type of irrigation equipment.</p> <p>Do not apply directly to water or areas where surface water is present. Do not spray to runoff.</p>
OR	
WEEDS CONTROLLED	Eastern black nightshade, velvetleaf, redroot pigweed
WEEDS SUPPRESSED	Common ragweed
TIMING	Post-emergent (3 to 8 leaf weed stage) to weeds
PRODUCT RATE	0.21 L/ha
TANK MIX PARTNER RATE	0.2% v/v AGRAL 90 Non-ionic Wetting & Spreading Agent Liquid
REMARKS	<p>Make only one application per year.</p> <p>Apply NCS COMETA 480 Herbicide pre-emergence to weeds in established sod OR prior to, or post-seeding and prior to emergence of weeds OR postemergence to weeds (regardless of whether prior to, or post-seeding)</p> <p>Cool-season turf species and varieties may vary in their tolerance to herbicides, including NCS COMETA 480 Herbicide. Since not all cool-season turf varieties have been tested for tolerance to NCS COMETA 480 Herbicide, first use of NCS COMETA 480 Herbicide should be limited to a small area of each variety or mixture to confirm tolerance prior to adoption as a general field practice. Additionally, consult your seed supplier for information on the tolerance of specific cool-season turf varieties to NCS COMETA 480 Herbicide. Do not apply NCS COMETA 480 Herbicide to creeping bentgrass.</p> <p>Apply in 250 L/ha of water.</p> <p>Make application close to anticipated weed seed germination. Apply at grass seeding or close to seeding for best performance. Weed control is most effective on young, actively growing weeds.</p> <p>Avoid spraying on newly germinated turfgrass plants. Efficacy will be reduced under moisture stress or from applications to mature weeds.</p> <p>NCS COMETA 480 Herbicide may reduce density of fine fescue seedings. NCS COMETA 480 Herbicide can be used on grass seed blends that contain less than 20% by weight of hard or fine fescue.</p>

	<p>Wait until the newly germinated turf has been mowed two times or four weeks after emergence (whichever is longer) before making a postemergence application.</p> <p>Use only ground boom equipment when applying this product.</p> <p>Do not apply NCS COMETA 480 Herbicide through any type of irrigation equipment.</p> <p>Do not apply directly to water or areas where surface water is present. Do not spray to runoff.</p>
--	---

Flax (Eastern Canada Only)

CROP	Flax (Eastern Canada Only)
WEEDS CONTROLLED	Lamb's-quarters, redroot pigweed, velvetleaf, wild mustard
WEEDS SUPPRESSED	Common ragweed
TIMING	Pre-emergent to flax and weeds
PRODUCT RATE	0.30 L/ha
REMARKS	<p>Make only one application of NCS COMETA 480 Herbicide per year.</p> <p>Apply in 200 L water and a minimum pressure of 206 kPa.</p> <p>Apply by ground equipment only.</p> <p>Applications of NCS COMETA 480 Herbicide to emerged flax can result in severe crop injury.</p> <p>Do not graze treated immature crops or cut for forage or hay.</p>

Pearl Millet (Eastern Canada Only)

CROP	Pearl Millet (Eastern Canada Only)
WEEDS CONTROLLED	Lamb's-quarters, redroot pigweed, velvetleaf, wild mustard
WEEDS SUPPRESSED	Common ragweed
TIMING	Pre-plant
PRODUCT RATE	0.30 L/ha
REMARKS	<p>Make only one application per year. Apply in 100-200 L water/ha at 205-300 kPa.</p> <p>Do not apply to pearl millet grown in course textured soils.</p> <p>Do not apply more than 0.30 L/ha per crop <u>or</u> per year.</p> <p>Apply 7-14 days before planting pearl millet.</p> <p>Apply at up to the 2-leaf stage of weeds.</p> <p>Application to emerged pearl millet can result in severe crop injury.</p>

Sorghum (Eastern Canada Only)

CROP	Sorghum (Eastern Canada Only)
WEEDS CONTROLLED	Lamb's-quarters, redroot pigweed, velvetleaf, wild mustard

WEEDS SUPPRESSED	Common ragweed
TIMING	Pre-plant
PRODUCT RATE	0.30 L/ha
REMARKS	<p>Make only one application per year.</p> <p>Apply in 100-200 L water/ha at 205-300 kPa.</p> <p>Do not apply to sorghum grown in course textured soils nor to sudangrass, or sorghum-sudangrass hybrids.</p> <p>Do not apply more than 0.30 L/ha per crop or per year.</p> <p>Apply 7-14 days before planting sorghum.</p> <p>Apply at up to the 2-leaf stage of weeds.</p> <p>Application to emerged sorghum can result in severe crop injury.</p>
Rhubarb (Established)	
CROP	Rhubarb (Established)
WEEDS CONTROLLED	Lamb's-quarters, redroot pigweed, velvetleaf, wild mustard
WEEDS SUPPRESSED	Common ragweed
TIMING	Pre-emergent to crop, pre-emergent up to 2-leaf stage of weeds
PRODUCT RATE	0.30 L/ha
REMARKS	<p>Make only one application per year.</p> <p>Do not harvest within 42 days of application.</p> <p>Apply in 200 L/ha water at 206-300 kPa.</p> <p>CAUTION: Apply to dormant (prior to spring green-up) rhubarb. Applications of NCS COMETA 480 Herbicide to rhubarb that is not dormant will result in crop injury. Rainfall or irrigation after the NCS COMETA 480 Herbicide application may increase the risk of injury to emerging rhubarb. Repeated use of NCS COMETA 480 Herbicide in low Ph soils may cause injury to rhubarb plants. Wait at least 18 months after the last NCS COMETA 480 Herbicide application in rhubarb before replanting rhubarb.</p>

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, NCS COMETA 480 Herbicide is a Group 27 herbicide. Any weed population may contain or develop plants naturally resistant to NCS COMETA 480 Herbicide and other Group 27 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

Where possible, rotate the use of NCS COMETA 480 Herbicide or other Group 27 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.

Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.

Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.

Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.

Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.

For further information and to report suspected resistance, contact company representatives at 1-437-880-8283.

RECROPPING GUIDELINES

Certain crops have been shown to be sensitive to low residues of NCS COMETA 480 Herbicide in the soil. The minimum recropping interval is the time between the last application of NCS COMETA 480 Herbicide and the anticipated date of planting of the next crop.

To avoid injury to subsequent crops after an application at the recommended rate of NCS COMETA 480 Herbicide, the following recropping intervals should be observed.

CROP	MINIMUM RECROPPING INTERVAL (MONTHS)
Winter Wheat	3
Spring wheat	10
Field corn (grain or silage)	
Seed corn	
Sweet corn	
Alfalfa	11
Potato	
Soybean	
Transplanted tomato	
White beans	
All other crops	Bioassay

Observe the recropping guidelines of any product(s) to be tank mixed with NCS COMETA 480 Herbicide.

In the event of crop failure, only field corn (grain or silage), seed corn, or sweet corn can be planted as a salvage crop in the same fields in which the failed crop was treated with NCS COMETA 480 Herbicide. An application of NCS COMETA 480 Herbicide must NOT be made to the salvage crop.

Land treated with NCS COMETA 480 Herbicide CAN ONLY BE PLANTED TO A CROP NOT LISTED IN THE PREVIOUS TABLE if a field bioassay can be successfully performed. The bioassay must indicate normal growth with no yield reductions.

A bioassay may be conducted in any year following the year of application to assess the tolerance of the tested crop intended for planting in the year following the bioassay. Do not conduct a bioassay in the year of application. When conducting a field bioassay, it is very important to select a representative area(s) of the field previously treated with NCS COMETA 480 Herbicide to plant the test crop(s).

FIELD BIOASSAY

1. Representative Sample:

Ensure that soil parameters such as soil texture, depth of topsoil layer, soil pH and drainage of the test area selected are representative of the remainder of the field.

2. Sample Size:

The seeded area of each selected bioassay crop must be large enough to ensure that reliable results are obtained. The seedbed preparations and seeding of the bioassay crop(s) should be conducted the same way as when the entire field would be planted.

3. Other Residual Herbicides:

It is important that other herbicide products which are known to have residual activity were not applied to the field between the last application of NCS COMETA 480 Herbicide and the bioassay testing period. Avoid the use of other pesticides during the duration of the bioassay as they may damage the indicator crop(s).

4. Comparisons:

Ideally, an untreated check strip in a neighbouring field should be established and monitored for comparison since growing conditions can vary greatly from year to year and may result in erroneous results.

5. Assessment:

The site should be monitored regularly throughout the growing season. Watch for any damage to the crop such as thinning, yellowing or stunting. A yield sample should be taken and compared to an adjacent untreated field.

DO NOT ROTATE TO OTHER CROPS UNTIL THE BIOASSAY INDICATES NORMAL GROWTH WITH NO YIELD REDUCTIONS.

Failure to follow these recropping guidelines could result in injury to seeded crop(s).

ADHERE TO THESE PRODUCT LABELS FOR BROADLEAF WEEDS CONTROLLED, PRECAUTIONS, WARNINGS, RESTRICTIONS, TANK MIXING AND SPRAYING INSTRUCTIONS, ROTATIONAL CROPS AND DETAILED DIRECTIONS FOR USE REGARDING DOSAGE VARIATIONS ACCORDING TO SOIL TYPE.

SPRAYER CLEANUP:

To avoid subsequent injury to other crops, thoroughly clean application equipment immediately after spraying. Ensure that all traces of the product are removed. The following procedures are recommended:

1. Drain and flush tank, boom and all hoses for several minutes with clean water containing a household detergent. **DO NOT** clean the sprayer near desirable vegetation, wells or other water sources.

2. Fill the sprayer tank with clean water and add one litre of household ammonia (containing 3% ammonia) per 100 litres of water. Allow the solution to agitate for 15 minutes prior to flushing the solution through the boom and nozzles. Drain the system.
3. Remove the nozzles and screens and wash separately in a bucket containing the ammonia solution.
4. Thoroughly rinse the tank, hoses, booms, nozzles and screens with clean water for a minimum of 5 minutes to remove all traces of ammonia.
5. Dispose of all rinsings in accordance with provincial regulations.

All products listed are trademarks of their respective companies.