

2025-1761  
2025-05-16

|       |          |           |
|-------|----------|-----------|
| GROUP | <b>3</b> | FUNGICIDE |
|-------|----------|-----------|

# STARPRO<sup>®</sup> XC

FOLIAR FUNGICIDE

SUSPENSION

FOR CONTROL OR SUPPRESSION OF LISTED DISEASES IN WHEAT AND BARLEY

COMMERCIAL

REGISTRATION NUMBER 35516 PEST CONTROL PRODUCTS ACT

CAUTION – EYE IRRITANT

Active Ingredients

**Prothioconazole** 216 g/L

**Tebuconazole** 216 g/L

Contains bronopol at 0.045% as a preservative.

**READ THE LABEL AND LEAFLET BEFORE USING**

NET CONTENTS: 1 L to bulk

ALBAUGH, LLC  
1525 NE 36th Street  
Ankeny IA 50021  
United States  
1-800-247-8013

For medical or treatment information from exposure to this product, call 1-888-347-6732 (7 days/week, 24-hr).

For 24-hour chemical spill, leak, fire, exposure, or accident response information, call CHEMTREC toll free at 1-800-424-9300.

STARPRO XC Foliar Fungicide is a broad-spectrum systemic fungicide for the control or suppression of listed diseases on wheat, and barley.

**NOTICE TO USER:**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

**PRECAUTIONS:**

**KEEP OUT OF REACH OF CHILDREN.** May irritate eyes. Avoid contact with eyes. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, or smoking. If product comes in contact with clothing, remove all contaminated clothing, wash skin with soap and water and dress in clean clothing. Launder applicator clothing separate from other laundry. **DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

**PROTECTIVE CLOTHING AND EQUIPMENT:**

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and boots during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair. Coveralls and gloves are not required during application within closed cabs or cockpits.

**FIRST AID:**

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION:**

Treat symptomatically. This product contains petroleum distillates. Vomiting may cause aspiration pneumonia.

**ENVIRONMENTAL PRECAUTIONS:**

Toxic to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

#### **STORAGE:**

- Store this product away from food or feed.
- Do not contaminate water, food, or feed by storage or disposal.
- Keep away from fire or open flame or other sources of heat.
- Do not store at temperatures below freezing.
- If stored for 1 year or longer, shake well before using.
- Store the tightly closed container away from feeds, seeds, fertilizer, plants and foodstuffs.
- Do not use or store in or around the home.
- Keep in original container during storage.

#### **DISPOSAL:**

##### **Recyclable Container :**

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

##### **Returnable Container:**

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

**Refillable Container:** For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

##### **Disposal of Unused, Unwanted Product:**

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for cleanup of spills.

STARPRO is a registered trademark of Albaugh, LLC.

Booklet

|       |          |           |
|-------|----------|-----------|
| GROUP | <b>3</b> | FUNGICIDE |
|-------|----------|-----------|

# STARPRO<sup>®</sup> XC

FOLIAR FUNGICIDE

SUSPENSION

FOR CONTROL OR SUPPRESSION OF LISTED DISEASES IN WHEAT, AND BARLEY

COMMERCIAL

REGISTRATION NUMBER 35516 PEST CONTROL PRODUCTS ACT

CAUTION – EYE IRRITANT

Active Ingredients:

|                        |         |
|------------------------|---------|
| <b>Prothioconazole</b> | 216 g/L |
| <b>Tebuconazole</b>    | 216 g/L |

Contains bronopol at 0.045% as a preservative.

**READ THE LABEL AND LEAFLET BEFORE USING  
KEEP OUT OF REACH OF CHILDREN**

NET CONTENTS: 1 L to bulk

ALBAUGH, LLC  
1525 NE 36th Street  
Ankeny IA 50021  
United States  
1-800-247-8013

For medical or treatment information from exposure to this product, call 1-888-347-6732  
(7 days/week, 24-hr).

For 24-hour chemical spill, leak, fire, exposure, or accident response information, call  
CHEMTREC toll free at 1-800-424-9300.

## GENERAL INFORMATION

### NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

### PRODUCT DESCRIPTION

STARPRO XC Foliar Fungicide is a broad-spectrum systemic fungicide for the control or suppression of listed diseases on wheat, and barley.

## SAFETY AND HANDLING

### PRECAUTIONS:

**KEEP OUT OF REACH OF CHILDREN.** May irritate eyes. Avoid contact with eyes. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, or smoking. If product comes in contact with clothing, remove all contaminated clothing, wash skin with soap and water and dress in clean clothing. Launder applicator clothing separate from other laundry. **DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.. Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

### PROTECTIVE CLOTHING AND EQUIPMENT:

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and boots during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair. Coveralls and gloves are not required during application within closed cabs or cockpits..

### FIRST AID:

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person..

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

### TOXICOLOGICAL INFORMATION:

Treat symptomatically. This product contains petroleum distillates. Vomiting may cause aspiration pneumonia.

### **ENVIRONMENTAL PRECAUTIONS**

Toxic to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under **DIRECTIONS FOR USE**.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Tebuconazole is persistent and will carryover.

### **STORAGE:**

- Store this product away from food or feed.
- Do not contaminate water, food, or feed by storage or disposal.
- Keep away from fire or open flame or other sources of heat.
- Do not store at temperatures below freezing.
- If stored for 1 year or longer, shake well before using.
- Store the tightly closed container away from feeds, seeds, fertilizer, plants and foodstuffs.
- Do not use or store in or around the home.
- Keep in original container during storage.

### **DISPOSAL**

#### **Recyclable Container Disposal:**

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

#### **Returnable Container:**

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

#### **Refillable Container:**

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

**Disposal of Unused, Unwanted Product:**

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for clean-up of spills.

**Disposal of Unused Spray Solution:**

If any spray solution remains in the tank after spraying is finished, it should be sprayed on the perimeter of the area just sprayed, **NOT ON THE CROP**, away from water supplies, ditches, and irrigation canals. Spray buffer zones indicated in the **DIRECTIONS FOR USE** must also be respected.

**DIRECTIONS FOR USE**

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

**Introduction**

STARPRO XC Foliar Fungicide is a broad-spectrum systemic fungicide for control of a wide range of diseases on wheat (including spring, durum and winter), and barley. STARPRO XC Foliar Fungicide will protect the crop from yield and quality losses due to disease.

**CROP, DISEASE, RATE AND TIMING**

| <b>USE DIRECTIONS</b>                    |   |                             |  |
|--|---|-----------------------------|--|
| <b>CROP</b>                              | <b>DISEASE</b>  | <b>RATE<br/>(mL/ha)</b>     | <b>REMARKS</b>   |
| <b>Wheat</b><br>(winter, spring & durum) | <p><b>Suppression of:</b><br/>Fusarium Head Blight<br/>(<i>Fusarium</i> spp.)</p> <p><b>Control of:</b><br/>Rusts<br/>(<i>Puccinia recondita</i>,<br/><i>Puccinia graminis</i>,<br/><i>Puccinia striiformis</i>)</p> <p>Leaf &amp; Glume Blotch<br/>(<i>Septoria tritici</i>,<br/><i>Stagonospora nodorum</i>)</p> <p>Tan Spot<br/>(<i>Pyrenophora tritici-repentis</i>)</p> <p>Powdery Mildew<br/>(<i>Erysiphe graminis</i>)</p> | 455<br><br>(apply with NIS) | <p>For optimum suppression of Fusarium Head Blight, apply STARPRO XC Foliar Fungicide as a preventative spray within the time period from when at least 75% of the wheat heads on the main stem are fully emerged to when 50% of the heads on the main stem are in flower. Application at this timing will also control the listed leaf diseases.</p> <p>For optimal results, apply STARPRO XC Foliar Fungicide with a registered NIS surfactant at 0.125% v/v.</p> <p>Apply by ground or aerial application equipment. For ground application, apply specified dosage in a minimum of 100 L of water per hectare. For aerial application, apply specified dosage in a minimum of 50 L of water per hectare. A maximum of one application may be applied per year.</p> |

|               |   |                            |  |
|---------------|---|----------------------------|--|
| <b>Barley</b> | <p><b>Suppression of:</b><br/>Fusarium Head Blight<br/>(<i>Fusarium</i> spp.)</p> <p><b>Control of:</b><br/>Net Blotch<br/>(<i>Pyrenophora teres</i>)</p> <p>Spot Blotch<br/>(<i>Cochliobolus sativus</i>)</p> <p>Scald<br/>(<i>Rhynchosporium secalis</i>)</p> <p>Leaf Blotch<br/>(<i>Septoria passerinii</i>)</p> <p>Leaf, Stem and Stripe<br/>Rusts (<i>Puccinia hordei</i>,<br/><i>Puccinia graminis</i>, <i>Puccinia<br/>striiformis</i>)</p> <p>Powdery Mildew<br/>(<i>Erysiphe graminis</i>)</p> | 455<br>(apply with<br>NIS) | <p>For optimum suppression of Fusarium head blight, apply STARPRO XC Foliar Fungicide as a preventative spray within the time period when 70 to 100% of the barley main stem heads are fully emerged, to 3 days after full head emergence. Application at this timing will also control the listed leaf diseases.</p> <p>For optimal results, apply STARPRO XC Foliar Fungicide with a registered NIS surfactant at 0.125% v/v.</p> <p>Apply by ground or aerial application equipment. For ground application, apply specified dosage in a minimum of 100 L of water per hectare. For aerial application, apply specified dosage in a minimum of 50 L of water per hectare. A maximum of one application may be applied per year.</p> |
|---------------|---|----------------------------|--|

**TANK MIXING:**

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact ALBAUGH LLC at 1-800-247-8013 for information before applying any tank mix that is not specifically recommended on this label.

**RESISTANCE MANAGEMENT ADVISORY FOR CEREAL CROPS:**

Repeated application of standalone DMI Fungicides should not be used on the same crop in one season against risky pathogens such as cereal powdery mildew in areas of high disease pressure for that particular pathogen. Mixture products, tank-mixtures or alternation with fungicides having a different mode of action have been shown to protect against the development of resistant forms of disease.

**Use Limitations**

- A maximum of one application of STARPRO XC Foliar Fungicide may be applied per crop year.
- Applications may not be made within 36 days of harvest.
- Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with STARPRO XC Foliar Fungicide.
- Straw cut after harvest may be fed or used for bedding.
- Do not enter treated fields for 12 hours after application.

Field sprayer application: **DO NOT** apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. **DO NOT** apply with sprays finer than the American Society of Agricultural and Biological Engineers (ASABE) S572 (572.1 to 572.3) medium classification. Boom height must be 60 cm or less above the crop or ground.

Conventionally piloted aircraft application: **DO NOT** apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 km/h at flying height at the site of application. **DO NOT** apply with sprays finer than the American Society of Agricultural and Biological Engineers (ASABE) S572 (572.1 to 572.3) medium classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotorspan.

Apply only by aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate specified for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices or a Global Positioning System (GPS).

### **Use Precautions**

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

### **Product Specific Precautions**

Read and understand the entire label before opening this product. If you have questions, call the registrant at 1-800-247-8013 or obtain technical advice from the distributor or your provincial or territorial agricultural representative. Application of this specific product must meet and/or conform to the following:

Volume: Apply the specified rate in a minimum spray volume of 50 litres per hectare for conventionally piloted aircraft.

### **SPRAY BUFFER ZONES**

A spray buffer zone is NOT required for uses with hand-held application equipment permitted on this label.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands) sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

| Method of application | Crop/Site                              |             | Spray Buffer Zones (metres) Required for the Protection of: |                  |                                     |                  | Terrestrial Habitat: |
|-----------------------|--|-------------|---|------------------|-------------------------------------|------------------|----------------------|
|                       |  |             | Freshwater Habitat of Depths:                               |                  | Estuarine/Marine Habitat of Depths: |                  |                      |
|                       |  |             | Less than 1 m   | Greater than 1 m | Less than 1 m                       | Greater than 1 m |                      |
| Field sprayer         | Wheat (spring, winter, during), Barley |             | 1   | 1                | 1                                   | 1                | 1                    |
| Aerial                | Wheat (spring, winter, during), Barley | Fixed wing  | 4   | 1                | 1                                   | 1                | 15                   |
|                       |  | Rotary wing | 1   | 1                | 1                                   | 1                | 10                   |

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASABE) category indicated on the labels for those tank mix partners.

The spray buffer zones for application of this product with conventional application equipment can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Drift Mitigation portion of the Canada.ca website.

**Rotational Crops:**

A rotational plantback interval of 120 days must be observed for crops not listed on the label.

Treated areas may be replanted with any crop specified on this label as soon as practical after the last application STARPRO XC Foliar Fungicide.

**Pre-harvest Intervals**

| <u>Crop</u>                      | <u>PHI</u> |
|----------------------------------|------------|
| Barley                           | 36 days    |
| Wheat (spring, durum and winter) | 36 days    |

## **MIXING INSTRUCTIONS**

STARPRO XC Foliar Fungicide must be applied with properly calibrated, clean equipment. Prior to adding STARPRO XC Foliar Fungicide to the spray tank, ensure that the spray tank is thoroughly clean.

STARPRO XC Foliar Fungicide applied alone:

- Add one-half of the required amount of water to the spray or mixing tank and start agitation.
- Add the required quantity of STARPRO XC Foliar Fungicide to the water and complete filling with water to the required total volume.
- Maintain agitation throughout mixing and spraying.
- STARPRO XC Foliar Fungicide should be thoroughly dispersed prior to the addition of a non-ionic surfactant. STARPRO XC Foliar Fungicide is recommended to be used with a registered non-ionic surfactant, such as Agral 90 or AgSurf, at 0.125% vol/vol.

## **Application Precautions**

### **GROUND APPLICATION:**

1. Use well-maintained and calibrated conventional spray equipment, which provides adequate and uniform coverage.
2. Apply STARPRO XC Foliar Fungicide in a minimum of 100 L water/ha.
3. Spray screens should be no finer than 50 mesh.
4. Ensure that the by-pass line discharges at the bottom of the tank to minimize foaming.
5. Maintain pressure at no less than 275 kPa to ensure good foliage penetration and coverage.
6. Provincial spray buffer zones that are greater than the spray buffer zones indicated in the DIRECTIONS FOR USE, Section 9, should be respected.
7. If spray mixture remains in the tank overnight, or for long periods during the day, agitate thoroughly prior to application.

### **AERIAL APPLICATION-GENERAL INFORMATION:**

1. To ensure proper coverage and distribution and to minimize drift, check to see that swath width and droplet size are adequate, and that wind velocities are low.
2. Use mechanical flaggers only.

### **AERIAL APPLICATION- OPERATOR PRECAUTIONS:**

1. **DO NOT** allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.
2. It is desirable that the pilot has communication capabilities at each treatment site at the time of application.
3. The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.
4. All personnel on the job site must wash hands and face thoroughly before eating and drinking.

5. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.
6. Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-800-247-8013 or obtain technical advice from the distributor or your provincial agricultural representative.

### **RESISTANCE MANAGEMENT RECOMMENDATIONS**

For resistance management, please note that STARPRO XC Foliar Fungicide contains Group 3 fungicides. Any fungal population may contain individuals naturally resistant to STARPRO XC Foliar Fungicide and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

- Where possible, rotate the use of STARPRO XC Foliar Fungicide or other Group 3 fungicides with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that is effective on the target pathogen when such use is permitted.
- Do not apply more than the indicated maximum number of applications specified for each crop in the DIRECTIONS FOR USE.
- Fungicide use should be based on an integrated pest management (IPM) program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological, and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications.
- Monitor treated fungal populations for resistance development. Notify Albaugh LLC if reduced sensitivity of the pathogen to STARPRO XC Foliar Fungicide is suspected.
- If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product and switch to another fungicide with a different target site of action, if available.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information and to report suspected resistance, contact an Albaugh LLC representative at 1-800-247-8013.

STARPRO is a registered trademark of Albaugh, LLC.