

2023-3120
2025-06-24

GROUP	15	27	HERBICIDES
-------	----	----	------------

STOREN Herbicide

HERBICIDE

COMMERCIAL - AGRICULTURAL

SUSPENSION

A herbicide for use in field and seed corn.

FOR SALE AND USE IN EASTERN CANADA ONLY

ACTIVE INGREDIENTS:

Bicyclopyrone	8.91 g ai/L
Mesotrione.....	36.7 g ai/L
S-metolachlor (and R-enantiomer).....	322 g ai/L
Pyroxasulfone.....	17.9 g ai/L

Warning, contains the allergen sulfite

Contains 1,2-benzisothiazolin-3-one at 0.009%, 2-bromo-2-nitropropane-1,3-diol at 0.023%, 5-chloro-2-methyl-4-isothiazolin-3-one at 0.0012% and 2-methyl-4-isothiazolin-3-one at 0.0004% as preservatives.

**READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

**CAUTION - EYE IRRITANT
POTENTIAL SKIN SENSITIZER**

REGISTRATION NO.: **35533**
PEST CONTROL PRODUCTS ACT

NET CONTENTS: **1 L to Bulk**

Syngenta Canada Inc.
140 Research Lane, Research Park
Guelph, Ontario N1G 4Z3
Telephone: 1-877-964-3682

LABEL

1.0 NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

2.0 FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

3.0 TOXICOLOGICAL INFORMATION

There is no specific antidote. Treat symptomatically.

4.0 PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

May irritate eyes. Avoid contact with eyes. Potential skin sensitizer.

Avoid all drift or contact with other vegetation.

Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

Restricted-Entry Intervals (REI): DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 days to perform hand-set irrigation in corn. For all other postapplication activities, **DO NOT** enter or allow worker entry into treated areas during the

restricted-entry interval (REI) of 12 hours.

DO NOT use or store near heat or open flame. **DO NOT** apply beyond the field boundary.

DO NOT apply by air.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Syngenta Canada Inc. at 1-87-SYNGENTA/1-877-964-3682.

5.0 PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and chemical-resistant footwear during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab.

Wash the outside of the gloves before removing. Wash splashes from skin and eyes IMMEDIATELY with plenty of water.

If pesticide gets inside, remove clothing/PPE immediately. Wash yourself thoroughly and put on clean clothing.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water.

While using product, do not eat, drink or use tobacco (including smoking).

All users should wash hands and shower thoroughly with soap and water after using STOREN Herbicide before eating, drinking, chewing gum, using tobacco, applying cosmetics, or using the toilet.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product. Do not reuse them.

Wear freshly laundered clothes daily.

Do not wear contaminated shoes.

Wash clothing in detergent and hot water before reuse.

Store and wash all protective clothing separately from household laundry.

6.0 ENVIRONMENTAL PRECAUTIONS

Toxic to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE. Toxic to small wild mammals.

This product demonstrates the properties and characteristics associated with chemicals

detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

7.0 STORAGE

Store this product away from food or feed. Always store in original container with top tightly closed in a cool, dry place. Do not store STOREN Herbicide below $-10\text{ }^{\circ}\text{C}$. Storing STOREN Herbicide below $-10\text{ }^{\circ}\text{C}$ can negatively affect product quality. Do not use or store near heat or open flame.

8.0 DISPOSAL

Recyclable Containers

Disposal of container: DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

Returnable Containers

Disposal of container: DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For Refillable Bulk Containers

Disposal of container: DO NOT use this container for any other purpose. For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product.

Disposal of unused, unwanted product: For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for clean-up of spills.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (1-800-FASTMED)***

GROUP	15	27	HERBICIDES
-------	----	----	------------

STOREN Herbicide

HERBICIDE

COMMERCIAL - AGRICULTURAL

SUSPENSION

A herbicide for use in field and seed corn.

FOR SALE AND USE IN EASTERN CANADA ONLY

ACTIVE INGREDIENTS:

Bicyclopyrone	8.91 g ai/L
Mesotrione.....	36.7 g ai/L
S-metolachlor (and R-enantiomer).....	322 g ai/L
Pyroxasulfone.....	17.9 g ai/L

Warning, contains the allergen sulfite

Contains 1,2-benzisothiazolin-3-one at 0.009%, 2-bromo-2-nitropropane-1,3-diol at 0.023%, 5-chloro-2-methyl-4-isothiazolin-3-one at 0.0012% and 2-methyl-4-isothiazolin-3-one at 0.0004% as preservatives.

**READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

**CAUTION - EYE IRRITANT
POTENTIAL SKIN SENSITIZER**

REGISTRATION NO.: **35533**
PEST CONTROL PRODUCTS ACT

Syngenta Canada Inc.
140 Research Lane, Research Park
Guelph, Ontario N1G 4Z3
Telephone: 1-877-964-3682

PAMPHLET

TABLE OF CONTENTS

	Section Number
NOTICE TO USER	1.0
FIRST AID	2.0
TOXICOLOGICAL INFORMATION	3.0
PRECAUTIONS.....	4.0
PERSONAL PROTECTIVE EQUIPMENT (PPE)	5.0
ENVIRONMENTAL PRECAUTIONS	6.0
STORAGE	7.0
DISPOSAL	8.0
PRODUCT INFORMATION	9.0
DIRECTIONS FOR USE	10.0
Ground Application	10.1
Mixing Instructions.....	10.2
Application in Liquid Fertilizers	10.3
Equipment Specific Instructions	10.4
Spraying Instructions.....	10.5
Sprayer Cleanup.....	10.6
Buffer Zones	10.7
CROP USE DIRECTIONS	11.0
Field Corn	11.1
Seed Corn	11.2
Pre-harvest Intervals	11.3
Use Precautions	11.4
RE-CROPPING GUIDELINES	12.0
Field Bioassay	12.1
RESISTANCE MANAGEMENT RECOMMENDATIONS	13.0

1.0 NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

2.0 FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

3.0 TOXICOLOGICAL INFORMATION

There is no specific antidote. Treat symptomatically.

4.0 PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

May irritate eyes. Avoid contact with eyes. Potential skin sensitizer.

Avoid all drift or contact with other vegetation.

Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

Restricted-Entry Intervals (REI): **DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 days to perform hand-set irrigation in corn. For all other postapplication activities, **DO NOT** enter or allow worker entry into treated areas during the

restricted-entry interval (REI) of 12 hours.

DO NOT use or store near heat or open flame. **DO NOT** apply beyond the field boundary.

DO NOT apply by air.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Syngenta Canada Inc. at 1-87-SYNGENTA/1-877-964-3682.

5.0 PERSONAL PROTECTIVE EQUIPMENT (PPE)

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and chemical-resistant footwear during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab.

Wash the outside of the gloves before removing. Wash splashes from skin and eyes IMMEDIATELY with plenty of water.

If pesticide gets inside, remove clothing/PPE immediately. Wash yourself thoroughly and put on clean clothing.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water.

While using product, do not eat, drink or use tobacco (including smoking).

All users should wash hands and shower thoroughly with soap and water after using STOREN Herbicide before eating, drinking, chewing gum, using tobacco, applying cosmetics, or using the toilet.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product. Do not reuse them.

Wear freshly laundered clothes daily.

Do not wear contaminated shoes.

Wash clothing in detergent and hot water before reuse.

Store and wash all protective clothing separately from household laundry.

6.0 ENVIRONMENTAL PRECAUTIONS

Toxic to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE. Toxic to small wild mammals.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

7.0 STORAGE

Store this product away from food or feed. Always store in original container with top tightly closed in a cool, dry place. Do not store STOREN HERBICIDE below $-10\text{ }^{\circ}\text{C}$. Storing STOREN HERBICIDE below $-10\text{ }^{\circ}\text{C}$ can negatively affect product quality. Do not use or store near heat or open flame.

8.0 DISPOSAL

Recyclable Containers

Disposal of container: DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

Returnable Containers

Disposal of container: DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For Refillable Bulk Containers

Disposal of container: DO NOT use this container for any other purpose. For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product.

Disposal of unused, unwanted product: For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for clean up of spills.

**IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING,
CALL 1-800-327-8633 (1-800-FASTMED)**

9.0 PRODUCT INFORMATION

STOREN Herbicide is labeled for use in field corn and seed corn for control of many annual grass and broadleaf weeds. STOREN HERBICIDE is a combination of the herbicides bicyclopyrone, mesotrione, pyroxasulfone and S-metolachlor.

NOTE: STOREN HERBICIDE contains benoxacor that has been shown to enhance S-metolachlor metabolism in corn.

10.0 DIRECTIONS FOR USE

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Syngenta Canada Inc. at 1-87-SYNGENTA / 1-877-964-3682 for information before applying any tank mix that is not specifically recommended on this label.

Syngenta Canada Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) of all potential tank mixes under all environmental conditions or for all crop varieties. Tank mixes that are not specifically listed on this label should be tested on a small area first, under local conditions and using standard practices, to confirm the tank mix is suitable for widespread application.

To determine the physical compatibility of this product with other products, use a jar test.

Always read and follow label directions including WALES mixing order.

10.1. Ground Application

Field sprayer application: **DO NOT** apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. **DO NOT** apply with sprays finer than the

American Society of Agricultural and Biological Engineers (ASABE) 5272 (S572.1 to 572.3) medium classification. Boom height must be 60 cm or less above the crop or ground.

DO NOT apply by air.

DO NOT apply more than 3.92 L product/ha/season.

DO NOT apply any other mesotrione containing products to fields treated with STOREN HERBICIDE in the same year.

On coarse textured soils with less than 3% OM, preplant and preemergence applications of STOREN Herbicide may cause injury to corn under adverse environmental conditions.

10.2. Mixing Instructions

1. Ensure that the sprayer interior is clean, then fill the spray tank with $\frac{1}{2}$ the required amount of water and engage gentle agitation. Good agitation is indicated by a rippling or rolling action on the surface of the water.
2. Add any WG or DF formulation mix partners and agitate to ensure complete mixing.
3. Add STOREN HERBICIDE and any other SE or SC formulation and agitate to ensure complete mixing.
4. Add any EC formulation mix partners and agitate to ensure complete mixing.
5. Fill the tank to $\frac{3}{4}$ the required amount of water.
6. Add any solution (SN or SL) formulation mix partners and agitate to ensure complete mixing.
7. Finish filling the sprayer with water, maintaining good agitation.
8. Always ensure that agitation is maintained until spraying is completed. After any break in spraying operations, agitate thoroughly before spraying again. Reduce agitation once tank volume is reduced to about one third of capacity.
9. STOREN Herbicide must be sprayed the same day as mixing. Do not allow spray solution to remain in the tank overnight.
10. Do not mix, load or clean spray equipment where there is a potential to contaminate wells or aquatic systems.

When using chemical handling equipment to fill the sprayer, the following additional recommendations apply:

- WG and DF formulations are preferentially batch mixed.
- SC, SN, and SL formulations may be inducted or batch mixed.
- EC formulations are preferentially batch mixed.

10.3. Application in Liquid Fertilizers

Nitrogen solutions (such as 28-0-0 UAN), excluding suspension and sulphur containing fertilizers, may replace water as a carrier for pre-emergence applications of STOREN HERBICIDE in corn. Do not use nitrogen solutions as a carrier to corn that has emerged. Always predetermine the compatibility of STOREN HERBICIDE tank mixes with your liquid fertilizer carrier by mixing small proportional quantities in advance. Even if STOREN HERBICIDE is physically compatible with a fluid fertilizer, constant agitation is necessary to maintain a uniform mixture during application. DO NOT apply post emergent UAN or liquid fertilizer or severe crop injury will occur.

10.4. Spraying Instructions

1. Carrier Volume: Apply in a minimum spray volume of 150 L/ha as per the tabulated instructions below.
2. Sprayer Agitation: Use a jet agitator or liquid sparge tube which recirculates 7-10% of the tank per minute. DO NOT use an air sparger.
3. Pump: Screens should be used to protect the pump and prevent clogging. Use 16 mesh or coarser screens on the suction side of the pump. DO NOT place a screen in the recirculation line. Use 50 mesh or coarser screens between the pump and boom unless directed otherwise by the sprayer manufacturer.
4. Spray Nozzles: 80° or 110° drift reducing flat fan (e.g. those with a pre-orifice or turbulence chamber) or air induction nozzles are recommended. Use 50 mesh or coarser nozzle screens or as recommended by the nozzle manufacturer. DO NOT use flood type nozzles, controlled droplet application equipment, spray foils or hollow cone nozzles.
5. Pressure: As recommended by the nozzle manufacturer to achieve no finer than ASABE medium sized droplets.
6. Apply at uniform speed and avoid overlapping. Avoid operations that contribute to non-uniform application and the potential for crop injury.

Coarse sprays are less likely to drift, therefore, avoid combinations of pressure and nozzle type that will result in fine particles (mist). **DO NOT** spray when the wind is blowing towards a nearby sensitive crop, garden, terrestrial habitat (such as shelter-belt) or aquatic habitat.

10.5. Sprayer Cleanup

To avoid subsequent injury to other crops, thoroughly clean application equipment immediately after spraying each day. Ensure that all traces of the product are removed. Do not allow residue to dry in spray tank. Do not allow solution to remain in lines or tank overnight. The following procedures are recommended:

1. Drain and flush tank, boom and all hoses for several minutes with clean water. Ensure dead end lines are flushed. **Do not** clean the sprayer near desirable vegetation, wells or other water sources.
2. Prepare a cleaning solution of one part of household ammonia per 25 parts of water. A household detergent may also be added to the solution.
3. Use a pressure washer to clean the inside of the spray tank with this solution. Take care to wash all parts of the tank, including the inside top surface. If a pressure washer is not

available, completely fill the sprayer with the cleaning solution to ensure contact of the cleaning solution with all internal surfaces of the tank and plumbing. Start agitation in the sprayer and thoroughly recirculate the cleaning solution for at least 15 minutes. All visible deposits must be removed from the spraying system.

4. Flush hoses, spray lines, and nozzles for at least 1 minute with the cleaning solution.
5. Dispose of all rinsings in accordance with provincial regulations.
6. Repeat steps 2-5.
7. Remove nozzles, screens, and strainers and clean separately in the ammonia solution after completing the above procedures.
8. Rinse the complete spraying system with clean water.

10.6. Spray Buffer Zones

A spray buffer zone is NOT required for:

- low-clearance hooded or shielded sprayers that prevent spray contact with crop or foliage

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop	Spray Buffer Zones (metres) Required for the Protection of:				
		Freshwater Habitat of Depths:		Estuarine/Marine Habitats of Depths:		Terrestrial habitat
		Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer	Corn	2	1	1	1	5

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASABE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product for conventional application equipment can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Drift Mitigation portion of the Canada.ca website.

11.0 CROP USE DIRECTIONS

11.1. Field Corn

Field corn - Surface pre-plant¹, pre-emergent and post-emergent^{2,6}	
PRODUCT RATE	3.92 L/ha
WEEDS	<p>Broadleaf weeds controlled (including triazine, group 2 and glyphosate-tolerant biotypes): American nightshade³, Eastern black nightshade, lamb's-quarters, redroot pigweed, velvetleaf, wild mustard, common ragweed, giant ragweed (suppression) (1 – 10 cm), Canada fleabane (1 – 10 cm), common waterhemp, hairy galinsoga (0 – 10 cm) and palmer amaranth</p> <p>Grass weeds controlled (up to 2 leaf): Barnyard grass, crabgrass (smooth, hairy), fall panicum, foxtail (green, yellow, giant), witchgrass, annual bluegrass and yellow nutsedge (suppression) (1 – 15 cm)</p> <p>Grass weeds suppressed⁴: proso millet</p>
PRODUCT RATE	3.35 L/ha ⁵
WEEDS	<p>Short season suppression of: Broadleaf weeds: lamb's-quarters, redroot pigweed, common ragweed, velvetleaf, Eastern black nightshade, American nightshade, wild mustard, common waterhemp, Canada fleabane (1 – 10 cm) and hairy galinsoga (0 – 10 cm)</p> <p>Grass weeds (up to 2 leaf): Barnyard grass, crabgrass (smooth, hairy), fall panicum, foxtails (green, yellow, giant), proso millet, annual bluegrass and witchgrass.</p>

Field corn – Tank mixes - Surface pre-plant¹, pre-emergent and post-emergent²	
WEEDS	Control or suppression of annual weeds listed above and burn-down of emerged annual and perennial weeds
TANK MIX PARTNER RATE/HA	900 g ae/ha glyphosate containing products that are registered for burndown of annual and perennial weeds ^{7,8}
WEEDS	Control or suppression of annual weeds listed above and control of wild buckwheat and lady's thumb
TANK MIX PARTNER	0.6-1.2 L/ha AATREX® Liquid 480 Herbicide; Use the 1.2 L/ha rate for control of lady's thumb.
NOTES	<p>Ground application only</p> <p>For pre-emergent uses, apply in a minimum of 150 L of carrier (water or UAN) per hectare. For post-emergent uses, apply in a minimum of 150 L of water per hectare.</p>

¹ STOREN Herbicide may be applied to the soil surface up to 7 days before planting for minimum and no-till systems in corn. If possible, do not move the treated soil out of the row or move untreated soil to the surface during planting or weed control will be diminished.

² Apply post up to and including 4 leaf corn.

³ Pre-emergent treatments sometimes provide better control than pre-plant surface or post-emergent treatments.

⁴ Apply prior to proso millet emergence and before other grass weeds exceed the 2-leaf stage.

⁵ For surface preplant and pre-emergent timings, the 3.35 L/ha rate will provide suppression only through the critical crop establishment phase and is to be used in a planned weed management program with a post-emergent herbicide treatment. Monitor the species and growth stages of escapes in order to better select a post-emergent herbicide application.

⁶ Application must be made prior to weed emergence or if weeds have emerged a tank mix with

glyphosate containing products in glyphosate tolerant corn is recommended.

⁷ Post-emergent applications on Glyphosate tolerant corn only. Residual control or suppression of annual weeds and burn-down of emerged annual and perennial weeds.

⁸ Higher rates of glyphosate may be recommended in some situations. Refer to the glyphosate product label for further effective rates.

11.2. Seed Corn

Seed Corn	
APPLICATION TIMING	Pre-emergent only
WEEDS	Control or suppression of annual weeds listed above
PRODUCT RATE	3.35 or 3.92 L/ha
NOTES	Ground application only For pre-emergent uses, apply in a minimum of 150 L of carrier (water or UAN) per hectare.

11.3. Pre-Harvest Intervals (PHIs)

Do not harvest treated crop prior to the number of days specified in the table below.

Crops	PHI (days)
Field corn, forage	45
Field corn grain and stover	Harvest at maturity

11.4. Use Precautions

STOREN Herbicide may be used pre-plant, pre-emergent or post-emergent on field corn and pre-emergent in production seed corn. Not all seed corn inbreds varieties have been tested; the use of STOREN HERBICIDE must be approved by the seed company and comply with the directions given by the seed corn company.

Although STOREN Herbicide has a flexible recropping profile, certain crops may be sensitive to low concentrations in the soil. Therefore, careful consideration should be given to crop rotation plans prior to using STOREN Herbicide (refer to "*RECROPPING GUIDELINES*").

Dry conditions following application may reduce the pre-emergent activity of STOREN Herbicide. If an activating rain (>1.25 cm) is not received within 7-10 days after a pre-emergent application, cultivate or use other weed control measures if weeds develop.

Do not apply to soils that contain less than 1% or more than 10% organic matter.

Temporary crop injury (bleaching & stunting) may occur under extreme weather conditions or when the crop is under stress due to inadequate or abnormally high moisture levels or extreme temperatures as well as under conditions of insect or plant disease attack, carryover of pesticide residues, use of certain soil applied systemic insecticides, improperly placed fertilizers or soil insecticides. The symptoms are most visible where excessive rates have been applied, such as sprayer overlaps. Corn quickly outgrows these effects and continues to grow normally.

Do not make a foliar post-emergent application of any organophosphate or carbamate insecticide within 7 days before or 7 days after a STOREN HERBICIDE application or severe

corn injury may occur. Syngenta Canada Inc. will not be held responsible for losses or damage resulting from such use.

STOREN Herbicide may be applied prior to or following a pyrethroid-type insecticide such as FORCE® Insecticide or MATADOR® 120EC Insecticide.

Do not apply STOREN HERBICIDE through any type of irrigation system.

Protect sprayer operators from drift or mist. Additional information on spray drift management for GROUND APPLICATIONS is provided in the “Ground Application” section of DIRECTIONS FOR USE. When low volumes of spray are applied, complete coverage and thorough application are essential for most effective results. Schedule applications in accordance with local conditions. Consult your local agricultural authorities for specific use information.

12.0 RECROPPING GUIDELINES

Certain crops have been shown to be sensitive to low residues of STOREN HERBICIDE in the soil. The minimum recropping interval is the time between the last application of STOREN HERBICIDE and the anticipated date of planting of the next crop.

To avoid injury to subsequent crops after an application at the recommended rate of STOREN Herbicide, the following recropping intervals should be observed.

CROP	MINIMAL RECROPPING INTERVAL
Field Corn, Production Seed Corn and Silage Corn	Immediately (no restriction)
Winter Wheat	4.5 months
Spring Wheat	10 months
Soybeans (except on sand or loamy sand soils)*	11 months
Potato	11 months
All Other Crops	Bioassay (after 12 months)
ROTATIONAL CROPS USE PRECAUTIONS	
*Crop injury may be observed when rotating to soybean in coarse soils (sand, loamy sand) with a soil pH less than 6.5 and if adequate rainfall is not received in the year of application. The addition of atrazine tank mixed with STOREN herbicide may increase the severity of crop injury observed in the rotational crop.	

Observe the recropping guidelines of any product to be tank mixed with STOREN HERBICIDE.

WARNING:

Do not plant any crop other than corn or winter wheat (after 4.5 months) in the same year on land treated with STOREN HERBICIDE as injury may occur. In the event of crop failure, only field corn (grain or silage) or seed corn can be planted as a salvage crop in the same fields in which the failed crop was treated with STOREN Herbicide. An application of STOREN Herbicide must NOT be made to the salvage crop.

Crops that may be planted following STOREN HERBICIDE are provided in the recropping guidelines. However, when the rotational crop is subjected to stress conditions, e.g. abnormally hot, dry weather, preceded by extended periods of dry weather the previous season, injury may

occur. Land treated with STOREN Herbicide CAN ONLY BE PLANTED TO A CROP NOT LISTED IN THE PRECEEDING TABLE if a field bioassay can be successfully performed. The bioassay must indicate normal growth with no yield reductions.

A bioassay may be conducted in any year following the year of application to assess the tolerance of the tested crop intended for planting in the year following the bioassay. Do not conduct a bioassay in the year of application. When conducting a field bioassay, it is very important to select a representative area(s) of the field previously treated with STOREN Herbicide to plant the test crop(s).

12.1. Field Bioassay

1. **Representative Sample:**
Ensure that soil parameters such as soil texture, depth of topsoil layer, soil pH and drainage of the test area selected are representative of the remainder of the field.
2. **Sample Size:**
The seeded area of each selected bioassay crop must be large enough to ensure that reliable results are obtained. The seedbed preparations and seeding of the bioassay crop(s) should be conducted the same way as when the entire field would be planted.
3. **Other Residual Herbicides:**
It is important that other herbicide products which are known to have residual activity were not applied to the field between the last application of STOREN HERBICIDE and the bioassay testing period. Avoid the use of other pesticides during the duration of the bioassay as they may damage the indicator crop(s).
4. **Comparisons:**
Ideally, an untreated check strip in a neighbouring field should be established and monitored for comparison since growing conditions can vary greatly from year to year and may result in erroneous results.
5. **Assessment:**
The site should be monitored regularly throughout the growing season. Watch for any damage to the crop such as thinning, yellowing or stunting. A yield sample should be taken and compared to an adjacent untreated field.

DO NOT ROTATE TO OTHER CROPS UNTIL THE BIOASSAY INDICATES NORMAL GROWTH WITH NO YIELD REDUCTIONS.

Failure to follow these recropping guidelines could result in injury to seeded crop(s).

ADHERE TO THESE PRODUCT LABELS FOR BROADLEAF AND GRASSWEEDS CONTROLLED, PRECAUTIONS, WARNINGS, RESTRICTIONS, TANK MIXING AND SPRAYING INSTRUCTIONS, ROTATIONAL CROPS AND DETAILED DIRECTIONS FOR USE REGARDING DOSAGE VARIATIONS ACCORDING TO SOIL TYPE.

13.0 RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, STOREN HERBICIDE is a Group 15 and 27 herbicide. Any weed population may contain or develop plants naturally resistant to STOREN HERBICIDE and other Group 15 and/or 27 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

Where possible, rotate the use of STOREN HERBICIDE or other Group 15 and/or 27 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.

Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.

Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.

Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.

Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.

For further information and to report suspected resistance, contact company representatives at 1-877-SYNGENTA (1-877-964-3682) or at www.syngenta.ca.

AATREX[®], FORCE[®] and MATADOR[®] are trademarks of a Syngenta Group Company.