

2025-1975  
2025-06-19

Container Label

**Maxunitech Glufosinate 200 SN**

GROUP	<b>10</b>	HERBICIDE
-------	-----------	-----------

**FOR SALE FOR USE ON GLUFOSINATE AMMONIUM TOLERANT FIELD CORN AND ON GLUFOSINATE AMMONIUM TOLERANT SOYBEANS IN CANADA AND ON GLUFOSINATE AMMONIUM TOLERANT CANOLA IN EASTERN CANADA AND BRITISH COLUMBIA**

**SOLUTION  
COMMERCIAL**

**ACTIVE INGREDIENT: glufosinate-ammonium..... 200 g/L**

**REGISTRATION NUMBER: 35549 PEST CONTROL PRODUCTS ACT**

**READ THE LABEL AND ATTACHED BOOKLET BEFORE USING**



**POISON**

**WARNING: EYE and SKIN IRRITANT**

**NET CONTENTS: 1L - 1,000 L**

**Maxunitech North America, Inc.**  
11601 Shadow Creek Pkwy, Suite 111-573  
Pearland, Texas 77584, USA  
86-571-28007837 or 1-855-462-9621

**Canadian Agent:**  
John Gibson  
618 Wilkins Terrace Saskatoon, SK S7W 0C8  
1-306-514-0770

**EMERGENCY TELEPHONE NUMBERS:**

For product and use information, Maxunitech North America Inc., 86-571-28007837 or 1-855-462-9621.

For spills or transportation accidents, Chemtrec, 1-800-424-9300.

## **PRECAUTIONS:**

### **KEEP OUT OF REACH OF CHILDREN.**

Causes eye irritation. May irritate the skin. Do not get in eyes, skin or clothing. May be harmful if swallowed or absorbed through the skin. Wash thoroughly after using and before eating, drinking or smoking.

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing and loading. Gloves are not required during application within a closed cab.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

When tank-mixes are permitted, read and observe all label directions, including rates and restrictions for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.

Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact Maxunitech North America, Inc. at 1-855-462-9621 or 86-571-28007837..

## **FIRST AID:**

**Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.**

If swallowed, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15– 20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

## **TOXICOLOGICAL INFORMATION:**

**Note to Physician:** If this product is ingested, endotracheal intubation and gastric lavage should be performed as soon as possible, followed by charcoal and sodium sulfate administration.

## **ENVIRONMENTAL PRECAUTIONS:**

TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE To reduce runoff from treated areas into aquatic

habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

#### **STORAGE:**

Store this product away from food or feed.

**CANNOT** be stored below freezing. If stored for 1 year or longer, shake well before using. Store the tightly closed container away from feeds, seeds, fertilizers, plants and foodstuffs. Do not use or store in or around the home. Keep in original container during storage.

#### **DISPOSAL:**

**Recyclable Container Disposal:** DO NOT reuse this container for any purpose. This is a recyclable container and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site.

Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

**Returnable Container Disposal:** DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

**Refillable containers (containers that can be refilled for the user by the distributor/dealer) Disposal:** For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

**Disposal of Unused, Unwanted Product:** For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for cleanup of spills.

#### **NOTICE TO USER:**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

Booklet

**Maxunitech Glufosinate 200 SN**

<b>GROUP</b>	<b>10</b>	<b>HERBICIDE</b>
--------------	-----------	------------------

**FOR SALE FOR USE ON GLUFOSINATE AMMONIUM TOLERANT FIELD CORN AND ON GLUFOSINATE AMMONIUM TOLERANT SOYBEANS IN CANADA AND ON GLUFOSINATE AMMONIUM TOLERANT CANOLA IN EASTERN CANADA AND BRITISH COLUMBIA**

**SOLUTION  
COMMERCIAL**

**ACTIVE INGREDIENT: glufosinate-ammonium..... 200 g/L**

**REGISTRATION NUMBER: 35549 PEST CONTROL PRODUCTS ACT**

**READ THE LABEL AND THIS BOOKLET BEFORE USING**



**POISON**

**WARNING: EYE and SKIN IRRITANT**

**Maxunitech North America, Inc.**  
11601 Shadow Creek Pkwy, Suite 111-573  
Pearland, Texas 77584, USA  
86-571-28007837 or 1-855-462-9621

**Canadian Agent:**  
John Gibson  
618 Wilkins Terrace Saskatoon, SK S7W 0C8  
1-306-514-0770

**EMERGENCY TELEPHONE NUMBERS:**

For product and use information, Maxunitech North America Inc., 86-571-28007837 or 1-855-462-9621.

For spills or transportation accidents, Chemtrec, 1-800-424-9300.

**Maxunitech Glufosinate  
200 SN**

<i>Table of Contents:</i>		<i>Section Number:</i>
<b>GENERAL INFORMATION</b>	The Product	1
<b>SAFETY AND HANDLING</b>	Precautions, Protective Clothing and Equipment, and Re-entry Restriction	2
	First Aid and Toxicological Information	3
	Environmental Precautions	4
	Storage	5
	Disposal	6
<b>DIRECTIONS FOR USE</b>	Crops, Weeds, Rates, Timing	7
	Tank Mixes	8
	Resistance Management	
	Recommendations	9
	Application Instructions and Cautions	10
	Mixing Instructions	11
	Sprayer Cleanup	12
	Livestock Feeding and Preharvest Intervals	13
	Notices	14

## GENERAL INFORMATION

### Section 1: The Product

Maxunitech Glufosinate 200 SN provides control of a broad spectrum of grassy and broadleaf weeds in Canola varieties, Corn hybrids and Soybean varieties that are specially developed to be tolerant to glufosinate ammonium.

Maxunitech Glufosinate 200 SN may also be applied to glufosinate ammonium tolerant inbred Corn lines grown for seed production.

For Canola hybrid seed production, Maxunitech Glufosinate 200 SN may be used to remove segregating wild type plants within the female population. See Section 7 (Crops, Weeds, Rates, Timing - Canola Hybrid Seed Production).

## SAFETY AND HANDLING

### Section 2: Precautions, Protective Clothing and Equipment, and, Re-entry Restrictions

#### PRECAUTIONS:

#### KEEP OUT OF REACH OF CHILDREN.

Causes eye irritation. May irritate the skin. Do not get in eyes, skin or clothing. May be harmful if swallowed or absorbed through the skin.

Wash thoroughly after using and before eating, drinking or smoking.

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing and loading. Gloves are not required during application within a closed cab.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

When tank-mixes are permitted, read and observe all label directions, including rates and restrictions for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.

Apply only to agricultural crops when the potential for drift to areas of human habitation and human activity, such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

### Section 3: First Aid and Toxicological Information

#### FIRST AID:

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**If swallowed**, call a poison control centre or doctor immediately for treatment advice. Have

person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**If in eyes**, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**If on skin or clothing**, take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled**, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

#### **TOXICOLOGICAL INFORMATION:**

**Note to Physician:** If this product is ingested, endotracheal intubation and gastric lavage should be performed as soon as possible, followed by charcoal and sodium sulfate administration.

### **Section 4: Environmental Precautions**

TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

Avoid spray drift onto susceptible plants and USE ONLY FOR RECOMMENDED PURPOSES AND AT RECOMMENDED RATES.

### **Section 5: Storage**

Store this product away from food or feed.

**CANNOT** be stored below freezing.

If stored for 1 year or longer, shake well before using.

Store the tightly closed container away from feeds, seeds, fertilizers, plants and foodstuffs.

Do not use or store in or around the home.

Keep in original container during storage.

### **Section 6: Disposal**

#### **Recyclable Container Disposal:**

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in

the tank.

2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

**Returnable Container Disposal:** DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

**Refillable Container (containers that can be refilled for the user by the distributor/dealer) Disposal:** For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

**Disposal of Unused, Unwanted Product:**

For information on disposal of unused, unwanted product, contact the manufacturer or the registrant or the provincial or territorial regulatory agency. Contact the registrant or the provincial or territorial regulatory agency. in case of a spill, and for cleanup of spills.

## DIRECTIONS FOR USE

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

### Section 7: Crops, Weeds, Rates, Timing

Best control will be obtained when Maxunitech Glufosinate 200 SN is applied at the recommended leaf stages of the weeds.

In Canola grown in Eastern Canada and British Columbia, apply Maxunitech Glufosinate 200 SN from the cotyledon stage up until, but prior to, the bolting stage of Canola.

In Corn grown in Canada, apply Maxunitech Glufosinate 200 SN from the 1-8 leaf stage of the

Corn plant or 5-6 visible collars (the leaf is counted once the next leaf is visible in the whorl).

In Soybeans grown in Canada, apply Maxunitech Glufosinate 200 SN from the cotyledon to the flowering stage of the crop.

Crop	Weed	Recommended Time (Leaf Stage)								Rate		
		1	2	3	4	5	6	7	8			
CANOLA, CORN, and SOYBEANS tolerant to Maxunitech Glufosinate 200 SN	Crabgrass	-----	-----	-----	-----	-----	-----	-----	-----	x	1.5 L/ha	
	Giant Foxtail	-----	-----	-----	-----	-----	-----	-----	-----	x		
	Green Foxtail	-----	-----	-----	-----	-----	-----	-----	-----	x		
	Proso Millet	-----	-----	-----	-----	-----	-----	-----	-----	x		
	Chickweed	-----	-----	-----	-----	-----	-----	-----	-----	-----		x
	Cocklebur	-----	-----	-----	-----	-----	-----	-----	-----	x		
	Ragweed	-----	-----	-----	-----	-----	-----	-----	-----	x		
	Redroot Pigweed	-----	-----	-----	-----	-----	-----	-----	-----	x		
	Shepherd's-purse	-----	-----	-----	-----	-----	-----	-----	-----	x		

Crop	Weed	Recommended Time (Leaf Stage)								Rate		
		1	2	3	4	5	6	7	8			
CANOLA, CORN, and SOYBEANS tolerant to Maxunitech Glufosinate 200 SN	Barnyard Grass	-----	-----	-----	-----	-----	-----	-----	-----	x	2 L/ha	
	Bristly Foxtail	-----	-----	-----	-----	-----	-----	-----	-----	x		
	Fall Panicum	-----	-----	-----	-----	-----	-----	-----	-----	x		
	Old Witchgrass	-----	-----	-----	-----	-----	-----	-----	-----	x		
	Wild Oats	-----	-----	-----	-----	-----	-----	-----	-----	x		
	Yellow Foxtail	-----	-----	-----	-----	-----	-----	-----	-----	x		
	Eastern Black Nightshade	-----	-----	-----	-----	-----	-----	-----	-----	x		
	Lady's-thumb	-----	-----	-----	-----	-----	-----	-----	-----	x		
	Lamb's-quarters	-----	-----	-----	-----	-----	-----	-----	-----	x		
	Perrenial Sowthistle	-----	-----	-----	-----	-----	-----	-----	-----	x		
	Ragweed	-----	-----	-----	-----	-----	-----	-----	-----	x		
	Stinkweed	-----	-----	-----	-----	-----	-----	-----	-----	-----		x
	Velvetleaf	-----	-----	-----	-----	-----	-----	-----	-----	x		
	Wild Buckwheat	-----	-----	-----	-----	-----	-----	-----	-----	x		
Wild Mustard	-----	-----	-----	-----	-----	-----	-----	-----	x			

**Canada Thistle and Field Bindweed:** For season long suppression of Canada Thistle and Field Bindweed, apply Maxunitech Glufosinate 200 SN at a rate of 2 L/ha when the weeds are in the 1- 6 leaf stage.

**Quackgrass:** For season long suppression of Quackgrass, apply Maxunitech Glufosinate 200 SN at a rate of 2.5 L/ha plus ammonium sulphate at a rate of 6 L/ha (49% solution) OR 3 kg/ha (99%) when the Quackgrass is in the 1-4 leaf stage.

**Voluntary Crop Weeds:** Maxunitech Glufosinate 200 SN will not control volunteer crop weeds that emerge from previous plantings of glufosinate ammonium tolerant crops.

**ADDITIVES:** For enhanced control of Velvetleaf, add ammonium sulphate to the tank at a rate of 6 L/ha (49% solution) OR 3 kg/ha (99%).

**SECOND APPLICATION:** A second application of Maxunitech Glufosinate 200 SN can be made to fields treated initially with up to 2.5 L/ha if new weed germination or growth is present.

Apply when the new weed growth is in the correct leaf stage and up to the maximum leaf stage of the crop.

Application after the 8-leaf stage of Corn must be made with drop nozzles as a directed spray below the whorl of the Corn plant.

Application in Canola must be made prior to the bolting stage of the Canola.

Do not apply beyond the flowering stage of Soybeans.

Do not apply more than a total of 4.5 L/ha in one season.

**SPLIT APPLICATION PROGRAM:** For season long control of the above listed weeds a split application program with Maxunitech Glufosinate 200 SN can be utilised. The first application must be a minimum of 2 L/ha and at the proper weed stage. For the second application, the rate of 1.25 L/ha may be utilized provided Maxunitech Glufosinate 200 SN is applied as soon as the second flush of weeds occurs and before the maximum leaf stage of the crop.

**FOR CANOLA HYBRID SEED PRODUCTION** - Two applications are required to remove the segregating wild type plants. The first application should occur when the Canola is in the 2-4 leaf stage, the second application when the Canola plants are in the 4-6 leaf stage and the third application, if necessary, may be applied to the Canola up until, but prior to, the time of bolting. All applications of Maxunitech Glufosinate 200 SN for hybrid seed production should be made at 2.5 L/ha.

## Section 8: Tank Mixes

Consult the label of the tank mix partner for further instructions regarding directions for use, restrictions and precautions. Certain tank mix partners may have additional restrictions for leaf staging.

For control of weeds present and/or control of later flushes of annual broadleaf weeds and/or annual grasses (see Tank Mix Directions), Maxunitech Glufosinate 200 SN may be tank mixed with the following herbicides when used in Eastern Canada and British Columbia:

CROP	TANK MIX PRODUCT	RATE	DIRECTIONS
CANOLA	Centurion <sup>®</sup> or Select <sup>®</sup> + Amigo Adjuvant at 0.5% v/v	63 mL/ha	For control of Volunteer Barley and improved control of Wild Oats, apply Maxunitech Glufosinate 200 SN at a rate of 2.0 – 2.5 L/ha plus Centurion or Select at 63 mL/ha with adjuvant as recommended on the Centurion or Select label. Apply when the weeds are in the 1-4 leaf stage with a maximum of 2 tillers.
CORN	AAtrax Liquid 480 Herbicide	1.75 – 2.33 L/ha	Mix with the appropriate rate of Maxunitech Glufosinate 200 SN for the targeted weed species. Adjust atrazine rate to the level of residual activity desired. This tank mix will not control later flushes of triazine resistant weeds.

CORN	Distinct®	285 g/ha	Mix with the appropriate rate of Maxunitech Glufosinate 200 SN Herbicide for the targeted weed species. This tank mix will provide enhanced control of labeled annual broadleaf weeds.
CORN	Marksman®	2.5 - 3.7 L/ha	Mix with the appropriate rate of Maxunitech Glufosinate 200 SN for the targeted weed species. Adjust Marksman rate to the level of residual activity desired. This tank mix will provide residual control of annual grasses and annual broadleaf weeds.
SOYBEA N	Pursuit®	312 mL/ha	Mix with the appropriate rate of Maxunitech Glufosinate 200 SN for the targeted weed species. This tank mix will provide residual control of annual grasses and annual broadleaf weeds.
SOYBEA N	Basagran® Forte	1.75 L/ha	Mix with the appropriate rate of Maxunitech Glufosinate 200 SN for the targeted weed species. This tank mix will provide enhanced control of annual broadleaf weeds.

**Note:** Do not tank mix Maxunitech Glufosinate 200 SN with fertilizers or any other chemical additives unless recommended on this label.

### **Section 9: Resistance Management Recommendations**

For resistance management, Maxunitech Glufosinate 200 SN is a Group 10 herbicide. Any weed population may contain or develop plants naturally resistant to Maxunitech Glufosinate 200 SN and other Group 10 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

Where possible, rotate the use of Maxunitech Glufosinate 200 SN or other Group 10 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.

Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.

Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.

Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible, by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.

Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.

For further information or to report suspected resistance, contact Maxunitech North America, Inc. at 1-855-462-9621 or 86-571-28007837.

## **Section 10: Application Instructions and Cautions**

### **Application Instructions:**

For best results, apply to emerged, young, actively growing weeds. Weeds that emerge after application will not be controlled.

Maxunitech Glufosinate 200 SN will have an effect on weeds that are larger than the recommended leaf stage, however speed of activity and control may be reduced.

Apply Maxunitech Glufosinate 200 SN in a minimum of 110 L/ha of water, at a pressure of 175-275 kPa and at a ground speed of 6-10 kph. The use of 80° or 110° flat fan nozzles is highly recommended for optimum spray coverage and canopy penetration. Application of the spray at a 45° angle forward will result in better spray coverage. Follow directions elsewhere on the label for the correct rate and timing of application.

The speed of action of Maxunitech Glufosinate 200 SN is influenced by environmental factors.

At cool temperatures (below 10°C), poor moisture and low humidity, speed of action may be reduced. Generally, visual symptoms appear 2-4 days after application.

**IF RAINFALL OCCURS WITHIN 4 HOURS OF APPLICATION, EFFECTIVENESS MAY BE REDUCED.**

Weed control may be reduced when heavy dew, fog or mist are present at the time of application.

Maxunitech Glufosinate 200 SN works primarily as a contact herbicide.

Thorough coverage of the weeds to be controlled is essential.

**Application Cautions:**

**Field sprayer application:** **DO NOT** apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. **DO NOT** apply with sprays finer than the American Society of Agricultural and Biological Engineers (ASABE) S572 (572.1 to 572.3) medium classification. Boom height must be 60 cm or less above the crop or ground.

**DO NOT** apply by air.

- **GROUND APPLICATION ONLY.**
- Do not use flood jet nozzles, controlled droplet application equipment or air-assisted spray equipment.
- Uniform, thorough spray coverage is important to achieve consistent weed control.
- **DO NOT** apply if winds exceed 16 kph when using open boom sprayers. **DO NOT** apply if winds exceed 25 kph when using hooded sprayers.
- Avoid contact with desirable plants or crops either from direct application or from spray drift as severe damage may occur.
- Plant back Intervals:
- 70 days for Buckwheat, Barley, Millet, Oats, Rye, Sorghum, Triticale and Wheat.
- 120 days for all other crops except Field Corn, Canola and Soybeans. No plant back interval is required for Field Corn, Canola or Soybeans.
- Application of Maxunitech Glufosinate 200 SN to non-tolerant crops will result in severe crop injury or death of the crop.
- **SUSCEPTIBLE WEEDS:** Maxunitech Glufosinate 200 SN has an effect on all weeds and crops except for those crops which are developed to be tolerant to applications of Maxunitech Glufosinate 200 SN including when they become voluntary weeds.

**Spray buffer zones:**

A spray buffer zone is NOT required for:

- uses with hand-held application equipment permitted on this label.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

Method of Application	Crop	Spray buffer zones (metres) Required for the Protection of:		
		Freshwater Habitat of Depths:		Terrestrial Habitat
		Less than 1 m	Greater than 1 m	
Field Sprayer	Corn, Canola, Soybean	1	0	1

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASABE) category indicated on the labels for those tank-mix partners.

The spray buffer zones for conventional application equipment can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Drift Mitigation portion of the Canada.ca website.

## **Section 11: Mixing Instructions**

Maxunitech Glufosinate 200 SN must be applied with properly calibrated, clean equipment. Maxunitech Glufosinate 200 SN is specially formulated to mix readily in water.

Prior to adding Maxunitech Glufosinate 200 SN to the spray tank, ensure that the spray tank is thoroughly clean (see Section 12 “SPRAYER CLEANUP”).

1. Fill tank to three-quarters full with clean water prior to adding Maxunitech Glufosinate 200 SN.
2. Add the correct amount of Maxunitech Glufosinate 200 SN.
3. Add the remaining amount of water, begin agitation, and spray out immediately.

**Note:** With the exception of use with CENTURION or SELECT, when tank mixing Maxunitech Glufosinate 200 SN always add the tank mix partner to the tank first, ensure that it is thoroughly mixed, THEN add Maxunitech Glufosinate 200 SN. When using ammonium sulphate, ALWAYS add it to the tank first, prior to the addition of the tank mix partner and Maxunitech Glufosinate 200 SN. Refer to the following specific instructions for mixing with CENTURION or SELECT.

### **Maxunitech Glufosinate 200 SN and CENTURION or SELECT: Glufosinate Ammonium Tolerant Canola only**

When tank mixing Maxunitech Glufosinate 200 SN and CENTURION or SELECT, always add AMIGO adjuvant to the tank first, *then* add Maxunitech Glufosinate 200 SN to the tank followed by the CENTURION or SELECT.

**The addition of an anti-foaming agent may reduce foaming, especially when using soft water.**

1. Thoroughly clean the sprayer by flushing the system with water containing detergent (see Section 12, “Sprayer Cleanup”).
2. Fill clean spray tank half full with clean water. Start agitation system.
3. Add the correct amount of AMIGO. Continue to agitate until AMIGO is thoroughly mixed.
4. STOP agitation. Add the required amount of Maxunitech Glufosinate 200 SN and Crop Desiccant MP to the spray tank. Start agitation system.
5. Add the correct amount of CENTURION or SELECT along with the remaining amount of water necessary to fill the spray tank.
6. Continue to agitate or run the by-pass system and spray out immediately.
7. After any break in the spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to remix the spray materials. Do not allow the mixture to sit overnight.
8. If an oil film starts to build up in the tank, drain it and clean tank with strong detergent

solution.

9. Immediately after use, thoroughly clean the sprayer by flushing the system with water containing detergent (see Section 12, “Sprayer Cleanup”).

**Note:** The addition of an anti-foaming agent may reduce foaming, especially when using soft water.

**Note:** Ensure that all circuits (pipes, booms, etc.) have the correct Maxunitech Glufosinate 200 SN/water concentration before application is started.

## **Section 12: Sprayer Cleanup**

Before and after using Maxunitech Glufosinate 200 SN always complete a thorough cleaning of the spray tank, lines and filter. Spray equipment should be thoroughly rinsed using a strong detergent solution.

## **Section 13: Livestock Feeding and Preharvest Intervals (PHIs)**

### **CANOLA:**

Grain and meal from treated Canola can be fed to livestock.

DO NOT graze or feed other portions of the treated Canola to livestock; there are not sufficient data to support such use.

When Maxunitech Glufosinate 200 SN is tank mixed with CENTURION or SELECT, observe a PHI of 60 days from the date of treatment (or last treatment when a second application has been made).

### **CORN:**

DO NOT apply within 86 days of harvesting Corn grain.

DO NOT graze the treated fields within 20 days of application.

### **SOYBEAN:**

DO NOT apply within 70 days of harvesting Soybean seed.

DO NOT graze the treated fields within 20 days of application.

## **Section 14: Notices**

### **NOTICE TO USER:**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.