

2025-3495  
2025-09-22

Container Label

<b>GROUP</b>	<b>14</b>	<b>HERBICIDE</b>
--------------	-----------	------------------

**SPECIAL HERBICIDE**

Water dispersible granular herbicide for pre-seed, pre-emergent and chemfallow application for control of broadleaf weeds, for pre-harvest weed management in wheat, barley and triticale, and for harvest aid in canola, dry common beans, chickpeas, dry field peas, faba beans, flax, red lentil varieties, mustard, soybeans and sunflowers

**COMMERCIAL (AGRICULTURAL)**

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND  
INTERIOR OF BRITISH COLUMBIA (INCLUDING THE PEACE RIVER REGION) ONLY

**ACTIVE INGREDIENT:** Saflufenacil..... 70%

**REGISTRATION NO. 35610**

**PEST CONTROL PRODUCTS ACT**

READ THE LABEL AND BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

**NET CONTENTS:** 0.25 kg – 100 kg, Bulk

**SHARDA Cropchem Limited**  
2nd Floor, Prime Business Park  
Dashrathlal Joshi Road  
Vile Parle (West)  
Mumbai - 400056, India

Canadian Agent  
**SHARDA Cropchem Limited**  
63 Kingsview Blvd  
Etobicoke, Ontario, CA  
M9R 1V1  
1-844-810-5720  
1-416-840-5639

## PRECAUTIONS

1. **KEEP OUT OF REACH OF CHILDREN.**
2. **DO NOT** take internally. Harmful if swallowed.
3. **DO NOT** get in eyes or on skin. Causes eye and skin irritation. Avoid inhalation of vapor or spray.
4. Wash exposed areas of skin thoroughly after handling and before eating, drinking or smoking or going to the washroom. Take a shower immediately after work.
5. Ground and aerial scenarios:

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair, and a NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested during mixing and loading.

For application using ground equipment: Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes. Gloves are not required during application within a closed cab.

For application using aerial equipment: Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes. Gloves are not required during application within a closed cockpit.

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. The pilot is allowed to load premixed chemicals with a closed system.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

6. If clothing becomes contaminated, remove immediately and wash. Store and wash all protective clothing separately from household laundry. Wash in detergent and hot water before reuse. Wear freshly laundered clothes daily.
7. Apply only to agricultural crops when the potential for drift to areas of human habitation or areas of and human activity, such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

## FIRST AID

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

## **TOXICOLOGICAL INFORMATION**

Treat symptomatically.

## **ENVIRONMENTAL PRECAUTIONS**

TOXIC to non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

## **STORAGE**

1. Store this product away from food or feed.
2. Store the product in original, tightly-closed container and do not allow water to be introduced into this container.
3. **DO NOT** ship or store the product near seed or fertilizers.
4. Store the product in a cool, dry, locked, well-ventilated area without floor drain.
5. Herbicides should be shipped or stored separately from other pesticides to avoid cross contamination.

## **DISPOSAL**

### For recyclable containers

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

### For returnable containers

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For containers that can be refilled for the user by the distributor/dealer

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

### Disposal of unused, unwanted product

For information on disposal of unused, unwanted product, contact the registrant or the provincial/territorial regulatory agency. Contact the registrant and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

**SPECIAL HERBICIDE**

Water dispersible granular herbicide for pre-seed, pre-emergent and chemfallow application for control of broadleaf weeds, for pre-harvest weed management in wheat, barley and triticale, and for harvest aid in canola, dry common beans, chickpeas, dry field peas, faba beans, flax, red lentil varieties, mustard, soybeans and sunflowers

**COMMERCIAL (AGRICULTURAL)**

FOR SALE FOR USE IN THE PRAIRIE PROVINCES AND  
INTERIOR OF BRITISH COLUMBIA (INCLUDING THE PEACE RIVER REGION) ONLY

**ACTIVE INGREDIENT:** Saflufenacil..... 70%

**REGISTRATION NO. 35610**

**PEST CONTROL PRODUCTS ACT**

READ THE LABEL AND BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

**NET CONTENTS:** 0.25 kg – 100 kg, Bulk

**SHARDA Cropchem Limited**  
2nd Floor, Prime Business Park  
Dashrathlal Joshi Road  
Vile Parle (West)  
Mumbai - 400056, India

Canadian Agent  
**SHARDA Cropchem Limited**  
63 Kingsview Blvd  
Etobicoke, Ontario, CA  
M9R 1V1  
1-844-810-5720  
1-416-840-5639

## GENERAL INFORMATION

**SPECIAL HERBICIDE** is a water dispersible granular herbicide for broadleaf weed control.

**SPECIAL HERBICIDE** is rapidly absorbed by root and foliar uptake; once absorbed it exhibits mobility in plants. **SPECIAL HERBICIDE** is a potent inhibitor of protoporphyrinogen oxidase (PPO). Cell membrane damage induced by inhibition of PPO leads to plant death. Susceptible weeds develop injury symptoms within hours of application under active growing conditions; plant death occurs within 3 to 5 days depending upon growing conditions.

**SPECIAL HERBICIDE** is recommended for pre-seed and pre-emergent applications. **SPECIAL HERBICIDE** may also be applied in fallow crop lands and post-harvest, and as a pre-harvest weed management treatment and harvest aid in registered crops.

**SPECIAL HERBICIDE** does not control grass weeds. **SPECIAL HERBICIDE should always be tank mixed with glyphosate** for broad spectrum weed control.

**SPECIAL HERBICIDE** is a broad spectrum weed resistance management tool for activity on a range of broadleaf weeds.

## PRECAUTIONS

1. **KEEP OUT OF REACH OF CHILDREN.**
2. **DO NOT** take internally. Harmful if swallowed.
3. **DO NOT** get in eyes or on skin. Causes eye and skin irritation. Avoid inhalation of vapor or spray.
4. Wash exposed areas of skin thoroughly after handling and before eating, drinking or smoking or going to the washroom. Take a shower immediately after work.
5. Ground and aerial scenarios:

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair, and a NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested during mixing and loading.

For application using ground equipment: Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes. Gloves are not required during application within a closed cab.

For application using aerial equipment: Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes. Gloves are not required during application within a closed cockpit.

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. The pilot is allowed to load premixed chemicals with a closed system.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

6. If clothing becomes contaminated, remove immediately and wash. Store and wash all protective clothing separately from household laundry. Wash in detergent and hot water before reuse. Wear freshly laundered clothes daily.

7. Apply only to agricultural crops when the potential for drift to areas of human habitation or areas of and human activity, such as houses, cottages, schools and recreational areas, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

#### **FIRST AID**

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

#### **TOXICOLOGICAL INFORMATION**

Treat symptomatically.

#### **ENVIRONMENTAL PRECAUTIONS**

TOXIC to non-target terrestrial plants.

Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

#### **STORAGE**

1. Store this product away from food or feed.
2. Store the product in original, tightly-closed container and do not allow water to be introduced into this container.
3. **DO NOT** ship or store the product near food, feed, seed or fertilizers.
4. Store the product in a cool, dry, locked, well-ventilated area without floor drain.
5. Herbicides should be shipped or stored separately from other pesticides to avoid cross contamination.

## DISPOSAL

### For Recyclable containers

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

### For returnable containers

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For containers that can be refilled for the user by the distributor/dealer

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

### Disposal of unused, unwanted product

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

## NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

## REGISTERED CROPS

### PRE-SEED OR PRE-EMERGENT

**SPECIAL HERBICIDE** is registered for use prior to the following crops as a pre-seed or pre-emergent application.

Barley	Lentils*
Bromegrass, seedling (seed production, hay and forage)	Oats
Creeping red fescue and timothy, seedling (seed production and forage)	Peas (dried field)
Canary seed	Wheat (spring, winter and durum)
Chickpeas	Corn
Faba beans	Soybeans*

\* Rate restrictions apply. Refer to the crop specific section for details.

## CHEMFALLOW HARVEST AID

**SPECIAL HERBICIDE** is registered for use as a desiccant in the following crops:

Canola (all types)	Lentils (red lentil varieties only) <sup>1</sup>
Chickpeas	Mustard <sup>2</sup>
Dry common beans	Peas (dried field)
Faba beans	Soybeans
Flax	Sunflower

<sup>1</sup> Always tank mix **Special Herbicide** with glyphosate when applying as a desiccant to red lentils.

<sup>2</sup> All classes, including brown, oriental, canola quality *Brassica juncea*, *Brassica juncea* varieties with the **Clearfield®** trait, and yellow mustard.

## PRE-HARVEST WEED MANAGEMENT

**SPECIAL HERBICIDE** is registered for use as a pre-harvest treatment to improve dry down of volunteer canola (all types including Roundup Ready), common ragweed, Canada fleabane, redroot pigweed and wild buckwheat in the following crops:

Barley (including feed varieties)
Triticale
Wheat (including durum, spring and winter wheat)

## DIRECTIONS FOR USE

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Field sprayer application: DO NOT apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. DO NOT apply with sprays finer than the American Society of Agricultural and Biological Engineers (ASABE) S572 (572.1 to 572.3) medium classification. Boom height must be 60 cm or less above the crop or ground.

Conventionally piloted aircraft application: DO NOT apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with sprays finer than the American Society of Agricultural and Biological Engineers (ASABE) S572 (572.1 to 572.3) medium classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length MUST NOT exceed 65% of the wing- or rotorspan.

Apply only by fixed-wing or rotary aircraft equipment, which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate specified for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices or a Global Positioning System (GPS).

### Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

### Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-844-810-5720 or obtain technical advice from the distributor or your provincial or territorial agricultural representative. Application of this specific product must meet and/or conform to the following: Ensure thorough coverage of foliage. Consult nozzle manufacturer's recommendation for spray pressures for specific nozzles.

### SPRAY BUFFER ZONES

A spray buffer zone is NOT required for:

- uses with hand-held application equipment permitted on this label,
- low-clearance hooded or shielded sprayers that prevent spray contact with crop, fruit or foliage.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands).

Method of application	Crop	Spray Buffer zones (metres) required for the protection of:	
		Terrestrial Habitat	
Field sprayer	Lentils (pre-seed or pre-emergent)	3	
	Soybean (pre-seed or pre-emergent)	4	
	Barley, canary seed, chickpea, corn, faba beans, oats, dried field peas, forage grasses (seedling creeping red fescue, timothy, brome grass), wheat (pre-seed or pre-emergent); Chemfallow; Canola, faba beans, flax, lentil, chickpea, mustard, soybean, dried field peas, sunflower, dry common beans, wheat, barley, triticale (harvest aid)	10	
Aerial	Canola, faba beans, flax, lentil, chickpea, mustard, soybean, dried field peas, sunflower, dry common	Fixed wing	175
		Rotary wing	150

	bean, wheat, barley, triticale (harvest aid)		
--	---	--	--

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASABE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product for conventional application equipment can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Drift Management portion of the Canada.ca website.

### CROP USE RATES – SPECIAL HERBICIDE

For all **SPECIAL HERBICIDE** solo applications applied pre-seed, pre-emergent or as a chemfallow treatment, apply with MERGE® Adjuvant or MSO Concentrate with Leci-Tech at a rate of 0.5 – 1 L/ha.

### PRE-SEED OR PRE-EMERGENT

Crop	Use Rate (g/ha)
Lentils*	26
Soybean**	26 – 36
Barley	26 – 71
Bromegrass, seedling (seed production, hay and forage) *** Creeping red fescue and timothy, seedling***	26 – 71
Canary seed	26 – 71
Chickpea Kabuli	26 – 71
Corn (field, sweet****)	26 – 71
Faba beans	26 – 71
Oats	26 – 71
Peas (dried field)	26 – 71
Wheat (spring, durum, winter)	26 – 71
Chemfallow	26 – 71
* Rate restrictions apply. Do not use rates higher than 26 g/ha or injury could result. See the crop specific section for additional details.	
** Rate restrictions apply. Do not use rates higher than 36 g/ha or injury could result. See the crop specific section for additional details. Some soybeans cultivars may be more sensitive to saflufenacil and injury might occur.	
*** For seed production, forage and hay. <b>SPECIAL HERBICIDE</b> should not come into contact with the seed. Ensure adequate soil coverage for pre-emergent applications to forage grasses and grass grown for seed.	
**** Some sweet corn hybrids may be more sensitive to saflufenacil and injury might occur.	

### CHEMFALLOW

Crop	Use Rate (g/ha)	Application Timing
Chemfallow	26 – 71	Post-emergence in fallow croplands and post-harvest.

## HARVEST AID

**SPECIAL HERBICIDE** may be used as a harvest aid to accelerate the rate of crop dry down and improve crop uniformity to facilitate direct combining. Early application may result in yield and/or seed quality loss.

The dry down of crops will be best under favorable environmental conditions like warm temperatures, good moisture conditions and low humidity.

Harvesting of crops can be done when plant material is dry and seed moisture level allows efficient harvesting. Under ideal conditions, harvest can normally commence within 7-14 days after desiccation when applied at the appropriate crop stage recommendation. Adverse weather conditions such as rainfall, cool temperatures, shorter day length and high humidity may slow plant desiccation and keep seed moisture levels high which can delay commencement of harvest after the **SPECIAL HERBICIDE** application. Consult your Sharda Cropchem Limited representative for further information on the timing of harvest after a pre-harvest application.

Crop	Use Rate (g/ha)	Application Timing
Canola (all types), faba beans, chickpeas, dry common beans, flax, lentils <sup>2</sup> , mustard <sup>1</sup> , peas (dried field), soybeans, sunflower	52	Harvest aid

<sup>1</sup> All classes, including brown; oriental; canola quality *Brassica juncea*; *Brassica juncea* varieties with the **Clearfield** trait; and yellow mustard.

<sup>2</sup> Apply only to red lentil varieties.

## Aerial Application

**SPECIAL HERBICIDE** can be used for aerial application when used as a harvest aid or for pre-harvest weed management (see the “**PRE-HARVEST WEED MANAGEMENT**” section) in the registered crops.

Ensure thorough coverage of foliage. Consult nozzle manufacturer’s recommendation for spray pressures for specific nozzles.

## HARVEST AID - CROP SPECIFIC RECOMMENDATIONS STAND-ALONE

See the “**HARVEST AID**” section above, for further use instructions.

## DESICCATION - DRY COMMON BEANS / SOYBEANS

Rate	52 g/ha of <b>SPECIAL HERBICIDE</b> + adjuvant
Water Volume	200 L/ha ground; 50 L/ha aerial
Surfactant/Adjuvant	Add MERGE Adjuvant at a rate of 0.5 – 1 L/ha or MSO Concentrate with Leci-Tech at a rate of 1 L/ha.
Application Timing	Apply when stems are green to brown in colour and pods are mature (yellow – brown) and 80 – 90% of the original leaves have dropped.

Directions for Use – Ground	Ground application may be done with a standard boom sprayer.
Directions for Use – Aerial	For aerial application use a minimum water volume of 50 L/ha. Please see above for aerial application use and operator precautions.
Remarks	Do not graze or feed treated hay or straw to livestock.

### DESICCATION - CHICKPEAS

Rate	52 g/ha of SPECIAL HERBICIDE + adjuvant
Water Volume	200 L/ha ground; 50 L/ha aerial
Surfactant/Adjuvant	Add MERGE Adjuvant at a rate of 0.5 – 1 L/ha or MSO Concentrate with Leci-Tech at a rate of 1 L/ha.
Application Timing	For Desi type, apply at the time swathing would normally commence, when the majority of plants are yellow and most pods are mature and seeds have turned from green to yellow or brown. Upper part of plant may still be green.  For Kabuli type, apply when the majority of plants and pods are ripe and dry with seeds turned from green to white or tan, and detached from the pods. Dry down is less complete in Kabuli type due to its thick pod wall.
Directions for Use – Ground	Ground application may be done with a standard boom sprayer.
Directions for Use – Aerial	For aerial application use a minimum water volume of 50 L/ha. Please see above for aerial application use and operator precautions.
Remarks	Do not graze or feed treated hay or straw to livestock.

### DESICCATION – FIELD PEAS

Rate	52 g/ha of <b>SPECIAL HERBICIDE</b> + adjuvant
Water Volume	200 L/ha ground; 50 L/ha aerial
Surfactant/Adjuvant	Add MERGE Adjuvant at a rate of 0.5 – 1 L/ha or MSO Concentrate with Leci-Tech at a rate of 1 L/ha.
Application Timing	Apply when the majority of pods are brown (70-80%).
Directions for Use – Ground	Ground application may be done with a standard boom sprayer.
Directions for Use – Aerial	For aerial application use a minimum water volume of 50 L/ha. Please see above for aerial application use and operator precautions.
Remarks	Desiccation treated pea vines may be grazed or fed to livestock.

### DESICCATION – SUNFLOWER

Rate	52 g/ha of <b>SPECIAL HERBICIDE</b> + adjuvant
Water Volume	200 L/ha ground; 50 L/ha aerial
Surfactant/Adjuvant	Add MERGE Adjuvant at a rate of 0.5 – 1 L/ha or MSO Concentrate with Leci-Tech at a rate of 1 L/ha.
Application Timing	Apply when the backs of the heads and bracts are turning yellow, and seed moisture is 20 – 30%.
Directions for Use – Ground	Ground application may be done with a standard boom sprayer.

Directions for Use – Aerial	For aerial application use a minimum water volume of 50 L/ha. Please see above for aerial application use and operator precautions.
-----------------------------	---

### DESICCATION – CANOLA (ALL TYPES) AND MUSTARD<sup>1</sup>

Rate	52 g/ha of <b>SPECIAL HERBICIDE</b> + adjuvant
Water Volume	200 L/ha ground; 50 L/ha aerial
Surfactant/Adjuvant	Add MERGE Adjuvant at a rate of 0.5 – 1 L/ha or MSO Concentrate with Leci-Tech at a rate of 1 L/ha.
Application Timing	Apply when 65 – 80% of seeds have changed colour.
Directions for Use – Ground	Ground application may be done with a standard boom sprayer.
Directions for Use – Aerial	For aerial application use a minimum water volume of 50 L/ha. Please see above for aerial application use and operator precautions.
Remarks	<p>It is recommended that the application of <b>SPECIAL HERBICIDE</b> as a desiccant in canola and mustard be made to shatter resistant varieties.</p> <p>Under ideal conditions, harvest can normally commence within 14-21 days after desiccation when applied at the appropriate crop stage recommendation. Adverse weather conditions such as rainfall, cool temperatures, shorter day length and high humidity may slow the plant desiccation and keep seed moisture levels high which can delay commencement of harvest after the <b>Special Herbicide</b> application.</p>

<sup>1</sup>All classes, including brown; oriental; canola quality *Brassica juncea*; *Brassica juncea* varieties with the **Clearfield** trait; and yellow mustard

### DESICCATION – FLAX

Rate	52 g/ha of <b>SPECIAL HERBICIDE</b> + adjuvant
Water Volume	200 L/ha ground; 50 L/ha aerial
Surfactant/Adjuvant	Add MERGE Adjuvant at a rate of 0.5 – 1 L/ha or MSO Concentrate with Leci-Tech at a rate of 1 L/ha.
Application Timing	Apply when 75% of bolls have turned colour.
Directions for Use – Ground	Ground application may be done with a standard boom sprayer.
Directions for Use – Aerial	For aerial application use a minimum water volume of 50 L/ha. Please see above for aerial application use and operator precautions.

### DESICCATION – FABA BEANS

Rate	52 g/ha of <b>SPECIAL HERBICIDE</b> + adjuvant
Water Volume	200 L/ha ground; 50 L/ha aerial
Surfactant/Adjuvant	Add MERGE Adjuvant at a rate of 0.5 – 1 L/ha or MSO Concentrate with Leci-Tech at a rate of 1 L/ha.
Application Timing	Apply when 80% of lower pods have turned black, middle pods have turned yellow/tan, and top green pods have firm seed.
Directions for Use – Ground	Ground application may be done with a standard boom sprayer.

Directions for Use – Aerial	For aerial application use a minimum water volume of 50 L/ha. Please see above for aerial application use and operator precautions.
Remarks	Do not graze or feed treated hay or straw to livestock.

### PRE-HARVEST WEED MANAGEMENT

**SPECIAL HERBICIDE** may be used as a pre-harvest treatment in wheat (including durum, spring, and winter wheat), barley (including feed varieties) and triticale to improve dry down of volunteer canola (all types including Roundup Ready), common ragweed, Canada fleabane, redroot pigweed and wild buckwheat and to facilitate direct combining.

Early application may result in yield loss. **SPECIAL HERBICIDE** when used alone as a pre-harvest treatment will not affect seed germination if applied according to label recommendations.

### WHEAT, BARLEY AND TRITICALE

Rate	36 – 52 g/ha of <b>SPECIAL HERBICIDE</b> + adjuvant
Water Volume	100 – 200 L/ha ground; 50 L/ha aerial
Surfactant/Adjuvant	Add MERGE Adjuvant at a rate of 0.5 – 1 L/ha or MSO Concentrate with Leci-Tech at a rate of 1 L/ha.
Application Timing	Hard dough stage; a thumbnail impression remains on seed; less than 30% moisture.
Weeds Controlled	Improved dry down of volunteer canola (all types including Roundup Ready), common ragweed, Canada fleabane, redroot pigweed and wild buckwheat.
Directions for Use – Ground	Ground application may be done with a standard boom sprayer.
Directions for Use – Aerial	For aerial application use a minimum water volume of 50 L/ha. Please see above for aerial application use and operator precautions.
Remarks	Treated barley, wheat and triticale straw may be grazed or fed to livestock. Use higher water volume for dense crop stands and higher weed pressure.

### WEEDS CONTROLLED

**SPECIAL HERBICIDE** can be applied pre-seed or pre-emergence to the crop to control weeds listed below.

**SPECIAL HERBICIDE** should always be applied in combination with glyphosate for broad spectrum weed control including grassy weeds.

### Weeds Controlled For Rapid Burn down (Pre-seed or Pre-emergent)

#### Use Rate: 26 g/ha

Weed	Maximum Weed Stage
Kochia	15 cm
Canada fleabane	8 leaf
Cleavers	4-whorl stage
Lamb's-quarters	8 leaf
Narrow-leaved hawk's beard	8 cm
Redroot pigweed	8 leaf

Round-leaved mallow	8 leaf
Stinkweed	8 leaf
Volunteer canola (all types including Roundup Ready®)	8 leaf
Wild buckwheat	8 leaf
Wild mustard	8 leaf

**Use Rate: 36 g/ha**

**SPECIAL HERBICIDE** applied pre-seed or pre-emergence at 36 g/ha will provide rapid burndown control of the following weeds in addition to those listed above:

Weed	Maximum Weed Stage
Common ragweed	8 leaf
Giant ragweed	8 leaf
Lady's-thumb	6 leaf
Perennial sow-thistle (top growth burndown control)	8 leaf
Prickly lettuce (top growth only)	9 leaf
Shepherd's-purse	Full flower

**For Rapid Burndown and Suppression of Secondary Weed Flushes (Pre-seed or Pre-emergent)**

**Use Rate: 52 - 71 g/ha**

Weed	Maximum Weed Stage
Cleavers	4-whorl stage
Redroot pigweed	8 leaf
Stinkweed	8 leaf
Volunteer canola (all types including Roundup Ready)	8 leaf
Wild buckwheat	8 leaf
Wild mustard	8 leaf

### **TANK MIXES – PRE-SEED, PRE-EMERGENT AND CHEMFALLOWSPECIAL HERBICIDE + GLYPHOSATE**

For broad spectrum weed control, **SPECIAL HERBICIDE** should always be tank mixed with glyphosate present as isopropylamine salt, di-ammonium salt or potassium salt. **SPECIAL HERBICIDE** is compatible with all liquid glyphosate formulations, in which glyphosate is present as isopropylamine salt, di-ammonium salt or potassium salt, as a tank-mix partner.

Tank mixing **SPECIAL HERBICIDE** with glyphosate will provide control of all weeds controlled by glyphosate in addition to the broadleaf weeds listed on **SPECIAL HERBICIDE** label. Consult the glyphosate label for a complete list of weeds controlled by glyphosate.

Tank mixing is a recognized strategy to delay herbicide resistance as well as improve the weed spectrum controlled.

### TANK MIX CROP USE RATES

#### SPECIAL HERBICIDE + Glyphosate Use Rate (360 g/L equivalent) 1.25 – 2.5 L/ha

Crop	Use Rate (g/ha)
Lentils*	26
Soybean**	26 – 36
Barley	26 – 71
Bromegrass, seedling (seed production, hay and forage) *** Creeping red fescue and timothy, seedling***	26 – 71
Canary seed	26 – 71
Chickpea, Kabuli	26 – 71
Corn (field, sweet****)	26 – 71
Faba beans	26 – 71
Oats	26 – 71
Peas (dried field)	26 – 71
Wheat (spring, durum, winter)	26 – 71
Chemfallow	26 – 71
* Rate restrictions apply. Do not use rates higher than 26 g/ha or injury could result. See the crop specific section for additional details.	
** Rate restrictions apply. Do not use rates higher than 36 g/ha or injury could result. See the crop specific section for additional details. Some soybeans cultivars may be more sensitive to saflufenacil and injury might occur.	
*** For seed production, forage and hay. <b>SPECIAL HERBICIDE</b> should not come into contact with the seed. Ensure adequate soil coverage for pre-emergent applications to forage grasses and grass grown for seed.	
**** Some sweet corn hybrids may be more sensitive to saflufenacil and injury might occur.	

For tank mix applications of **SPECIAL HERBICIDE** with glyphosate, apply with either MERGE Adjuvant, MSO Concentrate with Leci-Tech at a rate of 0.5 – 1 L/ha.

Do not apply tank mix combinations by air.

When tank mixes are permitted, consult the labels of the tank mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASABE) category indicated on the labels for those tank mix partners.

### TANK MIX WEEDS CONTROLLED

#### Rapid Burn Down (Pre-seed or Pre-emergent)

**SPECIAL HERBICIDE** when tank mixed with glyphosate will provide rapid burn down of the following weeds in addition to weeds listed for treatment of **SPECIAL HERBICIDE**.

#### Use Rate: 26 g/ha

Weed	Maximum Weed Stage
Dandelion (top growth burndown control only)	15 cm

Flixweed	8 leaf
----------	--------

**CROP SPECIFIC RECOMMENDATIONS  
CHICKPEAS, FABA BEANS AND PEAS (dried field)**

Timing	Pre-seed or Pre-emergent
Rate	26 – 71 g/ha of <b>SPECIAL HERBICIDE</b> + 1.25 – 2.5 L/ha of glyphosate (360 g/L equivalent) + adjuvant
Water Volume	50 – 100 L/ha
Surfactant/Adjuvant	Add MERGE Adjuvant or MSO Concentrate with Leci-Tech at a rate of 0.5 – 1 L/ha.
Weeds Controlled	See the “ <b>WEEDS CONTROLLED</b> ” section for a complete list of weeds controlled by <b>SPECIAL HERBICIDE</b> . Refer to the glyphosate label for weeds controlled in addition to those controlled by <b>SPECIAL HERBICIDE</b> .

**CEREALS – BARLEY, CANARY SEED, OATS AND WHEAT (spring, durum, winter)**

Timing	Pre-seed or Pre-emergent
Rate	26 – 71 g/ha of <b>SPECIAL HERBICIDE</b> + 1.25 – 2.5 L/ha of glyphosate (360 g/L equivalent) + adjuvant
Water Volume	50 – 100 L/ha
Surfactant/Adjuvant	Add MERGE Adjuvant or MSO Concentrate with Leci-Tech at a rate of 0.5 – 1 L/ha.
Weeds Controlled	See the Section “ <b>WEEDS CONTROLLED</b> ” for a complete list of weeds controlled by <b>SPECIAL HERBICIDE</b> . Refer to the glyphosate label for weeds controlled in addition to those controlled by <b>SPECIAL HERBICIDE</b> .

**SEEDLING FORAGE GRASSES AND GRASS GROWN FOR SEED (seedling creeping red fescue, timothy, brome grass)**

**SPECIAL HERBICIDE** may be applied for weed control in seedling creeping red fescue, timothy and brome grass grown for seed production, forage and hay.

Timing	Pre-seed or Pre-emergent
Rate	26 – 71 g/ha of <b>SPECIAL HERBICIDE</b> + 1.25 – 2.5 L/ha of glyphosate (360 g/L equivalent) + adjuvant
Water Volume	50 – 100 L/ha
Surfactant/Adjuvant	Add MERGE Adjuvant or MSO Concentrate with Leci-Tech at a rate of 0.5 – 1 L/ha.
Weeds Controlled	See the “ <b>WEEDS CONTROLLED</b> ” section for a complete list of weeds controlled by <b>SPECIAL HERBICIDE</b> . Refer to the glyphosate label for weeds controlled in addition to those controlled by <b>SPECIAL HERBICIDE</b> .
Remarks	<b>SPECIAL HERBICIDE</b> should not come into contact with the seed. Ensure adequate soil coverage for pre-emergent applications to forage grasses and grass grown for seed.

## SOYBEANS

Timing	Pre-seed or Pre-emergent
Rate	26 – 36 g/ha of <b>SPECIAL HERBICIDE</b> + 1.25 – 2.5 L/ha of glyphosate (360 g/L equivalent) + adjuvant
Water Volume	50 – 100 L/ha
Surfactant/Adjuvant	Add MERGE Adjuvant or MSO Concentrate with Leci-Tech at a rate of 0.5 – 1 L/ha.
Weeds Controlled	See the “ <b>WEEDS CONTROLLED</b> ” section for a complete list of weeds controlled by <b>SPECIAL HERBICIDE</b> . Refer to the glyphosate label for weeds controlled in addition to those controlled by <b>SPECIAL HERBICIDE</b> .
Remarks	<p>Some soybean cultivars may be more sensitive to saflufenacil and injury might occur.</p> <p>When applying 36 g/ha pre-emergent to soybeans, DO NOT apply to coarse textured soils with less than 2% organic matter.</p> <p>When applying pre-emergent to soybeans, apply prior to when the soybeans cause the ground to crack and no more than 3 days after planting.</p>

## LENTILS (including Clearfield® Lentils)

Timing	Pre-seed or Pre-emergent
Rate	26 g/ha of <b>SPECIAL HERBICIDE</b> + 1.25 – 2.5 L/ha of glyphosate (360 g/L equivalent) + adjuvant
Water Volume	50 – 100 L/ha
Surfactant/Adjuvant	Add MERGE Adjuvant or MSO Concentrate with Leci-Tech at a rate of 0.5 – 1 L/ha.
Weeds Controlled	See the “ <b>WEEDS CONTROLLED</b> ” section for a complete list of weeds controlled by <b>SPECIAL HERBICIDE</b> . Refer to the glyphosate label for weeds controlled in addition to those controlled by <b>SPECIAL HERBICIDE</b> .
Remarks	<p>Rainfall shortly after product application can result in slight injury to the crop. Lentils will be more susceptible to injury on coarse texture and low organic matter soils. Injury will usually appear as leaf tissue necrosis on the outer edges of the leaves. Lentils will grow out of injury symptoms, and yield will not be impacted at recommended rates. The user should contact Sharda Cropchem Limited before applying any other soil applied herbicide with, before, or after applications of <b>SPECIAL HERBICIDE</b> + glyphosate. The addition of other soil applied herbicides may increase the sensitivity of lentils to <b>SPECIAL HERBICIDE</b> and injury may result.</p>

## CORN (field, sweet)

Timing	Pre-seed or Pre-emergent
Rate	26 – 71 g/ha of <b>SPECIAL HERBICIDE</b> + 1.25 – 2.5 L/ha of glyphosate (360 g/L equivalent) + adjuvant
Water Volume	50 – 100 L/ha
Surfactant/Adjuvant	Add MERGE Adjuvant or MSO Concentrate with Leci-Tech at a rate of 0.5 – 1 L/ha.

Weeds Controlled	See the “ <b>WEEDS CONTROLLED</b> ” section for a complete list of weeds controlled by <b>SPECIAL HERBICIDE</b> . Refer to the glyphosate label for weeds controlled in addition to those controlled by <b>SPECIAL HERBICIDE</b> .
Remarks	Some sweet corn hybrids may be more sensitive to saflufenacil and injury might occur.

### CHEMFALLOW

Timing	Chemfallow
Rate	26 – 71 g/ha of <b>SPECIAL HERBICIDE</b> + 1.25 – 2.5 L/ha of glyphosate (360 g/L equivalent) + adjuvant
Water Volume	50 – 100 L/ha
Surfactant/Adjuvant	Add MERGE Adjuvant or MSO Concentrate with Leci-Tech at a rate of 0.5 – 1 L/ha.
Weeds Controlled	See the “ <b>WEEDS CONTROLLED</b> ” section for a complete list of weeds controlled by <b>SPECIAL HERBICIDE</b> . Refer to the glyphosate label for weeds controlled in addition to those controlled by <b>SPECIAL HERBICIDE</b> .
Remarks	Apply to actively growing weeds less than 15 cm in height. Better coverage of the product results in enhanced control of weeds. For application to larger weeds or dense weed infestations, use a minimum water volume of 100 L per hectare.

### HARVEST AID – CROP SPECIFIC RECOMMENDATIONS TANK-MIX

**SPECIAL HERBICIDE** may be used as a harvest aid to accelerate the rate of crop dry down and improve crop uniformity to facilitate direct combining. Early application may result in yield and/or seed quality loss.

The dry down of crops will be best under favorable environmental conditions like warm temperatures, good moisture conditions and low humidity.

Harvesting of crops can be done when plant material is dry and seed moisture level allows efficient harvesting. Under ideal conditions, harvest can normally commence within 7-14 days after desiccation when applied at the appropriate crop stage recommendation. Adverse weather conditions such as rainfall, cool temperatures, shorter day length, and high humidity may slow plant desiccation and keep seed moisture levels high which can delay commencement of harvest after the **SPECIAL HERBICIDE** application. Consult your Sharda Cropchem Limited representative for further information on the timing of harvest after a pre-harvest application.

**SPECIAL HERBICIDE** may be applied in tank mix with glyphosate for additional pre-harvest weed control.

### DESICCATION – DRY COMMON BEANS\*/ SOYBEANS

Timing	Desiccation of DRY COMMON BEANS* / SOYBEANS
Rate	52 g/ha of <b>SPECIAL HERBICIDE</b> + 2.5 L/ha of glyphosate (360 g/L equivalent) + adjuvant
Water Volume	100 – 200 L/ha ground

Surfactant/Adjuvant	Add MERGE Adjuvant or MSO Concentrate with Leci-Tech at a rate of 0.5 – 1 L/ha.
Remarks	*Consult glyphosate label or Sharda Cropchem Limited representative for information on the use on specific varieties of dry common beans.
Application Timing	Apply when stems are green to brown in colour and pods are mature (yellow – brown) and 80 – 90% of the original leaves have dropped.
Directions for Use – Ground	Ground application may be done with a standard boom sprayer.
Remarks	Do not graze or feed treated hay or straw to livestock. See the “ <b>HARVEST AID – CROP SPECIFIC RECOMMENDATIONS TANK-MIX</b> ” section for the detailed use pattern. Consult the label of the tank mix partner product for further use instructions, precautions and restrictions. The most restrictive labeling applies to tank mixes.

### DESICCATION – CHICKPEAS

Timing	Desiccation of CHICKPEAS
Rate	52 g/ha of <b>SPECIAL HERBICIDE</b> + 900 g a.e/ha glyphosate (1.67 L/ha of 540 g/L glyphosate formulation) + adjuvant
Water Volume	100 – 200 L/ha ground
Surfactant/Adjuvant	Add MERGE Adjuvant or MSO Concentrate with Leci-Tech at a rate of 0.5 – 1 L/ha.
Application Timing	For Desi type, apply at the time swathing would normally commence, when the majority of plants are yellow and most pods are mature and seeds have turned from green to yellow or brown. Upper part of plant may still be green.  For Kabuli type, apply when the majority of plants and pods are ripe and dry with seeds turned from green to white or tan, and detached from the pods. Dry down is less complete in Kabuli type due to its thick pod wall.
Directions for Use – Ground	Ground application may be done with a standard boom sprayer.
Remarks	Do not graze or feed treated hay or straw to livestock. See the “ <b>HARVEST AID – CROP SPECIFIC RECOMMENDATIONS TANK-MIX</b> ” section for the detailed use pattern. Consult the label of the tank mix partner product for further use instructions, precautions and restrictions. The most restrictive labeling applies to tank mixes.

### DESICCATION – RED LENTIL VARIETIES

Timing	Desiccation of RED LENTIL VARIETIES
Rate	52 g/ha of <b>SPECIAL HERBICIDE</b> + 2.5 L/ha of glyphosate (360 g/L equivalent) + adjuvant
Water Volume	100 – 200 L/ha ground
Surfactant/Adjuvant	Add MERGE Adjuvant or MSO Concentrate with Leci-Tech at a rate of 0.5 – 1 L/ha.
Application Timing	Apply when lowermost pods (bottom 15%) are brown and rattle when shaken.
Directions for Use – Ground	Ground application may be done with a standard boom sprayer.

Remarks	Apply only to red lentil varieties. Do not graze or feed treated hay or straw to livestock. See the “ <b>HARVEST AID – CROP SPECIFIC RECOMMENDATIONS TANK-MIX</b> ” section for the detailed use pattern. Consult the label of the tank mix partner product for further use instructions, precautions and restrictions. The most restrictive labeling applies to tank mixes.
---------	---

### DESICCATION – FIELD PEAS

Timing	Desiccation of FIELD PEAS
Rate	52 g/ha of <b>SPECIAL HERBICIDE</b> + 2.5 L/ha of glyphosate (360 g/L equivalent) + adjuvant
Water Volume	100 – 200 L/ha ground
Surfactant/Adjuvant	Add MERGE Adjuvant or MSO Concentrate with Leci-Tech at a rate of 0.5 – 1 L/ha.
Application Timing	Apply when the majority of pods are brown (70-80%).
Directions for Use – Ground	Ground application may be done with a standard boom sprayer.
Remarks	See the “ <b>HARVEST AID – CROP SPECIFIC RECOMMENDATIONS TANK-MIX</b> ” section for the detailed use pattern. Consult the label of the tank mix partner product for further use instructions, precautions and restrictions. The most restrictive labeling applies to tank mixes.

### DESICCATION – CANOLA (ALL TYPES) AND MUSTARD<sup>1</sup>

Timing	Desiccation of CANOLA and MUSTARD
Rate	52 g/ha of <b>SPECIAL HERBICIDE</b> + 2.5 L/ha of glyphosate (360 g/L equivalent) + adjuvant
Water Volume	100 – 200 L/ha ground
Surfactant/Adjuvant	Add MERGE Adjuvant or MSO Concentrate with Leci-Tech at a rate of 0.5 – 1 L/ha.
Application Timing	Apply when 60 – 75% of seeds have changed colour.
Directions for Use – Ground	Ground application may be done with a standard boom sprayer.
Remarks	It is recommended that the application of <b>SPECIAL HERBICIDE</b> as a desiccant in canola and mustard be made to shatter resistant varieties. See the “ <b>HARVEST AID – CROP SPECIFIC RECOMMENDATIONS TANK-MIX</b> ” section for the detailed use pattern. Consult the label of the tank mix partner product for further use instructions, precautions and restrictions. The most restrictive labeling applies to tank mixes. Under ideal conditions, harvest can normally commence within 14-21 days after desiccation when applied at the appropriate crop stage recommendation. Adverse weather conditions such as rainfall, cool temperatures, shorter day length and high humidity may slow plant desiccation and keep seed moisture levels high which can delay commencement of harvest after the <b>SPECIAL HERBICIDE</b> application.

<sup>1</sup> All classes, including brown, oriental, canola quality *Brassica juncea*, *Brassica juncea* varieties with the **Clearfield** trait, and yellow mustard.

## DESICCATION – FLAX

Timing	Desiccation of FLAX
Rate	52 g/ha of <b>SPECIAL HERBICIDE</b> + 2.5 L/ha of glyphosate (360 g/L equivalent) + adjuvant
Water Volume	100 – 200 L/ha ground
Surfactant/Adjuvant	Add MERGE Adjuvant or MSO Concentrate with Leci-Tech at a rate of 0.5 – 1 L/ha.
Application Timing	Apply when 75% of bolls have turned colour.
Directions for Use – Ground	Ground application may be done with a standard boom sprayer.
Remarks	See the “ <b>HARVEST AID – CROP SPECIFIC RECOMMENDATIONS TANK-MIX</b> ” section for the detailed use pattern. Consult the label of the tank mix partner product for further use instructions, precautions and restrictions. The most restrictive labeling applies to tank mixes.

## DESICCATION – FABA BEANS

Timing	Desiccation of Faba Beans
Rate	52 g/ha of <b>SPECIAL HERBICIDE</b> + 2.5 L/ha of glyphosate (360 g/L equivalent) + adjuvant
Water Volume	100 – 200 L/ha ground
Surfactant/Adjuvant	Add MERGE Adjuvant or MSO Concentrate with Leci-Tech at a rate of 0.5 – 1 L/ha.
Application Timing	Apply when 80% of lower pods have turned black, middle pods have turned yellow/tan, and top green pods have firm seed.
Directions for Use – Ground	Ground application may be done with a standard boom sprayer.
Remarks	Do not graze or feed treated hay or straw to livestock. See the “ <b>HARVEST AID – CROP SPECIFIC RECOMMENDATIONS TANK-MIX</b> ” section for the detailed use pattern. Consult the label of the tank mix partner product for further use instructions, precautions and restrictions. The most restrictive labeling applies to tank mixes.

## PRE-HARVEST WEED MANAGEMENT – TANK MIX CROP SPECIFIC RECOMMENDATIONS

**SPECIAL HERBICIDE** may be used as a pre-harvest treatment in wheat (including durum, spring and winter wheat), barley (including feed varieties) and triticale to improve dry down of volunteer canola (all types including Roundup Ready), common ragweed, Canada fleabane, redroot pigweed and wild buckwheat. Tank mixing **SPECIAL HERBICIDE** with glyphosate will provide additional pre-harvest weed management to the crop dry down provided by glyphosate.

**The tank mix with glyphosate may affect seed germination. Tank mixing with glyphosate is not recommended when harvested grain is to be used for seed.**

## WHEAT AND BARLEY

Rate	36 – 52 g/ha of <b>SPECIAL HERBICIDE</b> + 2.5 L/ha of glyphosate (360 g/L equivalent) + adjuvant
Water Volume	100 – 200 L/ha ground

Surfactant/Adjuvant	Add MERGE Adjuvant or MSO Concentrate with Leci-Tech at a rate of 0.5 – 1 L/ha.
Application Timing	Hard dough stage; a thumbnail impression remains on seed; less than 30% moisture.
Weeds Controlled	Improved dry down of volunteer canola (all types including Roundup Ready), common ragweed, Canada fleabane, redroot pigweed and wild buckwheat. Consult the glyphosate label regarding additional pre-harvest weed control.
Directions for Use – Ground	Ground application may be done with a standard boom sprayer.
Remarks	Consult the label of the tank mix partner product for further use instructions, precautions and restrictions. The most restrictive labeling applies to tank mixes.  Use higher water volume for dense crop stands and higher weed pressure.  Tank mixing with glyphosate is not recommended when harvested grain is to be used for seed.

### MIXING INSTRUCTIONS

1. When using **SPECIAL HERBICIDE**, always start with a clean sprayer. Thoroughly clean the sprayer by flushing the system with water containing detergent. Refer to previously applied product labels for specific cleaning instructions.
2. Fill clean spray tank half full with clean water. Start agitation system.
3. Using a calibrated measuring device, add the correct amount of **SPECIAL HERBICIDE**. Continue to agitate.
4. **For tank mix with glyphosate**, add **SPECIAL HERBICIDE** followed by recommended amount of glyphosate.
5. If recommended, add the correct amount of MERGE Adjuvant or MSO Concentrate with Leci-Tech to the tank last.
6. Continue agitation while filling the remainder of the tank with water necessary to fill the spray tank.
7. Continue to agitate or run the by-pass system.
8. After any break in the spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to remix the spray materials. Do not allow the mixture to sit overnight.
9. If an oil film starts to build up in the tank, drain it and clean the tank with strong detergent solution.
10. Immediately after use, thoroughly clean the sprayer by flushing the system with clean water containing detergent.

Dispose of all rinsings in accordance with provincial/territorial regulations.

### PRE-HARVEST INTERVAL (PHI)

The following pre-harvest intervals should be observed for respective crops.

Crop	Pre-Harvest Interval (PHI) in days
Barley	60
Canary seed	60
Chickpea	60
Corn (field, sweet)	60
Faba beans	60

Lentils	60
Oats	60
Peas (dried field)	60
Soybean	60
Wheat (spring, winter, durum)	60

Creeping red fescue, timothy and bromegrass: forage and hay can be used as feed or grazed immediately after application of **SPECIAL HERBICIDE**.

### PRE-HARVEST INTERVAL (PHI) – HARVEST AID

The following pre-harvest intervals should be observed for respective crops when **SPECIAL HERBICIDE** is used as a harvest aid.

Crop	Pre-Harvest Interval (PHI) in days
Barley	3
Canola	3
Chickpeas	2
Dry common beans	2
Faba beans	2
Flax	3
Lentils	3
Mustard	3
Peas (dried field)	3
Soybeans	3
Sunflower	7
Triticale	3
Wheat	3

### FOLLOW CROPPING

The crops listed below can be safely grown after a spring application of **SPECIAL HERBICIDE**.

<b>Plant Back Crops</b> In case of crop failure, the following crops can be planted in the same season <sup>1</sup>	<b>Rotational Crops</b> The following crops can be planted anytime in the following season
Barley Canary seed Chickpeas Corn (field and sweet) Lentils* Oats Dry field peas Soybean* Wheat (spring, winter & durum)	Barley (spring, winter, malting) Canary seed Canola Chickpeas Corn (field and sweet) Dry common beans Flax Lentils Mustard Oats Dry field peas

	Soybean Triticale Wheat (spring, winter & durum)
--	--

\*Rate restrictions apply. Lentils and soybeans can only be grown as plant back crops provided that a maximum product use rate of 26 and 36 g/ha, respectively, was used in the previous crop.

<sup>1</sup> A second application of SPECIAL HERBICIDE cannot be made in the rescue crop.

The crops listed below can be safely grown after a fall application of **SPECIAL HERBICIDE**.

Rotational crops that can be planted in the following spring after application	Rotational crops that can be planted in the second spring after application
Barley Canary seed Canola Chickpeas Corn (field and sweet) Flax Lentils Oats Dry field peas Soybeans Wheat (spring, winter, and durum)	All crops

## SPRAYING INSTRUCTIONS

### Water volume and spray pressure Conventional ground application

Use sprayers equipped with standard flat nozzles. The use of 80°-110° stainless steel flat fan nozzles is recommended for optimum spray coverage with nozzles tilted 45° forward to ensure better coverage.

Thoroughly clean all screens to prevent nozzle clogging. Apply in a water volume of 50-100 L/ha and at a pressure of 240 kPa. For applications to dense weed infestations and thick canopies, use a higher water volume at pressures of 275 kPa.

Better coverage of the product results in enhanced control of weeds.

## RESTRICTIONS AND LIMITATIONS

1. Wash sprayer thoroughly after use to avoid damage to the next crop sprayed.
2. DO NOT APPLY USING AERIAL APPLICATION EQUIPMENT, unless specified otherwise.
3. **DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.
4. Field corn – Corn forage and silage can be harvested, used as feed or grazed 60 or more days after application of **SPECIAL HERBICIDE**.
5. Legume forage (chickpeas, faba beans, field peas and lentils) may be used as feed or grazed 60 or more days after application of **SPECIAL HERBICIDE**. Desiccation treated pea vines may be grazed or fed to livestock.
6. Small grains (wheat, barley and oats) – forage and hay can be used as feed or grazed 30 or more days after application of **SPECIAL HERBICIDE**. Pre-harvest treated barley, wheat and triticale straw may be grazed or fed to livestock.
7. Soybeans may be used as feed or grazed 60 or more days after application of **SPECIAL HERBICIDE**.

8. Soybeans, chickpeas and dry common beans – Do not graze or feed treated hay or straw to livestock when **SPECIAL HERBICIDE** is used as a harvest aid.
9. Grasses (creeping red fescue, timothy and bromegrass): forage and hay can be used as feed or grazed immediately after application of **SPECIAL HERBICIDE**.
10. This product may be tank mixed with (a fertilizer, a supplement, or with) registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions for Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Sharda Cropchem Limited at 1-844-810-5720 for information before applying any tank mix that is not specifically recommended on this label.

### **RESISTANCE-MANAGEMENT RECOMMENDATIONS**

For resistance management, **SPECIAL HERBICIDE** is a Group 14 herbicide. Any weed population may contain or develop plants naturally resistant to **SPECIAL HERBICIDE** and other Group 14 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of **SPECIAL HERBICIDE** or other Group 14 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.

- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Sharda Cropchem Limited at 1-844-810-5720.

All products listed are registered trademarks of their respective companies.