



EverGol[®] Rise

Seed Treatment Fungicide

GROUP	7, 11, 3, 4	FUNGICIDES
-------	-------------	------------

A Seed Treatment Fungicide for control or suppression of certain seed-borne and soil-borne diseases on labelled crops.

All seed treated with this product must be conspicuously coloured at the time of treatment. No open transfer is permitted in commercial seed treatment facilities and for mobile treaters. Only closed transfer, including closed mixing, loading, calibrating, and closed treatment equipment is permitted. For on-farm treatment open transfer including open mixing, loading, calibrating, and open treatment equipment is permitted.

SUSPENSION

COMMERCIAL (AGRICULTURAL)

ACTIVE INGREDIENTS:	Penflufen	50 g/L
	Trifloxystrobin	50 g/L
	Prothioconazole	50 g/L
	Metalaxyl	50 g/L

Contains 1,2 benzisothiazolin-3-one at 0.035% and 5-chloro-2-methyl-4-isothiazolin-3-one at 0.0009% and 2-methyl-4-isothiazolin-3-one at 0.0003% as preservatives.

REGISTRATION NO. 35641 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

PROTECT FROM FREEZING

NET CONTENTS: 0.946L - 1000L

Bayer CropScience Inc.
Suite 200, 160 Quarry Park Blvd. SE
Calgary AB T2C 3G3

For product information, call: 1-888-283-6847

In case of spills, poisoning, fire or other emergencies,
call: 1-800-334-7577 (24 hours a day)

EverGol® Rise Seed Treatment Fungicide is a seed treatment containing the fungicides penflufen, trifloxystrobin, prothioconazole and metalaxyl.

EverGol® Rise Seed Treatment Fungicide controls or suppresses certain seed, seedling and soil-borne diseases caused by *Rhizoctonia solani*, *Fusarium* spp., *Botrytis cinerea*, *Pythium* spp., *Ascochyta* spp., *Penicillium* spp. and *Phomopsis longicolla* on labelled crops.

DIRECTIONS FOR USE:

Refer to the attached booklet for specific use directions for each crop and detailed instructions on rates, diseases controlled, application timing and technique.

PRECAUTIONS:

- KEEP OUT OF REACH OF CHILDREN.
- Commercial seed treatment facilities must use closed transfer equipment only, including closed mixing, loading, calibrating, and treatment equipment. No open transfer of EverGol® Rise Seed Treatment Fungicide is permitted in commercial seed treatment facilities.
- Apply only in a way that this product will not contact workers or other persons either directly or through drift. Only workers wearing personal protective equipment may be in the area where treating, bagging, sewing and/or stacking is occurring.
- Use good personal hygiene, washing hands and exposed skin with warm water and soap before eating, drinking, or smoking. No food, drink or tobacco should be allowed in areas of chemical storage or use.
- Follow the personal protective equipment (PPE) and restriction requirements for the appropriate task and scenario:

Seed Type	Tasks	PPE/Engineering Controls
For Commercial Seed Treatment (including facility workers and mobile treaters)		
All Listed Seed Types	Mixing, loading, treating, and calibrating	Closed transfer only. Closed transfer includes closed mixing, loading, calibrating and closed treatment. No open transfer of EverGol® Rise Seed Treatment Fungicide is permitted. Wear a long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves.
	Clean-up and repair, or other maintenance of seed treatment equipment	Wear coveralls over a long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves.
	Bagging sewing or stacking of bagged treated seed, or when transferring treated seed to a storage bin.	Wear a long-sleeved shirt, long pants, a suitable NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested, socks, shoes and chemical-resistant gloves.
For On-Farm Seed Treatment (including handling and planting of seed treated on-farm)		
All Listed Seed Types	Mixing, loading, treating, calibrating, cleaning, and	Open or closed transfer system is permitted

	repairing.	Wear a long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves.
	Activities involving handling and planting seeds treated on-farm	Wear a long-sleeved shirt, long pants, a suitable NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested, socks, shoes and chemical-resistant gloves. Gloves and dust mask are not required within a closed cab.
For Planting and Handling Commercially Treated Seed		
All Listed Seed Types	Handling and planting treated seed	Wear a long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat patient symptomatically. This product contains petroleum distillates. Vomiting may cause aspiration pneumonia.

ENVIRONMENTAL PRECAUTIONS:

Toxic to aquatic organisms.

Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned up from the soil surface.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

STORAGE:

Store this product away from food or feed. Store in a cool, dry area. Do not store in direct sunlight. Do not allow prolonged storage in temperatures that exceed 40°C or go below -10°C.

DISPOSAL:

FOR RECYCLABLE CONTAINERS:

Disposal of container: DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

FOR RETURNABLE CONTAINERS:

Disposal of container: DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

FOR REFILLABLE CONTAINERS:

Disposal of container: For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT use this container for any other purpose.

TRIPLE RINSE INSTRUCTIONS (FOR RECYCLABLE CONTAINERS ONLY): This container should be triple-rinsed prior to its disposal. The rinsate can be added to the seed treatment. Follow these steps for rinsing of containers:

1. Transfer EverGol[®] Rise Seed Treatment Fungicide from the container to the seed treating equipment applicator tank.
2. Add 1/3 of the rinse water to the container and agitate the contents.
3. Transfer the rinsate to the applicator tank.
4. Rinse as above 2 more times.
5. Thoroughly mix the EverGol[®] Rise Seed Treatment Fungicide and rinse water in the applicator tank prior to use.

CAUTION: Be sure to adjust the undiluted application rate by the same volume of water added to compensate for the dilution of the product caused by the addition of rinsate.

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for cleanup of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.



EverGol[®] Rise

Seed Treatment Fungicide

GROUP	7, 11, 3, 4	FUNGICIDES
-------	-------------	------------

A Seed Treatment Fungicide for control or suppression of certain seed-borne and soil-borne diseases on labelled crops.

All seed treated with this product must be conspicuously coloured at the time of treatment. No open transfer is permitted in commercial seed treatment facilities and for mobile treaters. Only closed transfer, including closed mixing, loading, calibrating, and closed treatment equipment is permitted. For on-farm treatment open transfer including open mixing, loading, calibrating, and open treatment equipment is permitted.

SUSPENSION

COMMERCIAL (AGRICULTURAL)

ACTIVE INGREDIENTS:	Penflufen	50 g/L
	Trifloxystrobin	50 g/L
	Prothioconazole	50 g/L
	Metalaxyl	50 g/L

Contains 1,2 benzisothiazolin-3-one at 0.035% and 5-chloro-2-methyl-4-isothiazolin-3-one at 0.0009% and 2-methyl-4-isothiazolin-3-one at 0.0003% as preservatives.

REGISTRATION NO. 35641 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

PROTECT FROM FREEZING

NET CONTENTS: 0.946L - 1000L

Bayer CropScience Inc.
Suite 200, 160 Quarry Park Blvd. SE
Calgary AB T2C 3G3

For product information, call: 1-888-283-6847

In case of spills, poisoning, fire or other emergencies,
call: 1-800-334-7577 (24 hours a day)

Table of Contents:	Section Number
GENERAL INFORMATION	The Product 1
SAFETY AND HANDLING	Precautions, Protective Clothing and Equipment..... 2
	First Aid and Toxicological Information 3
	Environmental Precautions 4
	Storage..... 5
	Disposal..... 6
	Notices..... 7
DIRECTIONS FOR USE	Diseases Controlled 8
	Recommended Uses 9
	Application Cautions..... 10
	Use Limitations 11
	Resistance Management Recommendations 12

GENERAL INFORMATION

Section 1: The Product

EverGol[®] Rise Seed Treatment Fungicide is a seed treatment containing the fungicides penflufen, trifloxystrobin, prothioconazole and metalaxyl. EverGol[®] Rise Seed Treatment Fungicide controls or suppresses certain seed, seedling and soil-borne diseases caused by *Rhizoctonia solani*, *Fusarium* spp., *Botrytis cinerea*, *Pythium* spp., *Ascochyta* spp., *Penicillium* spp. and *Phomopsis longicolla* on labelled crops.

SAFETY AND HANDLING

Section 2: Precautions, Protective Clothing and Equipment

PRECAUTIONS:

- KEEP OUT OF REACH OF CHILDREN.
- Commercial seed treatment facilities must use closed transfer equipment only including closed mixing, loading, calibrating, and treatment equipment. No open transfer of EverGol[®] Rise Seed Treatment Fungicide is permitted in commercial seed treatment facilities.
- Apply only in a way that this product will not contact workers or other persons either directly or through drift. Only workers wearing personal protective equipment may be in the area where treating, bagging, sewing and/or stacking is occurring.
- Use good personal hygiene, washing hands and exposed skin with warm water and soap before eating, drinking, or smoking. No food, drink or tobacco should be allowed in areas of chemical storage or use.
- Follow the personal protective equipment (PPE) and restriction requirements for the appropriate task and scenario:

Seed Type	Tasks	PPE/Engineering Controls
For Commercial Seed Treatment (including facility workers and mobile treaters)		
All Listed Seed Types	Mixing, loading, treating, and calibrating	Closed transfer only. Closed transfer includes closed mixing, loading, calibrating and closed treatment. No open transfer of EverGol [®] Rise Seed Treatment Fungicide is permitted. Wear a long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves.
	Clean-up and repair, or other maintenance of seed treatment equipment	Wear coveralls over a long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves.
	Bagging sewing or stacking of bagged treated seed, or when transferring treated seed to a storage bin.	Wear a long-sleeved shirt, long pants, a suitable NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested,

		socks, shoes and chemical-resistant gloves.
For On-Farm Seed Treatment (including handling and planting of seed treated on-farm)		
All Listed Seed Types	Mixing, loading, treating, calibrating, cleaning, and repairing.	Open or closed transfer system is permitted Wear a long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves.
	Activities involving handling and planting seeds treated on-farm	Wear a long-sleeved shirt, long pants, a suitable NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested, socks, shoes and chemical-resistant gloves. Gloves and dust mask are not required within a closed cab.
For Planting and Handling Commercially Treated Seed		
All Listed Seed Types	Handling and planting treated seed	Wear a long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves

Section 3: First Aid and Toxicological Information

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat patient symptomatically. This product contains petroleum distillates. Vomiting may cause aspiration pneumonia.

Section 4: Environmental Precautions

ENVIRONMENTAL PRECAUTIONS:

Toxic to aquatic organisms.

Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Section 5: Storage

Store this product away from food or feed. Store in a cool, dry area. Do not store in direct sunlight. Do not allow prolonged storage in temperatures that exceed 40°C or go below -10°C.

Section 6: Disposal

FOR RECYCLABLE CONTAINERS:

Disposal of container: DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

FOR RETURNABLE CONTAINERS:

Disposal of container: DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

FOR REFILLABLE CONTAINERS:

Disposal of container: For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT use this container for any other purpose.

TRIPLE RINSE INSTRUCTIONS (FOR RECYCLABLE CONTAINERS ONLY): This container should be triple-rinsed prior to its disposal. The rinsate can be added to the seed treatment. Follow these steps for rinsing of containers:

1. Transfer EverGol® Rise Seed Treatment Fungicide from the container to the seed treating equipment applicator tank.
2. Add 1/3 of the rinse water to the container and agitate the contents.
3. Transfer the rinsate to the applicator tank.

4. Rinse as above 2 more times.
5. Thoroughly mix the EverGol[®] Rise Seed Treatment Fungicide and rinse water in the applicator tank prior to use.

CAUTION: Be sure to adjust the undiluted application rate by the same volume of water added to compensate for the dilution of the product caused by the addition of rinsate.

DISPOSAL OF UNUSED, UNWANTED PRODUCT:

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for cleanup of spills.

Section 7: Notices

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

DIRECTIONS FOR USE

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Section 8: Diseases Controlled

EverGol[®] Rise Seed Treatment Fungicide is a seed treatment containing the fungicides penflufen, trifloxystrobin, prothioconazole and metalaxyl. EverGol[®] Rise Seed Treatment Fungicide controls or suppresses certain diseases caused by *Rhizoctonia solani*, *Fusarium* spp., *Botrytis cinerea*, *Pythium* spp., *Ascochyta* spp., *Penicillium* spp. and *Phomopsis longicolla* on labelled crops.

Section 9: Recommended Uses

EverGol[®] Rise Seed Treatment Fungicide is a seed treatment formulation for use in commercial seed treatment operations, and for on-farm treating with conventional seed treating equipment which can accurately meter and apply flowable seed treatment formulations.

This product is recommended to be diluted with water or another suitable liquid just prior to application to ensure uniform coverage on the seed during the application process. Uniform application to seed is necessary to ensure optimum performance. Allow seeds to dry before bagging, storing or seeding.

CROP	DISEASE	APPLICATION RATE	DIRECTIONS
<p>Crop Subgroup 6-21E (Pulses, dried shelled beans, except soybeans): African yam-bean, dry seed; American potato bean, dry seed; Bean (<i>Lupinus</i> spp.), dry seed; Bean (<i>Vigna</i> spp.), dry seed; Broad bean (fava bean), dry seed; Goa bean, dry seed; Guar bean, dry seed; Horse gram, dry seed; Jackbean, dry seed; Lablab bean, dry seed; Morama bean, dry seed; Sword bean, dry seed; Velvet bean, dry seed; Winged pea, dry seed; cultivars, varieties, and/or hybrids of these commodities.</p> <p>Crop Subgroup 6-21F (Pulses, dried shelled peas): Chickpea, dry seed; Grass-pea, dry seed; Lentil, dry seed; Pea (<i>Pisum</i> spp.), dry seed; Pigeon pea, dry seed; cultivars, varieties, and/or hybrids of these commodities.</p>	<p>Control of:</p> <p>Seed decay / pre-emergence damping-off and post-emergence damping-off caused by <i>Rhizoctonia solani</i>, <i>Fusarium</i> spp. and <i>Pythium</i> spp.</p> <p>Seed decay / pre-emergence damping-off, post-emergence damping-off and seedling blight caused by seed-borne <i>Botrytis cinerea</i>.</p> <p>Seed rot and pre-emergence damping-off caused by <i>Ascochyta</i> spp. on lentil, field pea and chickpea only</p> <p>Suppression of:</p> <p>Seed-borne ascochyta blight caused by <i>Ascochyta</i> spp.</p>	<p>Apply 100 mL per 100 kg of seed</p>	<p>EverGol® Rise Seed Treatment Fungicide is designed for commercial or on-farm treating with conventional seed treating equipment which can accurately control application rates and provide a good distribution of the chemical onto the seed in the mixing chamber.</p> <p>Uniform application to seed is necessary to ensure optimum product performance.</p>
<p>Soybean</p>	<p>Control of:</p> <p>Seed rot / pre-emergence damping-off and post-emergence damping-off caused by <i>Rhizoctonia solani</i>, <i>Fusarium</i> spp., and <i>Pythium</i> spp.</p> <p>Early-season root rot and seedling blight caused by <i>Rhizoctonia solani</i> and <i>Fusarium</i> spp.</p> <p>Seedling blight caused by seed-borne <i>Botrytis cinerea</i>.</p> <p>Seed rot / pre-emergence damping-off of soybean caused by <i>Phomopsis longicolla</i></p> <p>Suppression of seed infection caused by <i>Penicillium</i> spp.</p>	<p>Apply 100 mL per 100 kg of seed</p>	

TANK-MIXTURES

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions and Environmental Precautions are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Bayer CropScience Inc. at 1-888-283-6847 or www.cropscience.bayer.ca for information before applying any tank mix that is not specifically recommended on this label.

Bayer CropScience Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) of all potential tank mixes under all environmental conditions or for all crop varieties. Conduct a germination test on a small portion of seed before committing the total seed lot to tank mixes that are not specifically listed on this label.

To determine the physical compatibility of this product with other products, use a jar test. Always read and follow label directions including WAMLEGS mixing order.

For control of certain insect pests and/or additional diseases not on the EverGol[®] Rise Seed Treatment Fungicide product label, EverGol[®] Rise Seed Treatment Fungicide may be combined with certain seed treatment insecticides and fungicides. Refer to the registered label of the tank-mix partner(s) for application rates, precautions and directions for use associated with those products, and follow the most restrictive label precautions and limitations.

CROP	TANK-MIX PARTNERS
Pea (field) Chickpea Lentil Bean (dry)	Stress Shield 600
Soybean	Stress Shield 600
	Allegiance FL or Acceleron D-309 (35 mL per 100kg of seeds) ¹
	Acceleron I-425, Acceleron I-374

¹ For control of *Phytophthora sojae* in soybean, the recommended rate of EverGol[®] Rise Seed Treatment Fungicide may be tank-mixed with 35 mL of Allegiance FL or Acceleron D-309 per 100 kg seed.

Section 10: Application Cautions

COLOURATION OF TREATED SEED: This product contains a pigment which will colour the treated seed. However, users are responsible for ensuring that the treated seed, when dried and ready for bagging, storage or seeding, has an unnatural colour. If the pigment contained in the

formulation does not colour the seed adequately, additional colourant must be added to the mixture while treating the seed. Regulations pertaining to colouration of treated seed enforced under the *Seeds Act* must be strictly adhered to when using this product.

SLURRY PREPARATION:

When preparing a slurry of EverGol® Rise Seed Treatment Fungicide with dilutants such as water or liquid inoculants, additional or continuous agitation or mixing may be necessary to prevent EverGol® Rise Seed Treatment Fungicide from settling out. Do not store diluted slurry mixtures.

SEED TREATMENT AND INOCULANTS:

Please check with inoculant manufacturers for application and compatibility details prior to use.

Seeds treated with EverGol® Rise Seed Treatment Fungicide and seed inoculants may not flow through seeding equipment at the same rate as untreated seeds. Recalibrate the seeding equipment prior to planting treated seeds. Mixing EverGol® Rise Seed Treatment Fungicide with seed inoculants may increase drying time while treating and extend the processing time.

SEED QUALITY:

Lab and field studies have shown that EverGol® Rise Seed Treatment Fungicide treated seed can be stored without loss in germination or disease performance. Many large-seed legumes are prone to mechanical damage during seed handling, and a reduction in germination within a seed lot can occur should handling cause seed damage irrespective of seed treatment application. EverGol® Rise Seed Treatment Fungicide treated or untreated seed may drop in germination with age and germination drops are accelerated under adverse storage conditions or in seed with a high level of mechanical damage. Use handling equipment designed to minimize mechanical damage when moving seed. Due to seed quality and seed storage conditions beyond the control of Bayer CropScience, no claims are made to guarantee the germination of carry-over seed or propagating materials for all crop seed. Treatment of highly mechanically scarred or damaged seed, or seed known to be of low vigour and poor quality, may result in reduced germination and/or reduction of seed and seedling vigour. Treat a small quantity of seed using equipment similar to that planned for treating the total seed lot. Conduct germination tests on a small portion of seed before committing the total seed lot to a selected seed treatment.

Section 11: Use Limitations

USE RESTRICTIONS:

- Treated seed must not be used for food, feed or oil processing.
- Do not graze or feed livestock on treated areas for four weeks after planting.

LABELLING TREATED SEED:

All containers or packages containing treated seed for sale or use in Canada must be labelled or tagged as follows: “This seed has been treated with EverGol® Rise Seed Treatment Fungicide, which contains penflufen, prothioconazole, metalaxyl and trifloxystrobin. Toxic to aquatic organisms. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned up from the soil surface. Dispose of all excess treated seed. Leftover treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. Wear long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seed. DO

NOT use for feed, food or oil processing. Store away from feeds and other foodstuffs.” Registered crops for EverGol[®] Rise Seed Treatment Fungicide, as well as canola, mustard, rapeseed, soybean, alfalfa, corn and cereal grains, may be replanted at any time. For all other crops, do not plant-back within 30 days of seeding with EverGol[®] Rise Seed Treatment Fungicide treated seed. Do not graze or feed livestock on treated areas for four weeks after planting.

ROTATIONAL CROPS:

Registered crops for EverGol[®] Rise Seed Treatment Fungicide, as well as canola, mustard, rapeseed, soybean, alfalfa, corn and cereal grains, may be replanted at any time. For all other crops, do not plant-back within 30 days of seeding with EverGol[®] Rise Seed Treatment Fungicide treated seed.

Section 12: Resistance Management Recommendations

For resistance management, please note that EverGol[®] Rise Seed Treatment Fungicide contains a Group 7,11, 3 and 4 fungicide. Any fungal population may contain individuals naturally resistant to EverGol[®] Rise Seed Treatment Fungicide and other Group 7, 11, 3 and 4 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

- Where possible, rotate the use of EverGol[®] Rise Seed Treatment Fungicide or other Group 7,11, 3 and 4 fungicides with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group when such use is permitted.
- Fungicide use should be based on an IPM program that includes scouting, historical information related to pesticide use and crop rotation and considers cultural, biological and other chemical control practices.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Bayer CropScience at 1-888-283-6847 or at www.bayercropscience.ca.