

2025-5537

2025-12-03

Container Label

GROUPS

3

4

FUNGICIDES

TARIAN™ MD SEED TREATMENT FUNGICIDE

SYSTEMIC FUNGICIDE SEED PROTECTANT FOR WHEAT, BARLEY, OATS, RYE AND TRITICALE

TARIAN MD SEED TREATMENT FUNGICIDE is a combination of the systemic fungicides tebuconazole and metalaxyl which control certain seed, seedlings, and soil borne diseases of wheat, barley, oats, rye and triticale.

COMMERCIAL (AGRICULTURAL)

SUSPENSION

ACTIVE INGREDIENTS: Tebuconazole..... 5.0 g/L
Metalaxyl..... 6.6 g/L

REGISTRATION NO. 35666 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND BOOKLET BEFORE USING

PROTECT FROM FREEZING

NET CONTENTS: 1 L - BULK

For wheat, barley, rye, triticale, and oats: On-farm or commercial seed treatment (facilities and mobile treaters) with open or closed transfer treatment equipment is permitted. Apply only in a way that this product will not contact workers or other persons, either directly or through drift. Only workers wearing personal protective equipment may be in the area when seed is being treated or bagged.

For Hazardous Materials [or Dangerous Goods] Incident Spill, Leak, Fire, Exposure, or Accident Call CHEMTREC 1-800-424-9300 / +1 703-527-3887

TARIAN™ is a trademark of NewAgco Inc. or its affiliates.

2025-5537

2025-12-03

LABELLING OF TREATED SEED:

All bags containing treated seed for sale or use in Canada must be labeled or tagged as follows:

- **KEEP TREATED SEED OUT OF REACH OF CHILDREN AND ANIMALS**
- This seed has been treated with **TARIAN MD SEED TREATMENT FUNGICIDE** containing tebuconazole and metalaxyl.
- During handling and planting treated seed, wear coveralls over a long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves. During handling of treated seeds, wear a NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested. Use a closed-cab tractor when planting. Gloves and respiratory protection are not required with a closed cab.
- **DO NOT** use for food, feed or oil processing.
- **DO NOT** graze or feed livestock on treated areas for 4 weeks after planting.
- **DO NOT** contaminate feed or foodstuffs with treated seed.
- Store this product away from food or feed
- Toxic to aquatic organisms. **DO NOT** expose treated seeds on soil surface. Any spilled or exposed seed must be incorporated into the soil or otherwise cleaned-up from the soil surface.
- Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements.
- Dispose of seed packaging in accordance with local requirements. **DO NOT** re-use bags from treated seed to handle food or feed products.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN AND ANIMALS. Wash hands thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Apply only in a way that this product will not contact workers or other persons, either directly or through drift. Only workers wearing personal protective equipment may be in the area where treating, bagging, sewing and/or stacking is occurring.

Follow the personal protective equipment (PPE) and restriction requirements for the appropriate task and scenario:

Seed Types	Tasks	Personal Protective Equipment/Engineering Controls
For commercial seed treatment (including facility workers and mobile treaters)		
Wheat, Barley, Oats, Rye and Triticale	Mixing, loading, treating, calibrating, bagging, sewing, stacking, and any other activities involving handling treated seed.	Open or closed transfer. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes.
	Cleaning and repairing	Wear chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes
For on-farm treatment (including handling and planting of seed treated on-farm)		
Wheat, Barley, Oats, Rye and Triticale	Mixing, loading, treating, calibrating, cleaning, repairing and any other activities involving handling and planting treated seeds on-farm.	Open or closed transfer. Wear a long-sleeved shirt, long pants, shoes and chemical-resistant gloves. Use a closed-cab tractor when planting. Gloves are not required with a closed cab.
For planting commercially-treated seeds (also include on seed tags)		

2025-5537

2025-12-03

Wheat, Barley, Oats, Rye and Triticale	Handling and planting	Wear coveralls over a long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves. During handling of treated seeds, wear a NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested. Use a closed-cab tractor when planting. Gloves and respiratory protection are not required with a closed cab.
--	-----------------------	--

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product.
- Wash the outside of gloves before removing.
- As soon as possible, wash thoroughly and change into clean clothing.

Work in a well-ventilated area when treating seed or while augering or handling treated seed. Augers used for handling treated seed should not be used to move seed for feed, food or oil processing.

If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact NewAgco Inc. Canada Inc. at 1-866-639-2420 or www.newagco.com.

FIRST AID:

IF SWALLOWED: Call a Poison Control Centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Centre or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

This compound does not cause any definite symptoms that would be diagnostic. Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS

- Toxic to aquatic organisms and non-target terrestrial plants.

2025-5537

2025-12-03

- Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface.
- This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

STORAGE:

Store this product away from food or feed. Store product in original container only, away from other pesticides and fertilizer. Store in a cool, dry place and avoid excessive heat. Keep container closed.

DISPOSAL:

FOR RECYCLABLE CONTAINERS: DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

FOR RETURNABLE CONTAINERS: DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

FOR REFILLABLE CONTAINERS: For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

DISPOSAL OF UNUSED, UNWANTED PRODUCT: For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for cleanup of spills.

DISPOSAL OF TREATED SEED: Dispose of all excess treated seed. Leftover treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. Dispose of seed packaging in accordance with local requirements. DO NOT re-use bags from treated seed to handle food or feed products.

NOTICE: READ CAREFULLY

Treatment of highly mechanically damaged seed or seed of known low vigor may result in reduced germination and/or reduction of seedling vigor.

When seed quality is questionable or unknown, conducting a germination test on a small portion of treated seed is recommended before treating the whole seed lot.

Due to seed quality and storage conditions which are beyond the control of NewAgco Inc., Inc., no claims are made to guarantee germination of seed which has been treated and kept longer than the period, if any that is stated in the DIRECTIONS FOR USE section of this label.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

GROUPS	3	4	FUNGICIDES
--------	---	---	------------

TARIAN™ MD SEED TREATMENT FUNGICIDE

SYSTEMIC FUNGICIDE SEED PROTECTANT FOR WHEAT, BARLEY, OATS, RYE AND TRITICALE

TARIAN MD SEED TREATMENT FUNGICIDE is a combination of the systemic fungicides tebuconazole and metalaxyl which control certain seed, seedling, and soil borne diseases of wheat, barley, oats, rye and triticale.

COMMERCIAL (AGRICULTURAL)

SUSPENSION

ACTIVE INGREDIENTS: tebuconazole..... 5.0 g/L
 metalaxyl..... 6.6 g/L

REGISTRATION NO. 35666 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND BOOKLET BEFORE USING

PROTECT FROM FREEZING

NET CONTENTS: 1 L - BULK

For wheat, barley, rye, triticale, and oats: On-farm or commercial seed treatment (facilities and mobile treaters) with open or closed transfer treatment equipment is permitted. Apply only in a way that this product will not contact workers or other persons, either directly or through drift. Only workers wearing personal protective equipment may be in the area when seed is being treated or bagged.

For Hazardous Materials [or Dangerous Goods] Incident Spill, Leak, Fire, Exposure, or Accident Call CHEMTREC 1-800-424-9300 / +1 703-527-3887

TARIAN™ is a trademark of NewAgco Inc. or its affiliates.

LABELLING OF TREATED SEED:

All bags containing treated seed for sale or use in Canada must be labeled or tagged as follows:

- This seed has been treated with **TARIAN MD SEED TREATMENT FUNGICIDE** containing tebuconazole and metalaxyl.
- Wear coveralls or a long-sleeved shirt and long pants, NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask), shoes plus socks and chemical resistant gloves when handling treated seed, stacking bags or transferring to storage bin.
- Do not use for food, feed or oil processing.
- Do not graze or feed livestock on treated areas for 4 weeks after planting.
- Do not contaminate feed or foodstuffs with treated seed.
- Toxic to aquatic organisms. Do not expose treated seeds on soil surface. Any spilled or exposed seed must be incorporated into the soil or otherwise cleaned-up from the soil surface.
- Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements.
- Dispose of seed packaging in accordance with local requirements. DO NOT re-use bags from treated seed to handle food or feed products.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN AND ANIMALS. Wash hands thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Apply only in a way that this product will not contact workers or other persons, either directly or through drift. Only workers wearing personal protective equipment may be in the area where treating, bagging, sewing and/or stacking is occurring.

Follow the personal protective equipment (PPE) and restriction requirements for the appropriate task and scenario:

Seed Types	Tasks	Personal Protective Equipment/Engineering Controls
For commercial seed treatment (including facility workers and mobile treaters)		
Wheat, Barley, Oats, Rye and Triticale	Mixing, loading, treating, calibrating, bagging, sewing, stacking, and any other activities involving handling treated seed.	Open or closed transfer. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes.
	Cleaning and repairing	Wear chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes
For on-farm treatment (including handling and planting of seed treated on-farm)		
Wheat, Barley, Oats, Rye and Triticale	Mixing, loading, treating, calibrating, cleaning, repairing and any other activities involving handling and planting treated seeds on-farm.	Open or closed transfer. Wear a long-sleeved shirt, long pants, shoes and chemical-resistant gloves. Use a closed-cab tractor when planting. Gloves are not required with a closed cab.
For planting commercially-treated seeds (also include on seed tags)		

2025-5537

2025-12-03

Wheat, Barley, Oats, Rye and Triticale	Handling and planting	Wear coveralls over a long-sleeved shirt, long pants, socks, shoes and chemical-resistant gloves. During handling of treated seeds, wear a NIOSH-approved N95 (minimum) filtering facepiece respirator (dust mask) that is properly fit tested. Use a closed-cab tractor when planting. Gloves and respiratory protection are not required with a closed cab.
--	-----------------------	--

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product.
- Wash the outside of gloves before removing.
- As soon as possible, wash thoroughly and change into clean clothing.

Work in a well-ventilated area when treating seed or while augering or handling treated seed. Augers used for handling treated seed should not be used to move seed for feed, food or oil processing.

If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact NewAgco Inc. Canada Inc. at 1-866-639-2420 or www.newagco.com.

FIRST AID:

IF SWALLOWED: Call a Poison Control Centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a Poison Control Centre or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

This compound does not cause any definite symptoms that would be diagnostic. Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS

- Toxic to aquatic organisms and non-target terrestrial plants.

2025-5537

2025-12-03

- Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface.
- This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

STORAGE:

Store this product away from food or feed. Store product in original container only, away from other pesticides and fertilizer. Store in a cool, dry place and avoid excessive heat. Keep container closed.

DISPOSAL:

FOR RECYCLABLE CONTAINERS:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

FOR RETURNABLE CONTAINERS:

DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

FOR REFILLABLE CONTAINERS:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

**TARIAN MD SEED TREATMENT FUNGICIDE TARIAN MD SEED TREATMENT FUNGICIDE
DISPOSAL OF UNUSED, UNWANTED PRODUCT:**

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for cleanup of spills.

DISPOSAL OF TREATED SEED:

- Dispose of all excess treated seed. Leftover treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements.
- Dispose of seed packaging in accordance with local requirements. DO NOT re-use bags from treated seed to handle food or feed products.

NOTICE: READ CAREFULLY

Treatment of highly mechanically damaged seed or seed of known low vigor may result in reduced germination and/or reduction of seedling vigor.

When seed quality is questionable or unknown, conducting a germination test on a small portion of treated seed is recommended before treating the whole seed lot.

Due to seed quality and storage conditions which are beyond the control of NewAgco Inc., Inc., no claims are made to guarantee germination of seed which has been treated and kept longer than the period, if any that is stated in the DIRECTIONS FOR USE section of this label.

2025-5537

2025-12-03

GENERAL INFORMATION: TARIAN MD SEED TREATMENT FUNGICIDE is a combination of the systemic fungicides tebuconazole and metalaxyl, which control or suppress the following seed, seedling, and soil borne diseases of wheat, barley, oats, rye and triticale.

Crop	Diseases
Wheat, Rye, Triticale	Diseases Controlled: Loose smut (<i>Ustilago tritici</i>) Common bunt or stinking smut (<i>Tilletia tritici</i> , <i>Tilletia laevis</i>) Seed rot and pre-emergent damping-off caused by seed- and soil-borne <i>Fusarium spp.</i> Seedling blight caused by seed-borne <i>Fusarium spp.</i> Damping-off caused by <i>Pythium spp.</i> Seed-borne <i>Septoria nodorum</i>
	Diseases Suppressed: Root and Crown rot caused by seed- and soil-borne <i>Fusarium spp.</i> Common root rot caused by seed- and soil-borne <i>Cochliobolus sativus</i> Seed rot and pre-emergent damping-off caused by seed- and soil-borne <i>Cochliobolus sativus</i> Seedling blight caused by seed-borne <i>Cochliobolus sativus</i>
Barley	Diseases Controlled: True loose smut (<i>Ustilago nuda</i>) Covered smut (<i>Ustilago hordei</i>) False loose smut (<i>Ustilago nigra</i>) Seed rot and pre-emergent damping-off caused by seed- and soil-borne <i>Fusarium spp.</i> Seedling blight caused by seed-borne <i>Fusarium spp.</i> Damping-off caused by <i>Pythium spp.</i>
	Diseases Suppressed: Root and Crown rot caused by seed- and soil-borne <i>Fusarium</i> species Common root rot caused by seed- and soil-borne <i>Cochliobolus sativus</i> Seed rot and pre-emergent damping-off caused by seed- and soil-borne <i>Cochliobolus sativus</i> Seedling blight caused by seed-borne <i>Cochliobolus sativus</i>
Oats	Diseases Controlled: Covered smut (<i>Ustilago kollerii</i>) Loose smut (<i>Ustilago avenae</i>) Seed rot and pre-emergent damping-off caused by seed- and soil-borne <i>Fusarium spp.</i> Seedling blight caused by seed-borne <i>Fusarium spp.</i> Damping-off caused by <i>Pythium spp.</i>
	Diseases Suppressed: Root and Crown rot caused by seed- and soil-borne <i>Fusarium</i> species Common root rot caused by seed- and soil-borne <i>Cochliobolus sativus</i> Seed rot and pre-emergent damping-off caused by seed- and soil-borne <i>Cochliobolus sativus</i> Seedling blight caused by seed-borne <i>Cochliobolus sativus</i>

DIRECTIONS FOR USE:

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

WHEAT, BARLEY, OATS, RYE AND TRITICALE:

TARIAN MD SEED TREATMENT FUNGICIDE is a ready to use formulation designed for commercial or on-farm treating with conventional seed treating equipment which can accurately control application rates and provide good distribution of the chemical onto the seed in the mixing chamber. The formulation contains a colouring agent which will colour the treated seed. Uniform application to seed is necessary to ensure seed safety and best disease control.

Use 300 ml of **TARIAN MD SEED TREATMENT FUNGICIDE** per 100 kg of seed.

1. Seed is protected by **TARIAN MD SEED TREATMENT FUNGICIDE** as soon as it is treated. Seed may be planted immediately.
2. Calibrate seeding equipment using treated seed to insure proper seeding rate.
3. Germination will not be affected by treatment as long as over application does not occur and seed is properly stored. All seed carried from one year to the next, whether treated or untreated should be checked for germination before planting.
4. Uniform coverage at the correct rate is important for satisfactory results. Under treatment will lead to loss of efficacy and over treatment could reduce germination. Refer to instructions supplied with your seed treater or to information supplied by NewAgco Inc. for assistance with proper treating.

TANK MIXES

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label. In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact NewAgco Inc. at 1-866-639-2420 for information before applying any tank mix that is not specifically recommended on this label.

When tank mixes are permitted, read and observe all label directions, including rates and restrictions for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.

RESISTANCE MANAGEMENT RECOMMENDATIONS:

For resistance management, please note that **TARIAN MD SEED TREATMENT FUNGICIDE** contains both Group 3 and Group 4 fungicides. Any fungal population may contain individuals naturally resistant to **TARIAN MD SEED TREATMENT FUNGICIDE** and other Group 3 or Group 4 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

- Where possible, rotate the use of **TARIAN MD SEED TREATMENT FUNGICIDE** or other Group 3, 4 fungicides with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that is effective on the target pathogen when such use is permitted.
- Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications.

2025-5537

2025-12-03

- Monitor treated fungal populations for resistance development. Notify NewAgco Inc. if reduced sensitivity of the pathogen to **TARIAN MD SEED TREATMENT FUNGICIDE** is suspected.
- If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information and to report suspected resistance contact NewAgco Inc. at 1-866-639-2420.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

TARIAN™ is a trademark of NewAgco Inc. or its affiliates.

All other products mentioned are trademarks of their respective holders.