

2025-4911
2025-10-31

| | | |
|--------------|----------|------------------|
| GROUP | 5 | HERBICIDE |
|--------------|----------|------------------|

CANTERRA 480 SC

SUSPENSION

HERBICIDE

COMMERCIAL (AGRICULTURAL)

Liquid herbicide for selective weed control in corn, sorghum and switchgrass

ACTIVE INGREDIENT:

Atrazine and related triazines 480 g/L

Contains 1,2-benzisothiazolin-3-one at 0.017% as a preservative.

**READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF THE REACH OF CHILDREN**

REGISTRATION NO.: 35706
PEST CONTROL PRODUCTS ACT

NET CONTENTS: **0.5 L – 250 L**

Sharda Cropchem Limited
2nd Floor, Prime Business Park, Dashrathlal Joshi Road,
VileParle (west), Mumbai - 400056
India

Canadian Agent:
Sharda Cropchem Limited
601 - 402 - 21st Street East
Saskatoon, SK, S7K 0C3
[1-888-931-2530](tel:1-888-931-2530)

LABEL

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

FIRST AID

IF POISONING IS SUSPECTED, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

Treat symptomatically. When large amounts have been ingested, gastric lavage may be required.

PRECAUTIONS

KEEP OUT OF THE REACH OF CHILDREN.

Harmful if swallowed. May cause eye irritation. Avoid contact with eyes or prolonged contact with skin. Avoid inhalation of vapours or spray mist.

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair.

Wear freshly laundered clothes daily. Wash the outside of the gloves before removing. Wash splashes from skin and eyes **IMMEDIATELY** with plenty of water. Avoid working in spray mist. Avoid all drift or contact with other vegetation. After spraying, wash hands and shower thoroughly with soap and water.

While using product, do not eat, drink or use tobacco (including smoking). Wash hands and exposed skin with soap and water thoroughly before eating, drinking, smoking, applying cosmetics or using the toilet. Store and

wash all protective clothing separately from household laundry. Do not wear contaminated shoes. Wash in detergent and hot water before reuse. Keep product away from food, drink and animal feed. Store product in original container, tightly closed, and in a safe place. Mixing and loading is not permitted within 30 m from aquatic habitats.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

ENVIRONMENTAL PRECAUTIONS

Toxic to non-target terrestrial plants and aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

The residues of atrazine are persistent and may carry over. It is recommended that this product not be used in areas treated with any products containing atrazine during the previous year.

STORAGE

Store this product away from food or feed. STORE IN A DRY PLACE.

DISPOSAL

Recyclable containers:

Disposal of container: DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

Refillable containers:

Disposal of container: For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

Disposal of unused, unwanted product: For information on disposal of unused, unwanted product,

contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for cleanup of spills.

SPRAY DRIFT MANAGEMENT FOR GROUND APPLICATIONS

GENERAL INFORMATION

Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

GROUND APPLICATION

Do not apply Canterra 480 SC at rates greater than 3.1 L/ha. Apply atrazine at a maximum of 1.5 kg a.i./ha per year either as a pre- or post-emergent application before corn reaches 30 cm in height. Treatments can be made once or twice per year, but the maximum amount of atrazine used in a year cannot exceed 1.5 kg a.i./ha. Atrazine may be applied alone or as a registered tank mix partner. Refer to the tank mix partners for further instructions.

Bulk sites must be diked around and under the bulk tank with an impermeable non-earthed dike. The dike must be large enough to hold and contain 110 percent of the capacity of the largest tank.

CAUTION: Do not graze treated immature crops or cut for fodder. In the case of corn, immature means before ear emergence. Do not harvest within 45 days of application on sweet corn and 60 days for field corn.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Sharda Cropchem Ltd. at [1-888-931-2530](tel:1-888-931-2530).

CANTERRA 480 SC

SUSPENSION

HERBICIDE

COMMERCIAL (AGRICULTURAL)

Liquid herbicide for selective weed control in corn, sorghum and switchgrass

ACTIVE INGREDIENT:

Atrazine and related triazines 480 g/L

Contains 1,2-benzisothiazolin-3-one at 0.017% as a preservative.

**READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF THE REACH OF CHILDREN**

REGISTRATION NO.: 35706
PEST CONTROL PRODUCTS ACT

Sharda Cropchem Limited
2nd Floor, Prime Business Park, Dashrathlal Joshi Road,
VileParle (west), Mumbai - 400056
India

Canadian Agent
Sharda Cropchem Limited
601 - 402 - 21st Street East
Saskatoon, SK, **S7K 0C3**
[1-888-931-2530](tel:1-888-931-2530)

BOOKLET

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

FIRST AID

IF POISONING IS SUSPECTED, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed, call a poison control centre or doctor **IMMEDIATELY** for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin **IMMEDIATELY** with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

Treat symptomatically. When large amounts have been ingested, gastric lavage may be required.

PRECAUTIONS

KEEP OUT OF THE REACH OF CHILDREN.

Harmful if swallowed. May cause eye irritation. Avoid contact with eyes or prolonged contact with skin. Avoid inhalation of vapours or spray mist.

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair.

Wear freshly laundered clothes daily. Wash the outside of the gloves before removing. Wash splashes from skin and eyes **IMMEDIATELY** with plenty of water. Avoid working in spray mist. Avoid all drift or contact with other vegetation. After spraying, wash hands and shower thoroughly with soap and water.

While using product, do not eat, drink or use tobacco (including smoking). Wash hands and exposed skin with soap and water thoroughly before eating, drinking, smoking, applying cosmetics or using the toilet. Store and

wash all protective clothing separately from household laundry. Do not wear contaminated shoes. Wash in detergent and hot water before reuse. Keep product away from food, drink and animal feed. Store product in original container, tightly closed, and in a safe place. Mixing and loading is not permitted within 30 m from aquatic habitats.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

ENVIRONMENTAL PRECAUTIONS

Toxic to non-target terrestrial plants and aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

The residues of atrazine are persistent and may carry over. It is recommended that this product not be used in areas treated with any products containing atrazine during the previous year.

STORAGE

Store this product away from food or feed. **STORE IN A DRY PLACE.**

DISPOSAL

Recyclable containers

Disposal of container: **DO NOT** reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

Refillable containers

Disposal of container: For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. **DO NOT** reuse this container for any other purpose.

Disposal of unused, unwanted product: For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for cleanup of spills.

SPRAY DRIFT MANAGEMENT FOR GROUND APPLICATIONS

GENERAL INFORMATION

Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings

For the protection of non-target habitats, over-spray or drift to any body of water or other environmentally sensitive habitat must be avoided. The interaction of equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions.

GROUND APPLICATION

Do not apply Canterra 480 SC at rates greater than 3.1 L/ha. Apply atrazine at a maximum of 1.5 kg a.i./ha per year either as a pre- or post-emergent application before corn reaches 30 cm in height. Treatments can be made once or twice per year, but the maximum amount of atrazine used in a year cannot exceed 1.5 kg a.i./ha. Atrazine may be applied alone or as a registered tank mix partner. Refer to the tank mix partners for further instructions.

Bulk sites must be diked around and under the bulk tank with an impermeable non-earthen dike. The dike must be large enough to hold and contain 110 percent of the capacity of the largest tank.

CAUTION: Do not graze treated immature crops or cut for fodder. In the case of corn, immature means before ear emergence. Do not harvest within 45 days of application on sweet corn and 60 days for field corn.

If this pest control product is to be used on a commodity that may be exported to other countries in the world and you require information on acceptable residue levels in these countries, please contact Sharda Cropchem Ltd at [1-888-931-2530](tel:1-888-931-2530).

DIRECTIONS FOR USE

GENERAL INFORMATION

Canterra 480 SC provides selective control of broadleaf weeds and wild oats in corn. This product may be applied before or after weeds emerge.

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

DO NOT overspray non-target terrestrial or aquatic habitats.

DO NOT apply this product through any type of irrigation system.

DO NOT apply by air.

This product may be tank mixed with (a fertilizer, a supplement, or with) registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Sharda Cropchem Limited at 1-888-931-2530 for information before applying any tank mix that is not specifically recommended on this label.

Field sprayer application: DO NOT apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. DO NOT apply with sprays finer than the American Society of Agricultural and Biological Engineers (ASABE) S572 (572.1 to 572.3) medium classification. Boom height must be 60 cm or less above the crop or ground.

SPRAY BUFFER ZONES:

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

| Method of Application | Spray Buffer Zone Required for the Protection of: | |
|-----------------------|---------------------------------------------------|---------------------|
| | Terrestrial Habitat (m) | Aquatic Habitat (m) |
| Field Sprayer | 10 | 10 |

When tank mixes are permitted, consult the labels of the tank mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASABE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product for conventional application equipment can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Drift Mitigation portion of the Canada.ca website.

WEEDS CONTROLLED:

Wormseed Mustard
Wild Mustard
Common Purslane
Ragweed
Smartweed
Lady's-Thumb

Wild Buckwheat
Lamb's-Quarters
Redroot Pigweed
Volunteer Clover
Wild Oats

NOTE: Some naturally occurring triazine tolerant biotypes of the above listed broadleaf weeds may not be controlled by atrazine alone. Refer to specific recommendations in the tank mix section of this label or consult your local agricultural authority.

TIMING OF APPLICATION:

Can be applied:

- Pre-plant incorporated (PPI)
- Pre-emergence
- Post-emergence

in either conventional tillage or minimum tillage programs.

If weeds escape a PPI or pre-emergence application, an additional post-emergence application may be necessary. Do not make more than one re-application per year and only in the spring.

LIQUID FERTILIZERS (CORN):

For pre-plant incorporated and pre-emergence application, nitrogen solutions or complete liquid fertilizers may replace the water as a carrier.

CAUTION: Do not apply after the corn has emerged as there is danger of liquid fertilizer causing crop injury. If there is any doubt about compatibility, check by mixing a small amount of Canterra 480 SC with the liquid fertilizer.

MIXING AND SPRAYING INSTRUCTIONS:

Protect sprayer operators from drift or mist. Additional information on spray drift management for GROUND APPLICATION is provided in the section "SPRAY DRIFT MANAGEMENT FOR GROUND APPLICATIONS". When low volumes of spray are applied, complete coverage and thorough application are essential for most effective results. Schedule applications in accordance with local conditions. Consult your local agricultural authorities for specific use information. Where possible, use a nurse tank to fill your spray tank.

Mixing Steps:

1. Spray equipment should be thoroughly flushed with clean water before mixing Canterra 480 SC.
2. Fill spray tank 1/2 full with carrier. Engage gentle agitation.
3. Pour in the required amount of wettable powder or wettable granule, if applicable.
4. Fill tank 3/4 full and add the Canterra 480 SC and other liquid products, if applicable.
5. Fill tank 9/10 full and add emulsifiable concentrate or liquid solutions, as required.
6. Finish filling tank. Maintain gentle agitation during mixing and spraying operations.
7. Do not let stand without agitation.
8. When using Canterra 480 SC, clean tank immediately after use with a strong detergent

and clean water.

- Spray pressure 200 - 300 kPa
- Spray volume - minimum 150 litres per hectare
- Use 16-mesh suction screen. 50 mesh screens elsewhere on sprayer.

APPLICATION INFORMATION - CORN

FOR CANTERRA 480 SC HERBICIDE ALONE, PPI AND PRE-EMERGENCE TREATMENTS

| SOIL TYPE | RATE OF APPLICATION/HA |
|-----------------|------------------------------------------|
| Light and Sandy | 2.1 - 3.0 ¹ L Canterra 480 SC |
| Loams and Clays | 3.1 L Canterra 480 SC |

¹ Use higher rate for heavy weed infestations.

Rainfall is required within 10 days or a shallow cultivation will be necessary.

Canterra 480 SC may be applied as an overall spray or as a band application. USE PROPORTIONATELY LESS PRODUCT FOR BAND APPLICATION.

PREPLANT INCORPORATED TANK MIXES (Silage, Field, Sweet Corn)

| WEEDS CONTROLLED | PRODUCT RATES |
|-----------------------------------------------------|----------------------------------------------------------------------------|
| Broadleaf Weeds and Wild Oats | 2.1 - 3.1 L/ha Canterra 480 SC |
| Yellow Nutsedge, Annual Grasses and Broadleaf Weeds | 2.1 - 3.1 L/ha Canterra 480 SC + 1.25 – 1.75 L/ha DUAL II Magnum Herbicide |

If tillage is intended after spraying, do not incorporate Canterra 480 SC deeper than 5.0 cm.

PRE-EMERGENT TANK MIXES (Silage, Field, Sweet Corn)

| WEEDS CONTROLLED | PRODUCT RATES | | |
|---------------------------------------------------------------------|-------------------------------------------|---|-----------------------------------------------------------------------------------|
| Broadleaf Weeds and Wild Oats | 2.1 - 3.1 L/ha Canterra 480 SC | | |
| Broadleaf Weeds, Wild Oats and Triazine Tolerant Weeds ³ | 2.1 - 3.1 L/ha Canterra 480 SC | + | 1.25 L/ha Banvel® II Herbicide ^{1,2} |
| Annual Grasses and Broadleaf Weeds | 2.1 - 3.1 L/ha Canterra 480 SC | + | 1.25 – 1.75 L/ha DUAL II Magnum Herbicide |
| Annual Grasses, Broadleaf Weeds and Triazine Tolerant Weeds | 2.1 - 3.1 L/ha Canterra 480 SC | + | 1.25 – 1.75 L/ha DUAL II Magnum Herbicide |
| | | + | 1.25 L/ha Banvel II Herbicide ^{1,2} |
| Annual Grasses, Broadleaf Weeds and Triazine Tolerant Weeds | 2.1 - 3.1 L/ha Canterra 480 SC | + | 1.25 – 1.75 L/ha |
| | | + | DUAL II Magnum Herbicide 0.3 L/ha CALLISTO® 480SC Herbicide ² |
| Annual Grasses, Broadleaf Weeds and Triazine and ALS Tolerant Weeds | 1.67 L/ha Canterra 480 SC | + | 0.165 L/ha Converge® Pro ^{1,2} |
| | OR 2.22 L/ha Canterra 480 SC | + | 0.22 L/ha Converge Pro ^{1,2} |

¹ DO NOT APPLY TO SWEET CORN.

² Eastern Canada only.

³ This tank mix will control triazine tolerant forms of the weeds listed on the Banvel II Herbicide label.

A firm, level seedbed free from large clumps and surface trash must be prepared to ensure uniform weed control.

Do not use Canterra 480 SC on loose and rough seedbeds.

POST-EMERGENT TANK MIXES (Silage, Field, Sweet Corn)

| WEEDS CONTROLLED | PRODUCT RATES | | |
|-----------------------------------------------------------------------|-----------------|-----------------|-------------------------------------------------|
| Broadleaf Weeds, Wild Oats and Triazine Tolerant Weeds ¹ | 2.1 - 3.1 L/ha | + | 0.6 L/ha |
| | Canterra 480 SC | | Banvel II Herbicide ² |
| | 2.1 - 3.1 L/ha | + | 1.0 L/ha |
| | Canterra 480 SC | | Pardner® Herbicide ³ |
| | 2.1 - 3.1 L/ha | + | 1.0 L/ha |
| | Canterra 480 SC | | Buctril® M ^{3,4} |
| Broadleaf Weeds and Triazine Tolerant Weeds ⁵ | 1.0 L/ha | + | 1.0 - 1.2 L/ha |
| | Canterra 480 SC | | Pardner Herbicide ³ |
| Annual Grasses and Broadleaf Weeds | 2.1 - 3.1 L/ha | + | 1.25 – 1.75 L/ha |
| | Canterra 480 SC | | DUAL II Magnum Herbicide |
| Annual Grasses, Broadleaf Weeds and Triazine Tolerant Weeds | 2.1 - 3.1 L/ha | + | 1.25 – 1.75 L/ha |
| | | Canterra 480 SC | DUAL II Magnum Herbicide |
| | | + | 0.3 L/ha |
| | | | CALLISTO 480SC Herbicide ^{6,7,8} |
| Eastern black nightshade, redroot pigweed, velvetleaf, common ragweed | 0.58 L/ha | + | 0.21 L/ha |
| | Canterra 480 SC | | CALLISTO 480SC Herbicide ^{6,8,9,10} |

¹ Tank mix with Banvel II Herbicide will control triazine tolerant forms of the weeds listed on their labels.

² DO NOT APPLY TO SWEET CORN.

³ WEAR CHEMICAL-RESISTANT NITRILE GLOVES WHEN MIXING/LOADING, SPRAYING AND DURING CLEAN UP AND REPAIR. SEE TANK MIX PARTNER LABEL.

⁴ CORN WILL BE DAMAGED IF TREATED WITH THIS TANK MIX AFTER THE 6-LEAF STAGE.

⁵ This treatment will provide control of:

- Redroot Pigweed (triazine susceptible) seedlings up to the 6-leaf stage.
- Velvetleaf seedlings up to the 6-leaf stage and less than 10 cm in height.
- Common Ragweed seedlings up to the 8-leaf stage.

⁶ Eastern Canada only.

⁷ Apply on spike to 2 leaf stage of corn.

⁸ Apply to field corn only.

⁹ Apply on 3 to 8 leaf stage of corn.

¹⁰ This tank mix with CALLISTO 480SC Herbicide must be applied with the non-ionic surfactant AGRAL® 90 at 0.2% v/v (2.0 L for each 1000 L of spray mixture).

- ALWAYS APPLY CANTERRA 480 SC BEFORE THE CORN REACHES 30 CM IN HEIGHT.

WARNING:

Injury may occur if land treated with Canterra 480 SC is planted to any crop other than field corn or sorghum in the same season. One year after application there is virtually no hazard to most rotational crops when annual weed label rates are applied. Sugar beets and tobacco are sensitive to Canterra 480 SC and should not be planted on land treated with Canterra 480 SC the previous year. When extended periods of dry weather occur during the year of treatment, there can be some injury to succeeding crops such as white beans, onions, peas, tomatoes and turnips. Injury is most likely to occur when the seedling crop is subjected to periods of stress e.g., during periods of abnormally hot, dry weather.

Under conditions of heavy rainfall sufficient to cause erosion, soil containing Canterra 480 SC may wash to lower areas of land and injury to existing or subsequent crops other than corn may result.

THOROUGH TILLAGE INCLUDING PLOUGHING SHOULD PRECEDE THE PLANTING OF CROPS OTHER THAN CORN. PLOUGH THE LAND FLAT AS OPPOSED TO SETTING THE FURROW ON EDGE. UNEVEN APPLICATION OF CANTERRA 480 SC OR APPLICATION IN EXCESS OF RECOMMENDED RATES WILL NOT INJURE CORN BUT MAY RESULT IN THE INJURY OF SUCCEEDING CROPS OTHER THAN CORN.

NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS:

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Sharda Cropchem Ltd. under the User Requested Minor Use Label Expansion program. For these uses, Sharda Cropchem Ltd. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

SWEET CORN TANK MIXTURE

| TIMING | WEEDS CONTROLLED | RATE/HA CANTERRA 480 SC | RATE PER HECTARE/TANK MIX PARTNER | REMARKS |
|--------------|----------------------------------------------------------------------|-------------------------|-----------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Pre-emergent | Annual grasses and broadleaf weeds including triazine tolerant weeds | 2.1 - 3.1 L | + 1.25 L DUAL II Magnum Herbicide | Make only one application per year. Apply as pre-emergent treatment. Apply in 150 L water/ha minimum. Do not harvest sweet corn within 50 days of treatment. Apply by ground equipment only. |

CONTROL OF BUR CUCUMBER (*Sicyos angulatus*)

| TIMING | RATE/HA CANTERRA 480 SC | RATE/HA PARDNER ¹ HERBICIDE | REMARKS |
|---------------|-------------------------|----------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Post-emergent | 3.1 L | 1.0 L | Make only one application per year. Apply in 150-300 L water/ha. Corn should be treated at the 4 to 8-leaf stage. Bur cucumber seedlings in the 1 to 4-leaf stage. Apply by ground equipment only. |

¹Consult the Pardner Herbicide label for specific instructions regarding soil-type variations, and rotational crop, grazing or other restrictions.

FIELD, SILAGE, AND SWEET CORN

| TIMING | WEEDS CONTROLLED | RATE/HA CANTERRA 480 SC | RATE/HA PARDNER ¹ HERBICIDE | REMARKS |
|----------------------------|------------------------------------------------------------------------------|-------------------------|----------------------------------------|----------------------------------------------------------------------------------------------------|
| Post-emergent ² | Volunteer adzuki bean (<i>Vigna angularis</i>), giant ragweed ³ | 3.1 L | 1.0 L | Make only one application per year. Apply in 200 L water/ha. Apply by ground equipment only. |

¹ Consult the Pardner Herbicide label for precautions.

² Apply at 4 to 8 leaf stage (corn), unifoliate to fourth trifoliate leaf stage (adzuki bean), between cotyledon and 10 leaf growth stage of giant ragweed.

³ Suppression only.

FIELD CORN (EASTERN CANADA ONLY)

| TIMING ¹ | WEEDS CONTROLLED | RATE/HA CANTERRA 480 SC | RATE PER HECTARE/TANK MIX PARTNER ² | REMARKS |
|----------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------|---------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------|
| Post-emergent ³ | Volunteer Adzuki Bean (<i>Vigna angularis</i>), large crabgrass (<i>Digitaria sanguinalis</i>) ⁵ , Giant Ragweed ⁴ , Common Cocklebur ⁴ | 0.58 L | 0.21 L CALLISTO 480SC Herbicide + 0.2% v/v AGRAL 90 in 100-200 L water/ha | Make only one application per year. Apply by ground equipment only. Pre-Harvest Interval is 100 days (grain/stover) and 90 days (forage). |

¹ Apply at 3-8 leaf stage of corn.

² Consult tank mix partner label for precautions.

³ Apply at the unifoliate to fourth trifoliate leaf stage of adzuki bean and when giant ragweed is at the cotyledon-10 leaf stage and when common cocklebur is up to the cotyledon – 6 leaf stage.

⁴ Suppression only.

⁵ Apply when large crabgrass is at the 1 – 6 leaf stage.

SORGHUM (EASTERN CANADA ONLY)

| TIMING¹ | WEEDS CONTROLLED | RATE/HA CANTERRA 480 SC | REMARKS |
|-------------------------------|-------------------------------|--------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Pre-emergent or Post-emergent | Broadleaf weeds and wild oats | 2.1 L | <p>Make only one application per year. Apply by ground equipment only. Apply in a minimum of 150 L of water/ha.</p> <p>Pre-Harvest Interval is 90 days for grain.</p> <p>Grazing Restrictions: Do not graze or feed forage from treated areas for 60 days following a pre-emergent or 45 days following a post-emergent application.</p> <p>Heavy rain immediately following application tends to cause excessive concentrations of herbicide in seed furrow, resulting in possible crop injury.</p> <p>Application to sorghum growing under stress caused by minor element deficiency or to sorghum growing on highly calcareous soil may result in crop injury.</p> <p>Injury may occur if both this herbicide and an at-planting systemic insecticide applied in-furrow are used.</p> <p>For all soil applied applications prior to crop emergence, do not apply to coarse textured soils, i.e., sand, loamy sand, sandy, or to medium – and fine-textured soils having less than 1% organic matter, or injury may occur.</p> |

SWITCHGRASS

| TIMING | WEEDS CONTROLLED | RATE/HA CANTERRA 480 SC | REMARKS |
|------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Pre-plant incorporated or pre-emergent to crop | Wormseed mustard, wild buckwheat, wild mustard, lamb's-quarters, common purslane, redroot pigweed, ragweed, volunteer clover, smartweed, wild oats, lady's-thumb | 2.1 – 3.1 L | <p>Make only one application per year.</p> <p>Apply in year of establishment only.</p> <p>Apply by ground equipment only.</p> <p>Apply in a minimum of 150 L of water/ha.</p> <p>Pre-harvest interval is 485 days.</p> <p>Do not graze treated areas. Do not use seeds for bird food.</p> |

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, Canterra 480 SC is a Group 5 herbicide. Any weed population may contain or develop plants naturally resistant to Canterra 480 SC and other Group 5 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

Where possible, rotate the use of Canterra 480 SC or other Group 5 herbicide within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.

Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.

Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.

Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.

Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.

For further information and to report suspected resistance, contact Sharda Cropchem Ltd. at [1-888-931-2530](tel:1-888-931-2530).

, AGRAL[®], CALLISTO[®], DUAL II Magnum[®], and Magnum[®] are trademarks of a Syngenta Group Company. Banvel[®], Buctril[®], Converge[®], and Pardner[®] are trademarks of their respective companies.