

2025-6181
2025-12-18

****Container Label****

GROUP	3	FUNGICIDE
-------	----------	-----------

JSAL Prothio 480 SC

Suspension

For Control or Suppression of Listed Diseases in Barley, Borage, *Brassica carinata*, Bushberries, Canola, Chickpeas, Crambe, Low Growing Berries - except strawberries, Corn, Cucurbits, Flax (Linseed), Lentils, Oats, Oriental Mustard, Peanut, Rapeseed, Safflower, Soybean, Sugar Beet, Sunflower, Wheat and Certain Other Small Grains.

COMMERCIAL (AGRICULTURAL)

SYMBOL – Triple Rinse



CAUTION

POISON

REGISTRATION NUMBER: **35768**
PEST CONTROL PRODUCTS ACT

ACTIVE INGREDIENT:
Prothioconazole.....480 g/L

Contains 1,2-benzisothiazolin-3-one at 0.04% as preservative.

READ THE LABEL BEFORE USING
KEEP OUT OF REACH OF CHILDREN

NET CONTENTS: 1 LITRE to BULK

Jiangsu Agrochem Laboratory Co., Ltd.
No. 98, Minjiang Rd, Hi-tech Development Zone,
Changzhou, Jiangsu, 213022, China
Tel: +86-519-8822 5306, Fax: +86-519-8510 8097

Canadian Representative
HarvestGuard Agronomy LTD.
88 Saskatchewan Street East, Moose Jaw Saskatchewan S6H 0V4,
Canada
Email: info@hgagro.ca Phone: 306-676-8888

For product information, call: 1-306-676-8888

GENERAL INFORMATION

JSAL Prothio 480 SC is a broad-spectrum systemic fungicide for the control or suppression of listed *Ascomycetes*, *Basidiomycetes* and *Deuteromycetes* diseases on barley, bushberries, *Brassica carinata*, borage, canola, chickpeas, crambe, low growing berries - except strawberries, corn, cucurbits, flax (linseed), lentils, oats, Oriental mustard, peanut, rapeseed, safflower, soybean, sugar beet, sunflower, wheat and certain other small grains.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, or smoking. If product comes in contact with clothing, remove all contaminated clothing, wash skin with soap and water and dress in clean clothing. Launder applicator clothing separate from other laundry. Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings. A restricted entry interval (REI) of 20 days is required for hand-harvesting sweet corn and for detasseling seed corn. An REI of 3 days is required for hand-line irrigation in cucurbits and bushberries. For all other activities do not re-enter treated fields until 24 hours after application. If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact Jiangsu Agrochem Laboratory Co., Ltd. at +1-306-676 8888.

PROTECTIVE CLOTHING AND EQUIPMENT:

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks, and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair. Gloves are not required during application within a closed cab or cockpit.

FIRST AID:

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label, or product name and Pest Control Product Registration number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS

Toxic to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

STORAGE

- Store this product away from food and feed.
- Keep away from direct sunlight.
- Keep away from fire or open flame or other sources of heat.
- Do not store at temperatures below freezing.
- If stored for 1 year or longer, shake well before using.
- Store the tightly closed container away from feeds, seeds, fertilizer, plants and foodstuffs.
- Do not use or store in or around the home.
- Keep in original container during storage.

DISPOSAL

Recyclable Containers:

Disposal of container: DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

Returnable Containers:

Disposal of container: DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Disposal of Unused, Unwanted Product:

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for cleanup of spills.

Disposal of Unused Spray Solution:

If any spray solution remains in the tank after spraying is finished, it should be sprayed on the perimeter of the area just sprayed, NOT ON THE CROP, away from water supplies, ditches, and irrigation canals. Spray buffer zones indicated in the DIRECTIONS FOR USE must also be respected.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

For more information contact:

Jiangsu Agrochem Laboratory Co., Ltd.

No. 98, Minjiang Rd, Hi-tech Development Zone,
Changzhou, Jiangsu, 213022, China

All other products mentioned are trademarks of their respective companies.

JSAL Prothio 480 SC

Suspension

For Control or Suppression of Listed Diseases in Barley, Borage, *Brassica carinata*, Bushberries, Canola, Chickpeas, Crambe, Low Growing Berries - except strawberries, Corn, Cucurbits, Flax (Linseed), Lentils, Oats, Oriental Mustard, Peanut, Rapeseed, Safflower, Soybean, Sugar Beet, Sunflower, Wheat and Certain Other Small Grains.

COMMERCIAL (AGRICULTURAL)

SYMBOL - Triple Rinse



CAUTION

POISON

REGISTRATION NUMBER: **35768**
PEST CONTROL PRODUCTS ACT

ACTIVE INGREDIENT:
Prothioconazole.....480 g/L

Contains 1,2-benzisothiazolin-3-one at 0.04% as preservative.

**READ THE LABEL BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

NET CONTENTS: 1 LITRE to BULK

Jiangsu Agrochem Laboratory Co., Ltd.
No. 98, Minjiang Rd, Hi-tech Development Zone,
Changzhou, Jiangsu, 213022, China
Tel: +86-519-8822 5306, Fax: +86-519-8510 8097

Canadian Representative
HarvestGuard Agronomy LTD.
88 Saskatchewan Street East, Moose Jaw Saskatchewan S6H 0V4,
Canada
Email: info@hgagro.ca Phone: 306-676-8888
For product information, call: 1-306-676-8888

JSAL Prothio 480 SC

Table of Contents:

Section Number:

GENERAL INFORMATION

The Product	1
-------------	---

SAFETY AND HANDLING

Precautions, Protective Clothing & Equipment, and Re-entry Restriction	2
First Aid and Toxicological Information	3
Environmental Precautions	4
Storage	5
Disposal	6

DIRECTIONS FOR USE

Crop, Disease, Rate, Timing	7
Resistance Management	8
Application Precautions	9
Mixing Instructions	10
Pre-harvest Intervals	11
Notices	12

For more information, contact:

Jiangsu Agrochem Laboratory Co., Ltd.

No. 98, Minjiang Rd, Hi-tech Development Zone,
Changzhou, Jiangsu, 213022, China

GENERAL INFORMATION

Section 1: The Product

JSAL Prothio 480 SC is a broad-spectrum systemic fungicide for the control or suppression of listed *Ascomycetes*, *Basidiomycetes* and *Deuteromycetes* diseases on barley, bushberries, *Brassica carinata*, borage, canola, chickpeas, crambe, low growing berries - except strawberries, corn, cucurbits, flax (linseed), lentils, oats, Oriental mustard, peanut, rapeseed, safflower, soybean, sugar beet, sunflower, wheat and certain other small grains.

SAFETY AND HANDLING

Section 2: Precautions, Protective Clothing & Equipment, and Re-entry Restriction.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, or smoking. If product comes in contact with clothing, remove all contaminated clothing, wash skin with soap and water and dress in clean clothing. Launder applicator clothing separate from other laundry. Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings. A restricted entry interval (REI) of 20 days is required for hand-harvesting sweet corn and for detasseling seed corn. An REI of 3 days is required for hand-line irrigation in cucurbits and bushberries. For all other activities do not re-enter treated fields until 24 hours after application. If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact Jiangsu Agrochem Laboratory Co., Ltd. at +1-306-676-8888.

PROTECTIVE CLOTHING AND EQUIPMENT:

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks, and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair. Gloves are not required during application within a closed cab or cockpit.

Section 3: First Aid and Toxicological Information

FIRST AID:

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label, or product name and Pest Control Product Registration number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Treat symptomatically.

Section 4: ENVIRONMENTAL PRECAUTIONS

Toxic to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

Section 5: STORAGE

- Store this product away from food and feed.
- Keep away from direct sunlight.
- Keep away from fire or open flame or other sources of heat.
- Do not store at temperatures below freezing.
- If stored for 1 year or longer, shake well before using.
- Store the tightly closed container away from feeds, seeds, fertilizer, plants and foodstuffs.
- Do not use or store in or around the home.
- Keep in original container during storage.

Section 6: DISPOSAL

Recyclable Containers:

Disposal of container: DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

Returnable Containers:

Disposal of container: DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Disposal of Unused, Unwanted Product:

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for cleanup of spills.

Disposal of Unused Spray Solution:

If any spray solution remains in the tank after spraying is finished, it should be sprayed on the perimeter of the area just sprayed, NOT ON THE CROP, away from water supplies, ditches, and irrigation canals. Spray buffer zones indicated in the DIRECTIONS FOR USE must also be respected.

DIRECTIONS FOR USE

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Field sprayer application: DO NOT apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. DO NOT apply with sprays finer smaller than the American Society of Agricultural and Biological Engineers (ASABE) S572 (572.1 to 572.3) medium classification. Boom height must be 60 cm or less above the crop or ground.

Airblast application: DO NOT apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. DO NOT direct spray above plants to be treated. Turn off outward pointing nozzles at row ends and outer rows. DO NOT apply when wind speed is greater than 16 km/h at the application site as measured outside of the treatment area on the upwind side.

Conventionally piloted aircraft application: DO NOT apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with sprays finer smaller than the American Society of Agricultural and Biological Engineers (ASABE) S572 (572.1 to 572.3) medium classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length MUST NOT exceed 65% of the wing- or rotorspan.

Apply only by aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate specified for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices or a Global Positioning System (GPS).

To ensure proper coverage and distribution and to minimize drift, check to see that swath width and droplet size are adequate, and that wind velocities are low.

Use mechanical flaggers only.

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the registrant at (306)6768888 or obtain technical advice from the distributor or your provincial or territorial agricultural representative. Application of this specific product must meet and/or conform to the following:

Volume: Apply the specified rate in a minimum spray volume of 50 litres per hectare for conventionally piloted aircraft.

Operator Precautions

- DO NOT allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.
- It is desirable that the pilot has communication capabilities at each treatment site at the time of application.
- The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.
- All personnel on the job site must wash hands and face thoroughly before eating and drinking.
- Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Tank Mixes:

This product may be tank mixed with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Jiangsu Agrochem Laboratory Co., Ltd. at 1-306-676-8888 for information before applying any tank mix that is not specifically recommended on this label.

Spray Buffer Zones:

A spray buffer zone is NOT required for :

- uses with hand-held application equipment permitted on this label

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of Application	Crop		Spray Buffer Zones (metres) Required for the Protection of:				
			Freshwater Habitat of Depths:		Estuarine/Marine Habitat of Depths:		Terrestrial Habitat:
			Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer	Wheat, barley, millet, rye, triticale, teosinte, buckwheat, canola, Oriental mustard, rapeseed, <i>Brassica carinata</i> , chickpeas, lentils, cucurbits, bushberries, low growing berry (except strawberry), sugar beets, peanuts		1	1	1	1	1
	Oats, soybean, corn, safflower, sunflower, flax, crambe, borage		1	1	1	1	0
Airblast	Bushberries	Early growth stage	15	5	5	2	1
		Late growth stage	5	3	3	1	1
Aerial	Soybean	Fixed wing	1	1	1	1	0

		Rotary wing	1	1	1	1	0
	Flax (linseed), crambe, borage	Fixed wing	5	1	2	1	0
		Rotary wing	4	1	1	1	0
	Corn, safflower, sunflower	Fixed wing	10	1	5	1	0
		Rotary wing	5	1	1	1	0
	Oats	Fixed wing	10	1	1	1	0
		Rotary wing	10	1	1	1	0
	Canola, Oriental mustard, rapeseed, <i>Brassica carinata</i>	Fixed wing	10	2	2	1	10
		Rotary wing	10	1	1	1	10
	Wheat, barley	Fixed wing	10	3	5	1	10
		Rotary wing	10	1	1	1	10
	Lentils	Fixed wing	10	5	5	1	15
		Rotary wing	10	2	1	1	10
	Chickpeas, sugarbeets	Fixed wing	15	10	5	1	15
		Rotary wing	15	5	1	1	15

When tank mixes are permitted, consult the labels of the tank mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASABE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product for conventional application equipment can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Drift Mitigation portion of the Canada.ca website.

When applied as directed, JSAL Prothio 480 SC will control the listed diseases, unless otherwise indicated as suppression.

Section 7: Crop, Disease, Rate, Timing

CEREALS (Barley, Wheat and Oats)			
CROP	DISEASE CONTROLLED	RATE	TIMING

Barley	Fusarium head blight or scab (<i>Fusarium</i> spp.), (Suppression only)	Apply once, at 315-420 mL/ha (150-200 g a.i./ha) Adjuvant: Apply with a non-ionic surfactant, e.g. AgSurf or Agral 90, at 0.125% v/v	Fusarium head blight outbreaks in barley occur when the weather is warm and wet at ear emergence. The application of JSAL Prothio 480 SC for protection against Fusarium head blight (scab) should be considered when these environmental conditions are forecasted for this stage of barley development. Timing of application is critical: For optimum suppression of Fusarium head blight, apply JSAL Prothio 480 SC as a preventative spray within the time period when 70 to 100% of the barley main stem heads are fully emerged, to 3 days after full head emergence. Application of the 420 mL/ha rate is suggested in situations where disease pressure is expected to be high. Such situations may occur when prolonged periods of warm wet weather are forecast during barley ear emergence, when barley is grown in a crop rotation that has contained corn or when susceptible cultivars are grown. Use of the 420 mL/ha rate will typically provide highest levels of mycotoxin reduction. Spray Coverage is essential for optimum efficacy: Spray equipment must be setup to provide good coverage to barley heads. To achieve thorough barley head coverage using ground application equipment, it is recommended to use forward and backward mounted nozzles or nozzles that have a two-directional spray. Nozzles should be operated within the spray pressure recommendations suggested by the manufacturer.
	Net blotch (<i>Pyrenophora teres</i>) Scald (<i>Rhynchosporium secalis</i>) Spot blotch (<i>Cochliobolus sativus</i>)	Apply once, at 210-315 mL/ha (100-150 g a.i./ha) Adjuvant: Apply with a non-ionic surfactant, e.g. AgSurf or Agral 90 at 0.125% v/v	Apply JSAL Prothio 480 SC as a preventive foliar spray when the earliest disease symptoms appear on the leaves and stems. Barley fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.
Note: A maximum of 2 applications (735 mL/ha) of JSAL Prothio 480 SC, with a minimum 7 day application interval, may be applied per barley crop, per year. Do not apply within 30 days of harvest. Applications may be made by ground or aerial spray equipment.			

CROP	DISEASE CONTROLLED	RATE	TIMING
Wheat (spring, durum and winter)	Fusarium head blight or scab (<i>Fusarium</i> spp.) (Suppression only)	Apply once, at 315-420 mL/ha (150-200 g a.i./ha)	Fusarium head blight outbreaks in wheat occur when the weather is warm and wet at head emergence and flowering. The application of JSAL Prothio 480 SC for protection against Fusarium head blight (scab) should be considered when these environmental conditions are forecasted for this stage of wheat development.
	Glume blotch (<i>Stagonosora nodorum</i>)	Adjuvant: Apply with a non-ionic surfactant, AgSurf or Agral 90 at 0.125% v/v	Timing of application is critical: For optimum suppression of Fusarium head blight or control of Glume blotch, apply JSAL Prothio 480 SC within the time period from when at least 75% of the wheat heads on the main stem are fully emerged to when 50% of the heads on the main stem are in flower. Application of the 420 mL/ha rate is suggested in situations where disease pressure is expected to be high. Such situations may occur when prolonged periods of warm, wet weather are forecast during anthesis, when wheat is grown in a crop rotation that has contained corn or when more susceptible cultivars or wheat types (i.e. durum) are grown. Use of the 420 mL/ha rate will typically provide highest levels of mycotoxin reduction. Spray Coverage is essential for optimum efficacy: Spray equipment must be setup to provide good coverage to wheat heads. To achieve thorough wheat head coverage using ground application equipment, it is recommended to use forward and backward mounted nozzles or nozzles that have a two-directional spray. Nozzles should be operated within the spray pressure recommendations suggested by the manufacturer.
	Speckled leaf blotch (<i>Septoria tritici</i>)	315 mL/ha (150 g a.i./ha)	Apply JSAL Prothio 480 SC as a preventive foliar spray when the earliest disease symptoms appear on the leaves and stems. Wheat fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.
	Tan spot (<i>Pyrenophora tritici-repentis</i>)	Adjuvant: Apply with a non-ionic surfactant, AgSurf or Agral 90 at 0.125% v/v	
	Leaf rust (<i>Puccinia recondita</i>)		
Note: A maximum of 2 applications (735 mL/ha) of JSAL Prothio 480 SC, with a minimum 7 day application interval, may be applied per wheat crop per year. Do not apply within 30 days of harvest. Applications may be made by ground or aerial spray equipment.			

CROP	DISEASE CONTROLLED	RATE	TIMING
Oats	Crown rust (<i>Puccinia coronate</i>)	315 mL/ha (150 g a.i./ha) Adjuvant: Apply with a non-ionic surfactant, AgSurf or Agral 90 at 0.125% v/v	Apply JSAL Prothio 480 SC as a preventive foliar spray when the earliest disease symptoms appear on the leaves and stems. Fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.

Note: A maximum of 2 applications (630 mL/ha) of JSAL Prothio 480 SC, with a minimum 7 day application interval, may be applied per oat crop per year. Do not apply within 30 days of harvest. Applications may be made by ground or aerial spray equipment.

CROP	DISEASE CONTROLLED	RATE	TIMING
Pearl millet, Proso millet, Rye, Triticale, Teosinte, Buckwheat	Fusarium head blight or Scab (<i>Fusarium</i> spp.) – except buckwheat (Suppression only) Rust (<i>Puccinia</i> spp.)	315-420 mL/ha (150-200 g a.i./ha) Adjuvant: Apply with a non-ionic surfactant, AgSurf or Agral 90 at 0.125% v/v	Apply JSAL Prothio 480 SC as a preventive foliar spray when the earliest disease symptoms appear on the leaves and stems. Fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development.

Note: A maximum of 2 applications (735 mL/ha) of JSAL Prothio 480 SC, with a minimum 7 day application interval, may be applied per year. Do not apply within 30 days of harvest. Applications may be made by ground equipment only.

OILSEED CROPS (Canola, Rapeseed, Oriental Mustard, *Brassica carinata*)

CROP	DISEASE CONTROLLED	RATE	TIMING
Canola, Rapeseed, Oriental mustard, <i>Brassica carinata</i>	Sclerotinia stem rot (<i>Sclerotinia sclerotiorum</i>)	315-368 mL/ha (150-175 g a.i./ha)	Apply JSAL Prothio 480 SC when the crop is in the 20 - 50% bloom stage. Best protection will be achieved when the fungicide is applied prior to petals beginning to fall, and will allow for the maximum number of petals to be protected. The 315 mL/ha rate is the recommended rate for most oilseed crops, however, the higher rate is recommended for fields with a history of heavy disease pressure or for dense crop stands. Good spray coverage of the plants is essential. A second application at a rate of 315 mL/ha may be applied 7-10 days later, up to full bloom, if disease persists or weather conditions are favourable for disease development. When conditions favouring disease are severe, use the shorter interval.

Note: The lowest labelled rate of a non-ionic surfactant, AgSurf or Agral 90, may be tank mixed with JSAL Prothio 480 SC. The application may be made by ground or aerial spray equipment. A maximum of two applications (683 mL/ha of JSAL Prothio 480 SC) may be applied per crop year. Do not apply within 36 days of harvest.

PULSE CROPS (Chickpeas, Lentils)

CROPS	DISEASE CONTROLLED	RATE	TIMING
-------	--------------------	------	--------

Chickpeas	Ascochyta blight (<i>Ascochyta rabiei</i>)	315-420 mL/ha (150-200 g a.i./ha)	Apply JSAL Prothio 480 SC at the first sign of disease. After the initial application, repeat applications may be made on 10-14 day intervals. Apply the higher rate when conditions favour disease development, or when growing less disease resistant varieties.
	Grey mould (<i>Botrytis cinerea</i>) (Suppression only)	420 mL/ha (200 g a.i./ha)	A maximum of 1260 mL/ha of JSAL Prothio 480 SC may be applied per crop year for chickpeas. Maximum of three applications per year.
Lentils	Ascochyta blight (<i>Ascochyta</i> spp.)	315-420 mL/ha (150-200 g a.i./ha)	Apply JSAL Prothio 480 SC at the beginning of flowering or at first sign of disease. After the initial application, one additional application may be made 10-14 days afterwards if conditions remain favourable for continued or increased disease development. Apply the higher rate when conditions favour disease development, or when growing less disease resistant varieties.
	White mould (<i>Sclerotinia sclerotiorum</i>)		
	Anthracnose* (<i>Colletotrichum lentis</i>) (Suppression only)	320-420 mL/ha (154-200 g a.i./ha)	
	Grey mould (<i>Botrytis cinerea</i>) (Suppression only)	420 mL/ha (200 g a.i./ha)	A maximum of 840 mL/ha of JSAL Prothio 480 SC may be applied per crop year for lentils. Maximum of two applications per year.
Note: The lowest labelled rate of a non-ionic surfactant, AgSurfor Agral 90, may be tank mixed with JSAL Prothio 480 SC. Applications may be made by ground or aerial spray equipment. Do not apply within 7 days of cutting or swathing of the crop. *Including biotypes resistant to Group 11 (strobilurin) fungicides.			

Soybean			
CROP	DISEASE CONTROLLED	RATE	TIMING
Soybean	Asian soybean rust (<i>Phakopsora pachyrhizi</i>)	210 mL/ha (100 g a.i./ha)	Apply JSAL Prothio 480 SC when first symptoms of disease can be found or that the risk of infection is imminent.
	Frogeye leaf spot (<i>Cercospora soja</i>)		Do not apply more than one application per season.
Note: The lowest labelled rate of a non-ionic surfactant, AgSurfor Agral 90, may be tank mixed with JSAL Prothio 480 SC. The application may be made by ground or aerial spray equipment. Do not apply within 20 days of harvest.			

Corn (field, sweet and popcorn)			
CROP	DISEASE CONTROLLED	RATE	TIMING

Corn (field, sweet and popcorn, including seed production)	Rusts (<i>Puccinia sorghi</i> , <i>Puccinia polysora</i>) Eyespot (<i>Kabatiellazeae</i> or <i>Aureobasidium zeae</i>) Northern blight (<i>Setosphaeria turcica</i>)	315 mL/ha (150 g a.i./ha)	Apply JSAL Prothio 480 SC as a preventive foliar spray when the earliest disease symptoms appear on the leaves and stems. Fields should be observed closely for early disease symptoms, particularly under prolonged conditions favorable for disease development. Under high disease pressure, it is recommended to use a non-ionic surfactant with JSAL Prothio 480 SC (do not apply a non-ionic surfactant prior to tassel emergence as crop injury may occur). Refer to the section “Note” of this table for directions.
	Provides control of all diseases listed above, plus the following: Grey leaf spot (<i>Cercospora zeae-maydis</i>) Fusarium and Gibberella ear rots (<i>Fusarium</i> spp. and <i>Gibberella</i> spp.) (Suppression only) Stalk rot pathogens <i>Fusarium</i> spp., <i>Gibberella</i> spp. and <i>Colletotrichum</i> spp. which may cause stalk lodging. (Suppression only)	420 mL/ha (200 g a.i./ha)	Fusarium Ear Rot outbreaks in corn occur when the weather is warm and wet during silking. It is also driven by hybrid selection and field history of severe fusarium outbreaks. The application of JSAL Prothio 480 SC for protection against Fusarium Ear Rot should be considered when these conditions are forecasted for a specific corn field. Timing of application is critical: For optimum suppression of Fusarium Ear Rot apply JSAL Prothio 480 SC from the development stage of corn between the tip of stigmata visible (silking, BBCH 63) to the stigmata drying (silk browning, BBCH 67). JSAL Prothio 480 SC will reduce both disease symptoms and levels of mycotoxin in the grain. Spray Coverage is essential for optimum efficacy: Spray equipment must be setup to provide good coverage to the cobs. To achieve thorough cob coverage using ground application equipment, it is recommended to use drop-nozzles. Nozzles should be operated within the spray pressure recommendations suggested by the manufacturer.
Note: The lowest labelled rate of a non-ionic surfactant, AgSurf or Agral 90, may be tank mixed with JSAL Prothio 480 SC. Do not apply a non-ionic surfactant prior to tassel emergence. The application may be made by ground or aerial spray equipment. Do not apply within 14 days of harvest. Do not apply more than one application per season.			

Peanut		
DISEASE CONTROLLED	RATE	TIMING
<u>Soil-Borne:</u> <i>Rhizoctonia</i> pod rot (<i>Rhizoctonia solani</i>) (Suppression only)	420 mL/ha (200 g a.i./ha)	For Soil-Borne Diseases Applications of fungicides with a different mode of action should be made prior to and following applications of JSAL Prothio 480 SC to discourage development of resistant strains of fungi. In fields with a history of pod rot or in situations where conditions favour disease development apply JSAL Prothio 480 SC at about mid-season and continue JSAL Prothio 480 SC applications at 14-day intervals. JSAL Prothio 480 SC must be carried by rainfall or irrigation into the root and pod zone for control of pod rot caused by <i>Rhizoctonia solani</i>. Drought conditions will decrease the effectiveness of JSAL Prothio 480 SC against pod rot.

<u>Foliar:</u> Early leaf spot (<i>Cercospora arachidicola</i>) Leaf rust (<i>Puccinia arachidis</i>) (Suppression only)	365-420 mL/ha (175-200 g a.i./ha)	Foliar Disease Spray Program Apply the specified rate in a preventive spray schedule. Apply up to 4 sprays using a 14-day interval Use the higher use rate when conditions are favourable for severe disease pressure and/or when growing less disease resistant varieties.
Note: Apply up to four (4) applications of JSAL Prothio 480 SC per year. When planting varieties with good to excellent levels of resistance to foliar diseases, the application interval may be extended up to 21 days in the absence of soil borne diseases. A maximum of 1.7 L per hectare of JSAL Prothio 480 SC may be applied per year. JSAL Prothio 480 SC may be applied up to 14 days before harvest. Do not feed hay or threshings or allow livestock to graze in treated areas. Applications may be made by ground spray equipment only.		

Cucurbit Vegetables			
CROP	DISEASE CONTROLLED	RATE	TIMING
Cucurbit vegetables: Chinese waxgourd (Chinese preserving melon); citron melon; cucumber; gherkin; gourd, edible (includes hyotan, cucuzza, hechima, Chinese okra); <i>Momordica</i> spp (includes balsam apple, balsam pear, bittermelon, Chinese cucumber); muskmelon (include cantaloupe); pumpkin; squash, summer; squash, winter (includes butternut squash, calabaza, hubbard squash, acorn squash, spaghetti squash); watermelon	Fusarium wilt and Fusarium blight (<i>Fusarium</i> spp.) Gummy stem blight (<i>Didymella</i> spp.)	420 mL/ha (200 g a.i./ha) (soil or foliar)	Apply JSAL Prothio 480 SC at planting via drip /drench equipment. Follow with foliar treatments during the season.
	Powdery mildew (<i>Sphaerotheca fuliginea</i>)	210-420 mL/ha (100-200 g a.i./ha)	For Powdery Mildew Control apply JSAL Prothio 480 SC as a preventive foliar spray when the earliest sign of disease is first noticed., repeat at 5 - 10 day intervals with JSAL Prothio 480 SC or another approved fungicide from a different chemical group.
General Comments: Apply up to one (1) soil application and two (2) foliar applications of JSAL Prothio 480 SC at the high rate, or one soil application and four (4) foliar applications per year at the low rate. Repeat foliar applications using a 5- to 10-day spray interval if conditions remain favorable for continued or increasing disease development. Foliar applications may be applied by ground application equipment only. Apply with a non-ionic surfactant, AgSurf or Agral 90 at 0.125% v/v. A maximum of 1.26 L/ha of JSAL Prothio 480 SC may be applied per crop year. Do not apply within 7 days of harvest.			

Bushberry			
CROP	DISEASE CONTROLLED	RATE	TIMING
Bushberry subgroup: Aronia berry; blueberry, highbush; blueberry, lowbush; buffalo currant; cranberry, highbush; currant, black; currant, red; elderberry; European barberry; gooseberry; honeysuckle, edible; huckleberry; jostaberry; Juneberry (Saskatoon berry); lingonberry; salal; sea buckthorn; cultivars, varieties, and/or hybrids of these.	Septoria leaf spot (<i>Septoria</i> spp.) (Suppression only)	315 mL/ha (150 g a.i./ha).	Apply JSAL Prothio 480 SC at the first sign of disease. After the initial application, one additional application may be made 10- 14 days afterwards if conditions remain favourable for continued or increased disease development.
	Leaf rust (<i>Thekopsora minima</i>) (Suppression only) Valdensinia leaf spot (<i>Valdensinia heterodoxa</i>) - blueberry only (Suppression only)	400 mL/ha (190 g a.i./ha)	
	Monilinia blight (<i>Monilinia vaccinii-corymbosi</i>) - blueberry only	315-420 mL/ha (150-200 g a.i./ha)	Begin applications at early bloom for fruit rot. Make a second application of JSAL Prothio 480 SC or another approved fungicide 5- 10 days later.
General Comments: Apply up to two (2) applications of JSAL Prothio 480 SC per year. Applications may be made by ground application equipment only. Apply with a non-ionic surfactant, AgSurfor Agral 90 at 0.125% v/v.			

A maximum of 840 mL/ha of JSAL Prothio 480 SC may be applied per crop year. Do not apply within 7 days of harvest.

Low-Growing Berry, except strawberry

CROP	DISEASE CONTROLLED	RATE	TIMING
Low growing berry subgroup, except strawberry; Bearberry; bilberry; cloudberry; cranberry; partridgeberry; cultivars, varieties, and/or hybrids of these	Fruit rots caused by: <i>Coleophoma empetri</i> , <i>Glomerella cingulata</i> , <i>Phyllosticta vaccinii</i> , <i>Physalospora vaccinii</i> , <i>Allantophomopsis lycopodina</i> , <i>Allantophomopsis cytispora</i> , <i>Fusicoccum putrefaciens</i> , <i>Penicillium</i> spp., <i>Phomopsis vaccinii</i> , <i>Colletotrichum acutatum</i> , <i>Colletotrichum coccodes</i> Septoria leaf spot (<i>Septoria</i> spp.) (Suppression only)	365 mL/ha (0.175 kg a.i./ha) Adjuvant: Apply with a non-ionic surfactant, AgSurf or Agral 90 at 0.125% v/v	Begin applications at early bloom for fruit rot. Make a second application of JSAL Prothio 480 SC or another approved fungicide 5-10 days later.

General Comments: Apply up to two (2) applications of JSAL Prothio 480 SC per year. Repeat applications using a 5- to 10-day spray interval if conditions remain favorable for continued or increasing disease development. Applications may be made by ground spray equipment only.

A maximum of 730 mL/ha of JSAL Prothio 480 SC may be applied per crop year. Do not apply within 45 days of harvest.

Sunflower, Safflower

DISEASE CONTROLLED	RATE	TIMING
Sunflower rust (<i>Puccinia helianthi</i>) Safflower rust (<i>Puccinia carthami</i>)	420 mL/ha (200 g a.i./ha)	Apply JSAL Prothio 480 SC when average rust severity reaches 1% on the upper 4, fully expanded leaves prior to or during bloom.

Note: The lowest labelled rate of a non-ionic surfactant, AgSurf or Agral 90, may be tank mixed with JSAL Prothio 480 SC. The application may be made by ground or aerial spray equipment. A maximum of one application (420 mL/ha of JSAL Prothio 480 SC) may be applied to sunflower or safflower per crop year. Do not apply within 45 days of harvest. If applying JSAL Prothio 480 SC to greater than 310 ha per day of sunflower and/or safflower using open cab groundboom equipment, applicators must also wear a NIOSH/MSHA-approved respirator.

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Jiangsu Agrochem Laboratory Co., Ltd. under the User Requested Minor Use Label Expansion program. For these uses, Jiangsu Agrochem Laboratory Co., Ltd. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

CROP	DISEASE SUPPRESSED	RATE	TIMING
------	--------------------	------	--------

Sunflower, Safflower	Sclerotinia head rot (<i>Sclerotinia sclerotiorum</i>)	420 mL/ha (200 g a.i./ha)	Apply JSAL Prothio 480 SC when the crop is in the 10 - 50% disk flower bloom stage.
-------------------------	---	------------------------------	---

Note: The lowest labelled rate of a non-ionic surfactant, AgSurf or Agral 90, may be tank mixed with JSAL Prothio 480 SC. The application may be made by ground or aerial spray equipment. A maximum of one application (420 mL/ha of JSAL Prothio 480 SC) may be applied to sunflower or safflower per crop year. Do not *apply* within 45 days of harvest. If applying JSAL Prothio 480 SC to greater than 310 ha per day of sunflower and/or safflower using open cab groundboom equipment, applicators must also wear a NIOSH/MSHA-approved respirator.

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Jiangsu Agrochem Laboratory Co., Ltd. under the User Requested Minor Use Label Expansion program. For these uses, Jiangsu Agrochem Laboratory Co., Ltd. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

CROP	DISEASE CONTROLLED	RATE	TIMING
Flax (linseed), Crambe, Borage	Sclerotinia stem rot (<i>Sclerotinia sclerotiorum</i>)	315-368 mL/ha (150- 175 g a.i./ha)	Apply JSAL Prothio 480 SC when the crop is in the 20 - 50% bloom stage. Best protection will be achieved when the fungicide is applied prior to petals beginning to fall, and will allow for the maximum number of petals to be protected. The higher rate is recommended for fields with a history of heavy disease pressure or for dense crop stands. Good spray coverage of the plants is essential.

Note: The lowest labelled rate of a non-ionic surfactant, AgSurf or Agral 90, may be tank mixed with JSAL Prothio 480 SC. The application may be made by ground or aerial spray equipment. A maximum of one application (368 mL/ha of JSAL Prothio 480 SC) may be applied per crop year. Do not apply within 36 days of harvest.

CROP	DISEASE CONTROLLED	RATE	TIMING
Sugar beets	Foliar disease: Cercospora leaf spot (<i>Cercospora beticola</i>)	315-415 mL/ha (150-200 g a.i./ha)	Apply JSAL Prothio 480 SC at the first sign of disease. Use the higher rate and shorter intervals when conditions are favourable for severe disease pressure and/or when growing less disease-resistant varieties. Apply the specified rate of JSAL Prothio 480 SC in 100 - 200 L of water per hectare when using ground application equipment. Repeat applications using a 14 to 21-day spray interval depending on disease pressure. Use a 14-day spray interval under normal to heavy disease pressure and a 21-day spray interval under light disease pressure.
	Soil-borne disease: Rhizoctonia crown rot (<i>Rhizoctonia solani</i>)	415 mL/ha (200 g a.i./ha)	Apply JSAL Prothio 480 SC at the 4-leaf to row closure growth stage. Apply the specified rate of JSAL Prothio 480 SC in 50 - 100 L of water per hectare. Repeat applications using a 21-day spray interval.

Note: A maximum of 3 applications (1245 mL/ha of JSAL Prothio 480 SC) may be applied per crop year. Do not apply within 7 days of harvest. The lowest labelled rate of a non-ionic surfactant (AgSurf or Agral 90 at 0.125% v/v) may be tank mixed with JSAL Prothio 480 SC. The application may be made by ground or aerial application equipment. JSAL Prothio 480 SC is a Group 3 fungicide. To limit the potential for development of disease resistance, alternate every application of JSAL Prothio 480 SC with a non-Group 3 fungicide.

Rotation restrictions:

Treated areas may be replanted with any crop specified on this label as soon as practical after the last application. For crops not listed on this label, do not plant back within 30 days of last application.

Section 8: Resistance Management**Resistance Management Recommendations:**

For resistance management, **JSAL Prothio 480 SC** contains a Group 3 fungicide. Any fungal population may contain individuals naturally resistant to JSAL Prothio 480 SC and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist.

Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

- Alternate with fungicides having a different mode of action other than Group 3 every 7 to 10 days after each application of **JSAL Prothio 480 SC**.
- Do not apply more than the indicated maximum number of applications per year specified for each crop in the DIRECTIONS FOR USE, Section 7 table.
- Use tank mixtures with fungicides from a different group that is effective on the target pathogen when such use is permitted.
- Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications.
- Monitor treated fungal populations for resistance development. Notify Jiangsu Agrochem Laboratory Co., Ltd. if reduced sensitivity of the pathogen to JSAL Prothio 480 SC is suspected.
- If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different site of action, if available.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information and to report suspected resistance, contact Jiangsu Agrochem Laboratory Co., Ltd. at 1-306-676-8888.

Section 9: Application Precautions**GROUND APPLICATION:**

1. Use well-maintained and calibrated conventional spray equipment, which provides adequate and uniform coverage.
2. Apply JSAL Prothio 480 SC in a minimum of 100 L water/ha, unless otherwise specified in DIRECTIONS FOR USE, Section 7.
3. Spray screens should be no finer than 50 mesh.
4. Ensure that the by-pass line discharges at the bottom of the tank to minimize foaming.
5. Maintain pressure at no less than 275 kPa to ensure good foliage penetration and coverage.
6. Provincial spray buffer zones that are greater than the spray buffer zones indicated in the DIRECTIONS FOR USE, Section 7, should be respected.

7. If spray mixture remains in the tank overnight, or for long periods during the day, agitate thoroughly prior to application.

Section 10: Mixing Instructions

Mixing Instructions:

JSAL Prothio 480 SC must be applied with properly calibrated, clean equipment. Prior to adding JSAL Prothio 480 SC to the spray tank, ensure that the spray tank is thoroughly clean.

JSAL Prothio 480 SC applied alone:

Add one-half of the required amount of water to the spray or mixing tank and start agitation.

Add the required quantity of **JSAL Prothio 480 SC** to the water and complete filling with water to the required total volume.

Maintain agitation throughout mixing and spraying.

JSAL Prothio 480 SC should be thoroughly dispersed prior to the addition of a non-ionic surfactant. JSAL Prothio 480 SC is recommended to be used with a registered non-ionic surfactant, such as Agral 90 or AgSurf, at 0.125% vol/vol.

Section 11: Pre-harvest Intervals

Crop	PHI (days)
Wheat (spring, durum and winter), Barley, Oats, Pearl Millet, Proso Millet, Rye, Triticale, Teosinte, Buckwheat	30
Canola, Rapeseed, Oriental Mustard, <i>Brassica carinata</i> , Flax (Linseed), Crambe, borage	36
Corn (field, sweet, popcorn)	14
Chickpea, Lentils	7
Soybean	20
Sugar Beet	7
Bushberry	7
Peanut	14
Cucurbits	7
Low-Growing Berries (except strawberries), Safflower, Sunflower	45

Section 12: Notices

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

For more information contact:

Jiangsu Agrochem Laboratory Co., Ltd.

No. 98, Minjiang Rd, Hi-tech Development Zone,
Changzhou, Jiangsu, 213022, China

All other products mentioned are trademarks of their respective companies.