

2026-1213

2026-04-07

****Container Label****

GROUP	3	FUNGICIDE
-------	----------	-----------

JSAL ProthioTeb 250 EC

FUNGICIDE
EMULSIFIABLE CONCENTRATE

FOR CONTROL OR SUPPRESSION OF LISTED DISEASES IN WHEAT, BARLEY,
OATS AND TRITICALE

COMMERCIAL

WARNING – EYE AND SKIN IRRITANT

REGISTRATION NUMBER.....35885 PEST CONTROL PRODUCTS ACT

ACTIVE INGREDIENT:

Prothioconazole 125 g/L

Tebuconazole 125 g/L

READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

NET CONTENTS: 1 L to BULK

Jiangsu Agrochem Laboratory Co., Ltd.

No. 98, Minjiang Rd, Hi-tech Development Zone,

Changzhou, Jiangsu, 213022, China

Tel: +86-519-8822 5302, Fax: +86-519-8510 8097

Canadian Representative
HARVESTGUARD AGRONOMY LTD.
88 Saskatchewan Street East
Moose Jaw, Saskatchewan S6H 0V4
306-676-8888



In case of spills, poisoning or fire, telephone emergency response number 911

2026-1213

2026-04-07

JSAL ProthioTeb 250 EC Fungicide is a broad-spectrum systemic fungicide for the control or suppression of listed diseases on wheat, barley, oats and triticale.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Causes eye irritation. DO NOT get in eyes. May irritate the skin. Avoid contact with skin. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum/tobacco, smoking/vaping, or using the toilet. If the product comes in contact with clothing, remove all contaminated clothing, wash skin with soap and water and dress in clean clothing. Launder contaminated clothing regularly and separately from other laundry. Aircraft cockpit and vehicle cabs must be decontaminated regularly.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours after application.

Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

DO NOT allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted. It is desirable that the pilot has communication capabilities at each treatment site at the time of application.

If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact Jiangsu Agrochem Laboratory Co., Ltd. at +1-306-676-8888.

PERSONAL PROTECTIVE EQUIPMENT: Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair. Gloves and coveralls are not required during application within a closed cab or cockpit.

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person. **If in eyes:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

2026-1213

2026-04-07

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Treat symptomatically.

ENVIRONMENTAL PRECAUTIONS: Toxic to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE. To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

STORAGE:

- Store this product away from food or feed.
- Do not contaminate water, food, or feed by storage or disposal.
- Keep away from fire or open flame or other sources of heat.
- Do not store at temperatures below freezing.
- If stored for 1 year or longer, shake well before using.
- Store the tightly closed container away from feeds, seeds, fertilizer, plants and foodstuffs.
- Do not use or store in or around the home.
- Keep in original container during storage.
- In case of fire, leaking or damaged containers, call toll free 911.

DISPOSAL:

Recyclable Container: DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple - or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

Returnable Container: DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable Container: For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

2026-1213

2026-04-07

Disposal of Unused, Unwanted Product:

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for cleanup of spills.

Disposal of Unused Spray Solution:

If any spray solution remains in the tank after spraying is finished, it should be sprayed on the perimeter of the area just sprayed, NOT ON THE CROP, away from water supplies, ditches, and irrigation canals. Spray buffer zones indicated in the DIRECTIONS FOR USE must also be respected.

2026-1213

2026-04-07

GROUP	3	FUNGICIDE
-------	----------	-----------

****Booklet Label****

JSAL ProthioTeb 250 EC

FUNGICIDE
EMULSIFIABLE CONCENTRATE

FOR CONTROL OR SUPPRESSION OF LISTED DISEASES IN WHEAT, BARLEY, OATS
AND TRITICALE.

COMMERCIAL

WARNING – EYE AND SKIN IRRITANT

REGISTRATION NUMBER..... 35885 PEST CONTROL PRODUCTS ACT

ACTIVE INGREDIENT:

Prothoiconazole 125g/L

Tebuconazole 125g/L

READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN NET CONTENTS: 1 L to BULK

Jiangsu Agrochem Laboratory Co., Ltd.

No. 98, Minjiang Rd, Hi-tech Development Zone,

Changzhou, Jiangsu, 213022, China

Tel: +86 519-8822 5302, Fax: +86-519-8510 8097

Canadian Representative
HARVESTGUARD AGRONOMY LTD.
88 Saskatchewan Street East
Moose Jaw, Saskatchewan S6H 0V4
306-676-8888



Email: info@hgagro.ca Phone: 306-676-8888 In case of spills, poisoning or fire, telephone
emergency response number 911

2026-1213

2026-04-07

JSAL ProthioTeb 250 EC

Table of Contents:

Section Number:

GENERAL INFORMATION

Notices	1
The Product	2

SAFETY AND HANDLING

Precautions, Protective Clothing & Equipment, and Re-entry Restriction	3
First Aid and Toxicological Information	4
Environmental Precautions	5
Storage	6
Disposal	7

DIRECTIONS FOR USE

Crop, Disease, Rate, Timing	8
Use Limitations	9
Pre-harvest Intervals	10
Mixing Instructions	11
Application Precautions	12
Resistance Management Recommendations	13

For more information, contact:

Jiangsu Agrochem Laboratory Co., Ltd.

No. 98, Minjiang Rd, Hi-tech Development Zone,

Changzhou, Jiangsu, 213022, China

Tel: +86 519-8822 5302, Fax: +86-519-8510 8097

2026-1213

2026-04-07

GENERAL INFORMATION

Section 1: Notices

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

Section 2: The Product

JSAL ProthioTeb 250 EC is a broad-spectrum systemic fungicide for the control or suppression of listed diseases on wheat, barley, oats and triticale.

SAFETY AND HANDLING

Section 3: Precautions, Protective Clothing & Equipment, and Re-entry Restriction

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Causes eye irritation. DO NOT get in eyes. May irritate the skin. Avoid contact with skin.

Wash thoroughly with soap and water after handling and before eating, drinking, or smoking. If product comes in contact with clothing, remove all contaminated clothing, wash skin with soap and water and dress in clean clothing. Launder applicator clothing separate from other laundry.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours after application.

Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact Jiangsu Agrochem Laboratory Co., Ltd. at +86-519-8822 5302.

PERSONAL PROTECTIVE EQUIPMENT:

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair. Gloves and coveralls are not required during application within a closed cab or cockpit.

Section 4: First Aid and Toxicological Information
--

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Treat symptomatically.

Section 5: ENVIRONMENTAL PRECAUTIONS

Toxic to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE. To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

Section 6: STORAGE

Storage:

- Store this product away from food or feed.
- Do not contaminate water, food, or feed by storage or disposal.
- Keep away from fire or open flame or other sources of heat.
- Do not store at temperatures below freezing.
- If stored for 1 year or longer, shake well before using.
- Store the tightly closed container away from feeds, seeds, fertilizer, plants and foodstuffs.
- Do not use or store in or around the home.
- Keep in original container during storage.
- In case of fire, leaking or damaged containers, call 911.

Section 7: DISPOSAL

Recyclable Container: DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple - or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

Returnable Container: DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable Container: For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

Disposal of Unused, Unwanted Product:

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for cleanup of spills.

Disposal of Unused Spray Solution:

If any spray solution remains in the tank after spraying is finished, it should be sprayed on the perimeter of the area just sprayed, NOT ON THE CROP, away from water supplies, ditches, and irrigation canals. Buffer zones indicated in the DIRECTIONS FOR USE must also be respected.

DIRECTIONS FOR USE

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Introduction

JSAL ProthioTeb 250 EC is a broad-spectrum systemic fungicide for control of a wide range of diseases on wheat (spring, winter and durum), barley, oats and triticale (spring and winter). JSAL ProthioTeb 250 EC will protect the crop from yield and quality losses due to disease.

2026-1213

2026-04-07

Section 8: Crop, Disease, Rate and Timing			
USE DIRECTIONS			
CROP	DISEASE	RATE (mL/ha)	REMARKS
Wheat, (spring, winter and durum)	Suppression of: Fusarium Head Blight (<i>Gibberella zeae</i> / <i>Fusarium graminearum</i>) Control of: Rust (Leaf, Stem and Stripe) (<i>Puccinia recondita</i> , <i>Puccinia graminis</i> , <i>Puccinia striiformis</i>) Leaf & Glume Blotch (<i>Septoria tritici</i> , <i>Stagonospora nodorum</i>) Tan Spot (<i>Pyrenophora tritici-repentis</i>) Powdery mildew (<i>Erysiphe graminis</i>)	800	For suppression of Fusarium Head Blight, apply JSAL ProthioTeb 250 EC as a preventative spray within the time period from when at least 75% of the wheat heads on the main stem are fully emerged to when 50% of the heads on the main stem are in flower. Application at this timing will also control the listed leaf diseases. Apply by ground or aerial application equipment. For ground application, apply specified dosage in a minimum of 100 L of water per hectare. For aerial application, apply specified dosage in a minimum of 50 L of water per hectare.
Barley	Suppression of: Fusarium Head Blight (<i>Fusarium spp.</i>) Control of: Net Blotch (<i>Pyrenophora teres</i>) Spot Blotch (<i>Cochliobolus sativus</i>) Scald (<i>Rhynchosporium secalis</i>) Leaf Blotch (<i>Septoria passerinii</i>) Leaf, Stem and Stripe Rusts (<i>Puccinia hordei</i> , <i>Puccinia graminis</i> , <i>Puccinia striiformis</i>) Powdery Mildew (<i>Erysiphe graminis</i>)	800	For suppression of Fusarium head blight, apply JSAL ProthioTeb 250 EC as a preventative spray within the time period when 70 to 100% of the barley main stem heads are fully emerged, to 3 days after full head emergence. Application at this timing will also control the listed leaf diseases. Apply by ground or aerial application equipment. For ground application, apply specified dosage in a minimum of 100 L of water per hectare. For aerial application, apply specified dosage in a minimum of 50 L of water per hectare.

2026-1213

2026-04-07

Oats	Control of: Crown Rust (<i>Puccinia coronata</i>) Stem Rust (<i>Puccinia graminis</i>) Stagonospora (<i>Septoria</i>) leaf blotch and black stem (<i>Stagonospora avenae</i> syn. <i>Septoria avenae</i>)	800	Apply JSAL ProthioTeb 250 EC as a preventive foliar spray when the earliest disease symptoms appear on the leaves and stems. Fields should be observed closely for early disease symptoms, particularly when susceptible varieties are planted and/or under prolonged conditions favorable for disease development. Apply by ground or aerial application equipment. For ground application, apply specified dosage in a minimum of 100 L of water per hectare. For aerial application, apply specified dosage in a minimum of 50 L of water per hectare.
Triticale (spring and winter)	Suppression of: Fusarium Head Blight (<i>Gibberella zeae</i> / <i>Fusarium graminearum</i>) Control of: Rusts (<i>Puccinia recondita</i>, <i>Puccinia graminis</i>, <i>Puccinia striiformis</i>) Leaf & Glume Blotch (<i>Septoria tritici</i>, <i>Stagonospora nodorum</i>) Tan Spot (<i>Pyrenophora tritici-repentis</i>) Powdery mildew (<i>Erysiphe graminis</i>) Spot Blotch (<i>Cochliobolus sativus</i>)	800	For suppression of Fusarium Head Blight, apply JSAL ProthioTeb 250 EC as a preventative spray within the time period from when at least 75% of the heads on the main stem are fully emerged to when 50% of the heads on the main stem are in flower. Application at this timing will also control the listed leaf diseases. Apply by ground or aerial application equipment. For ground application, apply specified dosage in a minimum of 100 L of water per hectare. For aerial application, apply specified dosage in a minimum of 50 L of water per hectare.

Resistance Management Advisory for Cereal Crops:

Repeated applications of standalone DMI fungicides should not be used on the same crop in one season against high risk pathogens, such as cereal powdery mildew, in areas of high disease pressure for that particular pathogen. Mixture products, tank-mixtures or alternation with fungicides having a different mode of action applied at effective rates have been shown to protect against the development of resistant forms of disease.

Section 9: Use Limitations

- A maximum of one application of JSAL ProthioTeb 250 EC at 800 mL product/ha may be applied per year. DO NOT apply more than 136 g tebuconazole/ha/year on wheat (spring, winter, durum), barley, oats or triticale (spring and winter) if multiple products containing tebuconazole are used.
- Applications may not be made within 36 days of harvest.
- Do not allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with JSAL ProthioTeb 250 EC
- Straw cut after harvest may be fed or used for bedding.

Field sprayer application: DO NOT apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. DO NOT apply with sprays finer than the American Society of Agricultural and Biological Engineers (ASABE) S572 (572.1 to 572.3) medium classification. Boom height must be 60 cm or less above the crop or ground.

Conventionally piloted aircraft application: DO NOT apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with sprays finer than the American Society of Agricultural and Biological Engineers (ASABE) S572 (572.1 to 572.3) medium classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length MUST NOT exceed 65% of the wing- or rotorspan.

Apply only by aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate specified for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices or a Global Positioning System (GPS).

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides. Read and understand the entire label before opening this product. If you have questions, obtain technical advice from the distributor or your provincial agricultural representative.

Product Specific PrecautionsRead and understand the entire label before opening this product. If you have questions, call the registrant at +86-519-8822 5306 or obtain technical advice from the distributor or your provincial or territorial agricultural representative. Application of this specific product must meet and/or conform to the following:

Volume: Apply the specified rate in a minimum spray volume of 50 litres per hectare for conventionally piloted aircraft.

SPRAY BUFFER ZONES

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of Application	Crop/Site		Spray Buffer Zones (metres) Required for the Protection of:				
			Freshwater Habitat of Depths:		Estuarine/Marine Habitat of Depths:		Terrestrial Habitat:
			Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer	Wheat (spring, winter and durum), barley, oats and triticale (spring and winter)		1	0	0	0	1
Aerial	Wheat (spring, winter and durum), barley, oats and triticale (spring and winter)	Fixed wing	1	0	0	0	15
		Rotary wing	1	0	0	0	10

2026-1213
2026-04-07

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Jiangsu Agrochem Laboratory Co., Ltd. at +86-519-8822 5306 for information before applying any tank mix that is not specifically recommended on this label

The spray buffer zones for this product for conventional application equipment can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Drift Mitigation portion of the Canada.ca website.

Rotational Crops:
Do not plant back within 120 days of last application.

Section 10: Pre-harvest Intervals	
Crop	PHI
Wheat (spring, winter and durum)	36 days
Barley	36 days
Oats	36 days
Triticale (spring and winter)	36 days

Section 11: Mixing Instructions

Mixing Instructions:

JSAL ProthioTeb 250 EC must be applied with properly calibrated, clean equipment. Prior to adding JSAL ProthioTeb 250 EC to the spray tank, ensure that the spray tank is thoroughly clean.

JSAL ProthioTeb 250 EC applied alone:

- Add one-half of the required amount of water to the spray or mixing tank and start agitation.
- Add the required quantity of JSAL ProthioTeb 250 EC to the water and complete filling with water to the required total volume.
- Maintain agitation throughout mixing and spraying.

Section 12: Application Precautions

GROUND APPLICATION:

1. Use well-maintained and calibrated conventional spray equipment, which provides adequate and uniform coverage.
2. Apply JSAL ProthioTeb 250 EC in a minimum of 100 L water/ha.
3. Spray screens should be no finer than 50 mesh.
4. Ensure that the by-pass line discharges at the bottom of the tank to minimize foaming.
5. Maintain pressure at no less than 275 kPa to ensure good foliage penetration and coverage.
6. Provincial spray buffer zones that are greater than the spray buffer zones indicated in the DIRECTIONS FOR USE, Section 9, should be respected.
7. If spray mixture remains in the tank overnight, or for long periods during the day, agitate thoroughly prior to application.

Section 13: Resistance Management Recommendations

For resistance management, please note that JSAL ProthioTeb 250 EC contains two Group 3 fungicides. Any fungal population may contain individuals naturally resistant to JSAL ProthioTeb 250 EC and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

2026-1213

2026-04-07

To delay fungicide resistance:

- Where possible, rotate the use of JSAL ProthioTeb 250 EC or other Group 3 fungicides with products from different fungicide groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that is effective on the target pathogen when such use is permitted.
- Fungicide use should be based on an integrated pest management (IPM) program that includes scouting, historical information related to pesticide use and crop rotation and considers cultural, biological, and other chemical control practices.
- Monitor treated fungal populations for resistance development.
- If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product and switch to another fungicide with a different target site of action, if available.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.

For further information and to report suspected resistance, contact a Jiangsu Agrochem Laboratory Co., Ltd. representative at **306-306-676-8888676-8888**

For more information contact:

Jiangsu Agrochem Laboratory Co., Ltd.

No. 98, Minjiang Rd, Hi-tech Development Zone,

Changzhou, Jiangsu, 213022, China

Tel: +86-519-8822 5302, Fax: +86-519-8510 8097