

NCS BENTAMAX480 Herbicide

For selective post-emergence broadleaf weed control in soybeans, dry and snap common beans, peas, lima beans, fababeans, corn (grain, silage, sweet and seed), flax (including low linolenic acid varieties), peanuts, blueberries, turf, spring wheat (excluding durum), snow peas, seedling and established forage legumes and seedling forage grasses (seed production only for forage legumes and grasses), newly-planted fruit trees - apple, apricot, cherry, peach, pear and nectarine (directed use only)

COMMERCIAL (AGRICULTURAL)

ACTIVE INGREDIENT: Bentazon (present as the sodium salt) 480 g/L

Solution

REGISTRATION NO. 35926 PEST CONTROL PRODUCTS ACT



CAUTION

POISON

**EYE IRRITANT
POTENTIAL SKIN SENSITIZER**

**READ THE LABEL AND ATTACHED BOOKLET BEFORE USING.
KEEP OUT OF REACH OF CHILDREN.**

NET CONTENTS: 4 L to 200 L

Northern CropScience Inc.
Suite 300, 336 Sheppard Avenue East
North York, Ontario
M2N 3B4
1-437-880-8283

FOR CHEMICAL EMERGENCY: spill, leak, fire, exposure or accident, call CHEMTREC 1-800-424-9300.

For product and use information, call Northern CropScience Inc.: 1-437-880-8283.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

- Avoid inhalation of vapour, dust or spray mist.
- Harmful if swallowed. May irritate eyes. Avoid contact with eyes, skin and clothing. Potential skin sensitizer.
- **DO NOT** eat, drink or smoke/vape while using this product. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum/tobacco, smoking/vaping, or using the toilet. Aircraft cockpit and vehicle cabs must be decontaminated regularly.
- Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks, shoes, protective eyewear (goggles or face shield) and a full-face respirator with a NIOSH-approved organic-vapour-removing cartridge (with a prefilter) approved for pesticides, or a NIOSH-approved canister approved for pesticides during mixing, loading, application, clean-up and repair. Gloves, protective eyewear and a respirator are not required during application within a closed cab or cockpit
- If the product comes in contact with clothing/PPE, remove all contaminated clothing/PPE, wash with soap and water and dress in clean clothing. Wash the outside of gloves with soap and water before removing. Launder contaminated clothing/PPE regularly and separately from other laundry.
- **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift.
- **DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Clean spray equipment thoroughly after use. When tank mixing **NCS BENTAMAX 480 Herbicide** with **PINNACLE® SG HERBICIDE**, **PINNACLE® Soybean SG Herbicide**, **PINNACLE® SG Toss-N-Go® Herbicide**, **REFLEX Liquid Herbicide** or **2,4-D amine** or **ester**, refer to **PINNACLE® SG HERBICIDE**, **PINNACLE® Soybean SG Herbicide**, **PINNACLE® SG Toss-N-Go® Herbicide**, **REFLEX Liquid Herbicide** or **2,4-D amine** or **ester** label for sprayer cleanup.
- Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

ENVIRONMENTAL PRECAUTIONS

Toxic to non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

STORAGE

1. Store this product away from food or feed. Store in original tightly closed container.
2. Do not ship or store near food, feed, seed and fertilizers.
3. Store in cool, dry, locked, well-ventilated area without floor drain.
4. Herbicides should be shipped or stored separately from other pesticides to avoid cross-contamination.
5. Freezing will not harm **NCS BENTAMAX 480 Herbicide**. Should product freeze, warm to room temperature and shake well before using.

DISPOSAL

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

NCS BENTAMAX 480 Herbicide

For selective post-emergence broadleaf weed control in soybeans, dry and snap common beans, peas, lima beans, fababeans, corn (grain, silage, sweet and seed), flax (including low linolenic acid varieties), peanuts, blueberries, turf, spring wheat (excluding durum), snow peas, seedling and established forage legumes and seedling forage grasses (seed production only for forage legumes and grasses), newly-planted fruit trees - apple, apricot, cherry, peach, pear and nectarine (directed use only).

COMMERCIAL (AGRICULTURAL)

ACTIVE INGREDIENT: Bentazon (present as the sodium salt) 480 g/L

Solution

REGISTRATION NO. 35926 PEST CONTROL PRODUCTS ACT



CAUTION

POISON

**EYE IRRITANT
POTENTIAL SKIN SENSITIZER**

**READ THE LABEL AND BOOKLET BEFORE USING.
KEEP OUT OF REACH OF CHILDREN.**

NET CONTENTS: 4 L to 200 L

Northern CropScience Inc.
Suite 300, 336 Sheppard Avenue East
North York, Ontario
M2N 3B4
1-437-880-8283

FOR CHEMICAL EMERGENCY: spill, leak, fire, exposure or accident, call CHEMTREC 1-800-424-9300.

For product and use information, call Northern CropScience Inc.: 1-437-880-8283.

GENERAL INFORMATION

NCS BENTAMAX 480 Herbicide is a liquid herbicide for selective post-emergence control of many broadleaf weeds, yellow nutsedge, cleavers, stork's bill and volunteer canola in soybeans, dry and snap common beans, peas, lima beans, fababeans, corn (grain, silage, sweet and seed), flax (including low linolenic acid varieties), peanuts, spring wheat (excluding durum), snow peas, seedling forages (alfalfa*, red clover**, alsike clover**, sainfoin**, brome grass**, creeping red fescue**, meadow foxtail**, orchardgrass**, timothy and crested wheatgrass**), established forages (alfalfa*, red clover*, sainfoin* and sweet clover*) and newly-planted fruit trees - apple, apricot, cherry, peach, pear and nectarine. (Directed use only. Do not overspray.)

NCS BENTAMAX 480 Herbicide will also control yellow nutsedge in blueberries (directed spray only) and turf (sod farms and golf courses only).

* For seed production only.

** For seed production only in Western Canada.

NCS BENTAMAX 480 Herbicide does not control grasses.

NCS BENTAMAX 480 Herbicide is a herbicide with mainly contact action. Uptake into the plant occurs primarily through the leaves. Thorough coverage of foliage is important for consistent weed control. Failure to penetrate crop or weed leaf canopies with the spray will result in incomplete control of small weeds growing underneath.

Cool weather conditions or drought will delay herbicidal activity and if prolonged, may result in poor weed control. Use of **NCS BENTAMAX 480 Herbicide** in hot, humid weather may result in temporary leaf yellowing, leaf flecking, bronzing or burning. The crop usually outgrows this condition within 10 days (see Restrictions and Limitations).

DIRECTIONS FOR USE

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Timing of Application

Apply **NCS BENTAMAX 480 Herbicide** when broadleaf weeds are small and actively growing and before the weeds reach the maximum size recommended for treatment as listed in the **NCS BENTAMAX 480 Herbicide** Weed Control and Application Rate Table. **NCS BENTAMAX 480 Herbicide** should be applied when the main weed of concern is in the suggested growth stage for treatment.

Early treatment of weeds with **NCS BENTAMAX 480 Herbicide** is important to maximize crop yield potential through elimination of early weed competition and permits optimum coverage of the weeds, thereby increasing the performance of **NCS BENTAMAX 480 Herbicide**.

Although the timing of application should be primarily in relation to the stage of weed growth, the crop must be in a tolerant stage as shown below. The information on usual stage of crop for optimum weed control may also serve as a guide to spray timing.

Blueberries

Apply **NCS BENTAMAX 480 Herbicide** at the rate of 1.75 L/ha plus **ASSIST® OIL CONCENTRATE** not to exceed 2 L/ha when yellow nutsedge is 15-20 centimetres tall.

Use as a directed spray only. Do not overspray.

Apply by ground in sufficient water volume of 100-400 L of water per hectare at 140 kPa spray pressure. Repeat application in 7-10 days but do not apply **NCS BENTAMAX 480 Herbicide** more than two times per year.

NCS BENTAMAX 480 Herbicide may not be applied within 25 days of harvest.

Snow Peas

Apply **NCS BENTAMAX 480 Herbicide** at the rate of 1.75 L/ha plus **ASSIST OIL CONCENTRATE** at 1 to 2 L/ha once a year at or after 3-leaf stage of the crop.

Apply by ground in sufficient water volume of 300 L of water per hectare.

Do not apply **NCS BENTAMAX 480 Herbicide** within 30 days of harvest.

Turf

For optimum control in turf, treat when nutsedge is young and actively growing. An additional application may be necessary at an interval of 10 to 14 days. Do not apply more than two applications of **NCS BENTAMAX 480 Herbicide** per year. For best results, do not mow grass 3 to 5 days before or after application. Do not treat newly-seeded turf until seedlings are well established as injury may result.

ADDITIVES

Use **ASSIST® OIL CONCENTRATE** or **XA™ OIL CONCENTRATE** at the rate of 1.0 to 2.0 L/ha (depending on climatic conditions and the spray volume used) for improved control of broadleaf weeds and yellow nutsedge on all crops. Either **CITOWETT PLUS** or **ASSIST OIL CONCENTRATE** or **XA OIL CONCENTRATE** can be used on peas. Refer to the table in the Spraying Instructions section.

Additives in Soybeans Only

Add 6 L/ha ammonium sulphate or 10 L/ha 28% urea ammonium nitrate (UAN) for improved and more consistent control of velvetleaf and lamb's-quarters in soybeans. The addition of either form of nitrogen source to **NCS BENTAMAX 480 Herbicide** may cause slight leaf burn, but the new growth is normal and crop vigour is not reduced.

A high quality source of 28% UAN or ammonium sulphate should be used to avoid solids or contaminants that may interfere with spray application (clogging nozzle tips) or reduced product performance. Do not use brass or aluminum nozzles when spraying **NCS BENTAMAX 480 Herbicide** plus a nitrogen source.

Note: Do not add nitrogen source when tank mixing **NCS BENTAMAX 480 Herbicide** with **PINNACLE® SG HERBICIDE** or **PINNACLE® Soybean SG Herbicide** or **PINNACLE® SG Toss-N-Go® Herbicide** and **ASSIST OIL CONCENTRATE** or **XA OIL CONCENTRATE**.

Additive in Dry Beans (pinto, great northern, pink and small red)

The addition of ammonium sulphate at 1.5% v/v will result in more consistent weed control. The addition of ammonium sulphate may cause some leaf burn, but new growth is normal and yield

is not reduced. The potential for leaf burn is increased when relative humidity and temperature are high. Use with **ASSIST OIL CONCENTRATE**.

Do not apply **NCS BENTAMAX 480 Herbicide + ASSIST OIL CONCENTRATE** or **XA OIL CONCENTRATE** with the addition of either 28% **UREA AMMONIUM NITRATE** or **AMMONIUM SULPHATE** by air.

TANK MIXING

This product may be tank mixed with a fertilizer, a supplement or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Northern CropScience Inc. at 1-437-880-8283 for information before mixing any pesticide that is not specifically recommended on this label.

Soybeans

NCS BENTAMAX 480 Herbicide can be tank mixed with **PINNACLE® SG HERBICIDE** or **PINNACLE® Soybean SG Herbicide** or **PINNACLE® SG Toss-N-Go® Herbicide**, plus **ASSIST OIL CONCENTRATE** or **XA OIL CONCENTRATE** (in soybeans only) for improved control of lamb's-quarters and redroot pigweed. For further details refer to the **NCS BENTAMAX 480 Herbicide** Weed Control and Application Rate Table.

Dry Common Beans

NCS BENTAMAX 480 Herbicide can be tank mixed with **REFLEX® Liquid Herbicide** plus **ASSIST OIL CONCENTRATE** or **XA OIL CONCENTRATE** for suppression of redroot pigweed and control of lamb's-quarters, common ragweed and lady's-thumb. For further details, refer to the **NCS BENTAMAX 480 Herbicide** Weed Control and Application Rate Table.

NCS BENTAMAX 480 Herbicide and **REFLEX Liquid Herbicide** tank mix (ground application only) is registered in dry common beans only.

Dry Edible Beans in the Red River Valley of Manitoba

NCS BENTAMAX 480 Herbicide can be tank mixed with **REFLEX Liquid Herbicide** for selective post-emergence broadleaf weed control in dry edible beans in the Red River Valley of Manitoba. See **REFLEX Liquid Herbicide** label for weeds controlled in addition to those listed on the **NCS BENTAMAX 480 Herbicide** label.

Apply **NCS BENTAMAX 480 Herbicide** at the rate of 1.75 L/ha plus **REFLEX Liquid Herbicide** at 0.58 L/ha plus **AGRAL® 90** at 0.10% v/v at 1 to 2 trifoliate leaf stage of the crop and 3 to 4 leaf stage of the weeds.

Apply once a year, post-emergent and by ground only.

Do not apply this tank mix within 84 days of harvest.

Spring Wheat (excluding durum)

When tank mixed with **2,4-D amine** or **ester** formulation, **NCS BENTAMAX 480 Herbicide** will control lady's-thumb, bluebur, burdock (< 4-leaf), cocklebur, common plantain, daisy fleabane, false flax, flixweed, goat's beard, lamb's-quarters, mustard (except dog and tansy), prickly lettuce,

ragweed, redroot pigweed, Russian pigweed, Russian thistle, shepherd's purse, stinging nettle, stinkweed, sweet clover, volunteer canola, wild radish and wild sunflower.

All of the above-mentioned broadleaf weeds will be controlled if applied from the weed 2-4 leaf stage. For further details, refer to the **NCS BENTAMAX 480 Herbicide** Weed Control and Application Rate Table.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, **NCS BENTAMAX 480 Herbicide** is a Group 6 herbicide. Any weed population may contain or develop plants naturally resistant to **NCS BENTAMAX 480 Herbicide** and other Group 6 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of **NCS BENTAMAX 480 Herbicide** or other Group 6 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other chemical control practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative field options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance management and/or integrated weed management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Northern CropScience Inc. at 1-437-880-8283.

SPRAYING INSTRUCTIONS

Ground Application

Use sprayers equipped with standard flat fan pesticide nozzles with the recommended spray volume, pressure and additives. Tilt spray nozzles 45 degrees forward to ensure better coverage.

SPRAY VOLUME	SPRAY PRESSURE	ADDITIVES
Minimum 100 L/ha. Use larger water volumes (up to	Minimum 275 kPa. Use higher pressure (up to 425	Use 1 litre of ASSIST OIL CONCENTRATE or XA OIL

400 L/ha) for weeds at the upper limit of their recommended stage of treatment.	kPa) for weeds at the upper limit of their recommended stage for treatment.	CONCENTRATE per 100 litres of water with a maximum application rate of 2 L/ha. Under hot, humid conditions, restrict ASSIST OIL CONCENTRATE or XA OIL CONCENTRATE rate to 1 L/ha. Alternatively, CITOWETT PLUS may be used on peas only at a rate of 2.5 L per 1000 L of spray solution.
---	---	--

TIMING OF APPLICATION RATE TABLE

CROP	TOLERANT STAGE	USUAL STAGE OF CROP FOR OPTIMUM WEED CONTROL
Soybeans	Tolerant at any growth stage	Unifoliolate to two expanded trifoliolate leaves, usually 18-20 days after planting
Lima beans, dry common beans ¹ (<i>Phaseolus vulgaris</i> only - including but not limited to: white, kidney, black, pinto, great northern, pink, small red, cranberry and otebo) and most snap common beans including snap beans	Tolerant after 1st trifoliolate leaf has fully expanded	1 to 3 trifoliolate leaves
Corn (grain, silage, sweet, seed)	Tolerant at any growth stage	1 to 5-leaf stage
Peas (field and processing)	Tolerant after 3 pairs of leaves (or 3 nodes) are present	Soon after 3 pairs of leaves form
Fababeans	Tolerant after 2 to 3-leaf stage or crop is 10 cm high	Soon after 3-leaf stage
Peanuts	Tolerant at any growth stage	Unifoliolate to two expanded trifoliolate leaves
Flax (including low linolenic acid varieties)	Tolerant when crop is 5 cm or higher	Soon after crop reaches 5 cm
Spring wheat (excluding durum)	Tolerant at any growth stage	2 to 4 leaves
Turf (sod farms and golf courses) for yellow nutsedge control only	Tolerant on established turf	When turf is actively growing in late spring and early summer
Blueberries for yellow nutsedge control only	Tolerant as a directed spray only. Do not overspray.	
Snow peas		Soon after 3-leaf stage

¹ Dry common bean varieties may vary in their tolerance to herbicides, including **NCS BENTAMAX 480 Herbicide**. Since not all dry common bean varieties have been tested for tolerance to **NCS BENTAMAX 480 Herbicide**, first use of **NCS BENTAMAX 480 Herbicide** should be limited to a small area of each variety to confirm tolerance prior to adoption as a general field practice. Additionally, consult your seed supplier for information on the tolerance of specific varieties of dry common beans to **NCS BENTAMAX 480 Herbicide**.

Newly-planted fruit trees (apple, apricot, cherry, peach, pear, and nectarine)	Tolerant as a directed spray only. Do not overspray.	
--	--	--

TIMING OF APPLICATION RATE TABLE (Continued)

CROP	TOLERANT STAGE	USUAL STAGE OF CROP FOR OPTIMUM WEED CONTROL
Seed Production¹ for annual broadleaf weed control		
Forage Grasses (Seedling) • Bromegrass • Creeping Red Fescue • Meadow Foxtail • Orchardgrass • Timothy • Crested Wheatgrass	Tolerant at 1- 7-leaf growth stage ³	2 - 5-leaf of seedling forage grasses
Forage Legumes (Seedling) • Alfalfa • Red Clover • Alsike Clover • Sainfoin	Tolerant after third trifoliate stage ³	3- 5 trifoliate. Approximately 4-6 weeks after planting.
Forage Legumes (Established) • Alfalfa²	Tolerant before crop canopy closes, prior to flowering ³	
Forage Legumes (Established) ² • Red clover • Sainfoin • Sweet clover	Tolerant between 7.5 and 25 cm high ³	After crop is 7.5 cm and before crop canopy closes.

¹ All crops listed for seed production are Western Canada only with the exception of alfalfa and established sainfoin, sweet clover and red clover.

² For seed production, only one application of **NCS BENTAMAX 480 HERBICIDE** per season is recommended. Crop injury may occur under hot, humid conditions. Speed of recovery will be influenced by growing conditions and weed control. (**SEE RESTRICTIONS AND LIMITATIONS SECTION**)

³ Some leaf scorch may appear but the effect is transient and will outgrow within 3-4 weeks.

NCS BENTAMAX 480 HERBICIDE WEED CONTROL AND APPLICATION RATE TABLE

WEED SPECIES	RECOMMENDED TREATMENT HEIGHT (CM) FOR WEEDS	TOO LATE FOR BEST RESULTS	
		HEIGHT (CM)	LEAF STAGE
NCS BENTAMAX 480 HERBICIDE at 2.25 L/ha + ASSIST OIL CONCENTRATE or XA OIL CONCENTRATE at 1-2 L/ha			
Bird rape*	5.0 - 10.0	over 10	6-leaf
Buttercup	5.0 - 10.0	over 10	
Cleavers	4.0 - 8.0	over 8	1-3 whorl
Cocklebur	17.5 - 30.0	over 30	10-leaf
Common chickweed	1-3 weeks after emergence	Later than 3 weeks after emergence	
Common groundsel*	5.0 - 10.0	over 10	6-leaf
Common ragweed*	2.5 - 5.0	over 5	
Corn spurry	2.5 - 10.0	over 10	
Flower-of-an hour	5.0 - 10.0	over 10	
Giant ragweed	5.0 - 15.0	over 15	4-leaf
Hairy galinsoga	5.0 - 7.5	over 7.5	
Hairy nightshade	0.5 - 2.0	over 2	6-leaf
Jimsonweed	5.0 - 15.0	over 15	10-leaf
Lady's-thumb (smartweed)	7.5 - 20.0	over 20	10-leaf
Lamb's-quarters*	1.25 - 2.5	over 2.5	8-leaf
Low cudweed	2.5 - 5.0	over 5	6-leaf
Purslane	2.5 - 5.0	over 5	6-leaf
Redroot pigweed* (suppression only)	1.25 - 3.75	over 3.75	4-leaf
Russian thistle	2.5 - 7.5	over 7.5	
Shepherd's-purse	10.0 - 25.0	over 25	6-leaf
Stinkweed	5.0 - 15.0	over 15	6-leaf
Stork's bill	4.0 - 10.0	over 10	2-6-leaf
Velvetleaf**	10.0 - 15.0	over 15	6-leaf
Volunteer canola	2.0 - 15.0	over 15	8-leaf
Wild mustard	12.5 - 25.0	over 25	10-leaf
Wild radish	2.0 - 5.0	over 5	6-leaf
SOYBEANS ONLY: NCS BENTAMAX 480 HERBICIDE at 2.25 L/ha + PINNACLE SG HERBICIDE or PINNACLE Soybean SG Herbicide or PINNACLE SG Toss-N-Go Herbicide at 8.25 to 12.0 g/ha + ASSIST OIL CONCENTRATE or XA OIL CONCENTRATE at 1-2 L/ha			
Controls all the above listed weed species with the following improvements:			
+Lamb's-quarters*	1.25 - 10.0	over 10	8-leaf
Redroot pigweed*	1.25 - 10.0	over 10	6-leaf
NCS BENTAMAX 480 HERBICIDE at 1.75 L/ha + ASSIST OIL CONCENTRATE or XA OIL CONCENTRATE at 1-2 L/ha			
Bird rape*	2.5 - 5.0	over 5	
Cocklebur	7.5 - 17.5	over 17.5	
Flower-of-an-hour	2.5 - 5.0	over 5	
Lady's-thumb (smartweed)	2.5 - 7.5	over 7.5	
Shepherd's-purse	Rosette - 10.0	over 10	
Stinkweed	Rosette - 5.0	over 5	
Velvetleaf**	5.0 - 10.0	over 10	
Volunteer canola	2.0 - 15.0	over 15	8-leaf
Wild mustard	2.5 - 12.5	over 12.5	

+ Use 12.0 g/ha rate of **PINNACLE SG HERBICIDE** or **PINNACLE Soybean SG Herbicide** or **PINNACLE SG Toss-N-Go Herbicide** for advanced leaf staging.

++ Second flushes of these weeds will not be controlled.

* Triazine resistant strains of these weeds are controlled by **NCS BENTAMAX 480 HERBICIDE**

** **NCS BENTAMAX 480 HERBICIDE** will defoliate velvetleaf 15 cm or taller, but regrowth may occur.

*** Treat field bindweed before it is dark green and has begun to trail.

NCS BENTAMAX 480 HERBICIDE WEED CONTROL AND APPLICATION RATE TABLE Continued

NCS BENTAMAX 480 HERBICIDE WEED CONTROL AND APPLICATION RATE TABLE (Continued)			
WEED SPECIES	RECOMMENDED TREATMENT HEIGHT (CM) FOR WEEDS	TOO LATE FOR BEST RESULTS	
		HEIGHT (CM)	LEAF STAGE
DRY COMMON BEANS ONLY: NCS BENTAMAX 480 HERBICIDE at 1.75 L/ha + REFLEX Liquid Herbicide at 0.58 L/ha + ASSIST OIL CONCENTRATE or XA OIL CONCENTRATE at 2 L/ha			
Controls all the above listed weed species with the following improvements:			
++Common ragweed*	1.25 - 5.0	Over 5	6-leaf
++Lamb's-quarters*	1.25 - 2.5	over 2.5	8-leaf
++Redroot Pigweed* (suppression only)	1.25 - 3.75	over 3.75	4-leaf
SOYBEANS ONLY: NCS BENTAMAX 480 HERBICIDE at 1.75 L/ha + PINNACLE SG HERBICIDE or PINNACLE Soybean SG Herbicide or PINNACLE SG Toss-N-Go Herbicide at 8.25 to 12.0 g/ha + ASSIST OIL CONCENTRATE or XA OIL CONCENTRATE at 1-2 L/ha			
Controls all the above listed weed species with the following improvements:			
+Lamb's-quarters*	1.25 - 10.0	Over 10	8-leaf
Redroot Pigweed*	1.25 - 10.0	over 10	6-leaf
PERENNIAL WEEDS: NCS BENTAMAX 480 HERBICIDE at 1.75 L/ha + ASSIST OIL CONCENTRATE or XA OIL CONCENTRATE at 1-2 L/ha			
Repeat application 7 to 15 days after 1st application (if necessary):			
Canada thistle	15 - 20	Over 20	
Field bindweed*** (suppression only)	2.5 - 6	over 6	
Yellow nutsedge	15 - 20	over 20	
SPRING WHEAT (excluding durum): NCS BENTAMAX 480 HERBICIDE at 1.0 L/ha + 2,4-D amine or ester (470 g/L) at 0.75-1.0 L/ha			
WEEDS		RECOMMENDED APPLICATION TIMING	
Lady's-thumb		2 - 4 leaf	
Bluebur		2 - 4 leaf	
Burdock		< 4 leaf	
Cocklebur		2 - 4 leaf	
Common plantain		2 - 4 leaf	
Daisy fleabane		2 - 4 leaf	
False flax		2 - 4 leaf	
Flixweed		2 - 4 leaf	
Goat's beard		2 - 4 leaf	
Lamb's-quarters		2 - 4 leaf	
Mustard (except dog and tansy)		2 - 4 leaf	
Prickly lettuce		2 - 4 leaf	
Ragweed		2 - 4 leaf	
Redroot pigweed		2 - 4 leaf	
Russian pigweed		2 - 4 leaf	
Russian thistle		2 - 4 leaf	
Shepherd's purse		2 - 4 leaf	
Stinging nettle		2 - 4 leaf	
Stinkweed		2 - 4 leaf	
Sweet clover		2 - 4 leaf	
Volunteer canola		2 - 4 leaf	
Wild radish		2 - 4 leaf	
Wild sunflower		2 - 4 leaf	

+ Use 12.0 g/ha rate of PINNACLE SG HERBICIDE or PINNACLE Soybean SG Herbicide or PINNACLE SG Toss-N-Go Herbicide for advanced leaf staging.

++ Second flushes of these weeds will not be controlled.

* Triazine resistant strains of these weeds are controlled by NCS BENTAMAX 480 HERBICIDE.

** NCS BENTAMAX 480 HERBICIDE will defoliate velvetleaf 15 cm or taller, but regrowth may occur.

*** Treat field bindweed before it is dark green and has begun to trail.

NCS BENTAMAX 480 HERBICIDE BANDING TECHNIQUES

1. Spray a minimum of 25 cm wide band.
2. Minimize the amount of dust striking target weeds to ensure adequate coverage and penetration.
3. Do not use cultivation equipment when spraying.
4. Adjust the **NCS BENTAMAX 480 Herbicide** rate to proportion of the total area to be sprayed.

AIRCRAFT APPLICATIONS

DO NOT APPLY **NCS BENTAMAX 480 Herbicide** BY AIR TO: Corn, peas, fababeans, peanuts, flax, blueberries, snow peas, turf, newly-planted fruit trees, spring wheat, forage grasses and forage legumes.

Aerial applications are allowed on soybeans, dry and snap common beans only.

Apply **NCS BENTAMAX 480 Herbicide** when weeds are in the early stages of growth recommended for treatment. Crop canopy should not be so dense as to prevent spray from thoroughly covering weeds. Use spray volumes, pressure and additives as recommended below:

SPRAY VOLUME	SPRAY PRESSURE	ADDITIVES
50 to 100 L/ha	Minimum 275 kPa	0.125 to 0.25 L ASSIST OIL CONCENTRATE or XA OIL CONCENTRATE per hectare

Do not use **ASSIST OIL CONCENTRATE** or **XA OIL CONCENTRATE** in excess of 0.25 L/ha as substantial crop injury could occur.

Treat when wind is less than 8 kilometres per hour. Do not apply when weather conditions may cause spray drift from target areas to adjacent crops.

Apply only by aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate specified for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices or a Global Positioning System (GPS).

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.

Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the registrant at 1-437-880-8283 or obtain technical advice from the distributor or your provincial or territorial agricultural representative. Application of this specific product must meet and/or conform to the following volume: apply the specified rate in a minimum spray volume of 50 litres per hectare for conventionally piloted aircraft.

SPRAY DRIFT MANAGEMENT FOR GROUND AND AERIAL APPLICATION

Field Sprayer Application

DO NOT apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. DO NOT apply with sprays finer than the American Society of Agricultural and Biological Engineers (ASABE) S572 (572.1 to 572.3) Medium classification. Boom height must be 60 cm or less above the crop or ground.

Conventionally piloted aircraft application

DO NOT apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 8 km/h at flying height at the site of application. DO NOT apply with sprays finer than the American Society of Agricultural and Biological Engineers (ASABE) S572 (572.1 to 572.3) Medium classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length MUST NOT exceed 65% of the wing or rotorspan.

Spray Buffer Zones

A spray buffer zone is NOT required for:

- uses with hand-held application equipment permitted on this label,
- low-clearance hooded or shielded sprayers that prevent spray contact with crop, fruit or foliage

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive fresh water habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop	Spray Buffer Zones (metres) Required for the Protection of:				
		Freshwater Habitat of Depths:		Estuarine/Marine Habitat of Depths:		Terrestrial Habitat:
		Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	

Field sprayer	Snow peas, peas (field and processing), seedling forage grasses/legumes and established forage legumes (for seed production)		0	0	0	0	1
	Soybean, dry common beans, snap beans		0	0	0	0	2
Aerial	Soybean, dry common beans, snap beans	Fixed wing	0	0	0	0	55
		Rotary wing	0	0	0	0	45

When tank mixes are permitted, consult the labels of the tankmix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASABE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product for conventional application equipment can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Drift Mitigation portion of the Canada.ca website

MIXING

1. Clean spray tank and fill half full with clean water. Start agitation or by-pass system.
2. If required add correct amount of nitrogen source. **Note: Do not add nitrogen source when tank mixing NCS BENTAMAX 480 Herbicide with PINNACLE SG HERBICIDE, PINNACLE Soybean SG Herbicide, PINNACLE SG Toss-N-Go Herbicide, REFLEX Liquid Herbicide or 2,4-D amine or ester and ASSIST OIL CONCENTRATE or XA OIL CONCENTRATE.**
3. If required, add the correct amount of **PINNACLE SG HERBICIDE, PINNACLE Soybean SG Herbicide, PINNACLE SG Toss-N-Go Herbicide** (for use on soybeans only), **REFLEX Liquid Herbicide** (for use on dry common beans only) or **2,4-D amine or ester** (for use on spring wheat [excluding durum]). Continuous agitation is required to keep **PINNACLE SG HERBICIDE, PINNACLE Soybean SG Herbicide, PINNACLE SG Toss-N-Go Herbicide, REFLEX Liquid Herbicide or 2,4-D amine or ester** in suspension.
NOTE: On repeat tank loads, prepare a **PINNACLE SG HERBICIDE or PINNACLE Soybean SG Herbicide or PINNACLE SG Toss-N-Go Herbicide** /water slurry in a separate container with clean water before adding to spray tank.
4. Add correct amount of **NCS BENTAMAX 480 Herbicide** and agitate 2 to 3 minutes.
5. Add correct amount of **ASSIST OIL CONCENTRATE or XA OIL CONCENTRATE or CITOWETT PLUS** and agitate 2 to 3 minutes. When tank mixing **NCS BENTAMAX 480 Herbicide with PINNACLE SG HERBICIDE, PINNACLE Soybean SG Herbicide, PINNACLE SG Toss-N-Go Herbicide, REFLEX Liquid Herbicide or 2,4-D amine or ester**, only **ASSIST OIL CONCENTRATE or XA OIL CONCENTRATE** may be used.
6. Add remainder of water, agitate and spray.
7. If an oil film starts to build up in the tank, drain and clean tank with a strong detergent solution.

RESTRICTIONS AND LIMITATIONS

Do not treat any crops not listed on this label.

NOTE: Do not add nitrogen source when tank mixing NCS BENTAMAX 480 Herbicide with PINNACLE SG HERBICIDE, PINNACLE Soybean SG Herbicide, PINNACLE SG Toss-N-Go Herbicide, REFLEX Liquid Herbicide or 2,4-D amine or ester and ASSIST OIL CONCENTRATE or XA OIL CONCENTRATE.

Do not allow livestock to graze on **NCS BENTAMAX 480 Herbicide** treated turf.

Do not apply **NCS BENTAMAX 480 Herbicide** to any crops that have been subjected to stress from conditions such as hail damage, flooding, hot, humid weather, drought, widely fluctuating temperature conditions, prolonged cold weather or injury from prior herbicide applications, as crop injury may result.

Do not apply **NCS BENTAMAX 480 Herbicide** to newly-seeded turf until seedlings are well established.

Rainfall within 6 to 8 hours of application may reduce effectiveness of spray.

When **NCS BENTAMAX 480 Herbicide** is applied beyond recommended weed growth stages, limited to unsatisfactory weed control will result.

Cool weather conditions or drought will delay herbicide activity and if prolonged, may result in poor weed control.

NCS BENTAMAX 480 Herbicide and **PINNACLE SG HERBICIDE** or **PINNACLE Soybean SG Herbicide** or **PINNACLE SG Toss-N-Go Herbicide** tank mix is registered for use in soybeans only (ground application only).

NCS BENTAMAX 480 Herbicide and **REFLEX Liquid Herbicide** tank mix (ground application only) is registered in dry common beans only.

NCS BENTAMAX 480 Herbicide and **2,4-D amine** or **ester** tank mix is registered for use in spring wheat (excluding durum) only. **Do not apply this tank mix by air.** A fifty (50) day pre-harvest interval is required after application, before harvesting the treated crop.

Do not apply **NCS BENTAMAX 480 Herbicide** within 36 days of harvest of red clover grown for seed production.

WARNING

Do not apply **NCS BENTAMAX 480 Herbicide** when weather conditions may cause spray drift from treated areas to adjacent crops.

Lentils, adzuki and mung beans, cucumbers, sugar beets and sunflowers can be injured by **NCS BENTAMAX 480 Herbicide**.

ENVIRONMENTAL PRECAUTIONS

Toxic to non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

1. Avoid inhalation of vapour, dust or spray mist.
2. Harmful if swallowed. May irritate eyes. Avoid contact with eyes, skin and clothing. Potential skin sensitizer.
3. **DO NOT** eat, drink or smoke/vape while using this product. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum/tobacco, smoking/vaping, or using the toilet. Aircraft cockpit and vehicle cabs must be decontaminated regularly.
4. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks, shoes, protective eyewear (goggles or face shield) and a full-face respirator with a NIOSH-approved organic-vapour-removing cartridge (with a prefilter) approved for pesticides, or a NIOSH-approved canister approved for pesticides during mixing, loading, application, clean-up and repair. Gloves, protective eyewear and a respirator are not required during application within a closed cab or cockpit.
5. If the product comes in contact with clothing/PPE, remove all contaminated clothing/PPE, wash with soap and water and dress in clean clothing. Wash the outside of gloves with soap and water before removing. Launder contaminated clothing/PPE regularly and separately from other laundry.
6. **DO NOT** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.
7. Clean spray equipment thoroughly after use. When tank mixing **NCS BENTAMAX 480 Herbicide** with **PINNACLE SG HERBICIDE**, **PINNACLE Soybean SG Herbicide**, **PINNACLE SG Toss-N-Go Herbicide**, **REFLEX Liquid Herbicide** or **2,4-D amine or ester**, refer to **PINNACLE SG HERBICIDE**, **PINNACLE Soybean SG Herbicide**, **PINNACLE SG Toss-N-Go Herbicide**, **REFLEX Liquid Herbicide** or **2,4-D amine or ester** label for sprayer cleanup.
8. Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

STORAGE

1. Store this product away from food or feed. Store in original tightly closed container.
2. Do not ship or store near food, feed, seed and fertilizers.
3. Store in cool, dry, locked, well-ventilated area without floor drain.
4. Herbicides should be shipped or stored separately from other pesticides to avoid cross-contamination.
5. Freezing will not harm **NCS BENTAMAX 480 Herbicide**. Should product freeze, warm to room temperature and shake well before using.

DISPOSAL

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

®All other products listed are registered trademarks of their respective companies.