

MEY Canada GlufoSupreme 150 SN Herbicide

GROUP	10	HERBICIDE
-------	-----------	-----------

COMMERCIAL

SOLUTION



WARNING EYE AND SKIN IRRITANT

REGISTRATION NUMBER 35941 PEST CONTROL PRODUCTS ACT

ACTIVE INGREDIENT: glufosinate- ammonium 150 g/L

NET CONTENTS:
[1L – 1000 L]

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

Distributed by:

MEY Canada Corporation
855 2nd St. S.W. Suite 3500
Calgary, AB, Canada TSP 4J8
919.338.1628

In case of spills, poisoning or fire, telephone emergency response number
911 (24 hours a day).

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Harmful or fatal if absorbed through the skin.

Harmful if swallowed. Causes eye irritation. **DO NOT** get in eyes, on skin or on clothing.

Avoid breathing spray mist. Wash thoroughly after using and before eating, drinking or smoking.

Protective Clothing and Equipment: Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes, and protective eyewear (goggles or face shield) during mixing, loading, application, clean-up and repair. Gloves and protective eyewear (goggles or face shield) are not required during application within a closed cab and/or cockpit.

Restricted-entry Restriction: Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

Apply only when the potential for drift to areas of human habitation and areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact DELL TECH LABORATORIES LTD. at 519-858-5024 or consult the website <https://meycanadacorp.ca>.

FIRST AID: Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.

TOXICOLOGICAL INFORMATION:

Medical personnel should contact 911. **Note to Physician:** If this product is ingested, endotracheal intubation and gastric lavage should be performed as soon as possible, followed by charcoal and sodium sulfate administration.

ENVIRONMENTAL PRECAUTIONS:

TOXIC to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE. Avoid spray drift to susceptible plants and **USE ONLY FOR RECOMMENDED PURPOSES AND AT RECOMMENDED RATES.**

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

STORAGE:

- CANNOT be stored below freezing.
- If stored for 1 year or longer, shake well before using.
- Store the tightly closed container away from feeds, seeds, fertilizers, plants and foodstuffs.
- Do not use or store in or around the home.
- Keep in original container during storage.
- To prevent contamination, store this product away from food or feed.

DISPOSAL:

Recyclable Container Disposal (less than 23 liters):

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

Returnable Container Disposal:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable Container Disposal:

For disposal, this empty container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

Disposal of Unused, Unwanted Product:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills.

Disposal of Unused Spray Solution:

If any spray solution remains in the tank after spraying is finished, it should be sprayed on the perimeter of the area just sprayed, away from water supplies, ditches, and irrigation canals.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

2026-0660
2026-04-02

Booklet

MEY Canada GlufosSupreme 150 SN Herbicide

GROUP	10	HERBICIDE
--------------	-----------	------------------

COMMERCIAL

SOLUTION



WARNING EYE AND SKIN IRRITANT

REGISTRATION NUMBER 35941
PEST CONTROL PRODUCTS ACT

ACTIVE INGREDIENT: glufosinate- ammonium 150 g/L

NET CONTENTS: [1L – 1000 L]

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

Distributed by:

MEY Canada Corporation
855 2nd St. S.W. Suite 3500
Calgary, AB, Canada TSP 4J8
919.338.1628

In case of spills, poisoning or fire, telephone emergency response number
911 (24 hours a day).

MEY CANADA GLUFOSUPREME 150 SN HERBICIDE

<i>Table of Contents:</i>		<i>Section Number:</i>
GENERAL INFORMATION	The Product	1
SAFETY AND HANDLING	Precautions, Protective Clothing and Equipment	2
	First Aid and Toxicological Information	3
	Environmental Precautions	4
	Storage	5
	Disposal	6
DIRECTIONS FOR USE	POST EMERGENT USE	
	Crops, Weeds, Timing and Rates	7
	Tank Mixes	8
	Primocane Control in Raspberry	9
	DESICCATION USE	
	Crops, Instructions and Rates	10
	GENERAL INSTRUCTIONS	
	Herbicide Resistance Management	
	Recommendations	11
	Application Instructions and Precautions	12
	Mixing Instructions	13
	Sprayer Cleanup	14
	Livestock Feeding and Preharvest Intervals	15
	Notices	16

GENERAL INFORMATION

Section 1: The Product

MEY Canada GlufoSupreme 150 SN Herbicide provides control of a broad spectrum of grassy and broadleaf weeds in Canola and Soybean varieties or hybrids that are specially developed to be tolerant to glufosinate ammonium.

MEY Canada GlufoSupreme 150 SN Herbicide is registered for use on glufosinate ammonium tolerant Canola and Soybean varieties or hybrids

MEY Canada GlufoSupreme 150 SN Herbicide may also be applied to glufosinate ammonium tolerant lines or varieties grown for seed production.

For hybrid seed production in canola, MEY Canada GlufoSupreme 150 SN Herbicide may be used to remove segregating wild type plants within the female population. Two applications are required to do this.

Glufosinate ammonium tolerant Corn can be used as separator rows in hybrid Canola seed production but must be destroyed and not used for food or feed.

MEY Canada GlufoSupreme 150 SN Herbicide may be used as a harvest aid (desiccant) on Alfalfa grown for seed.

MEY Canada GlufoSupreme 150 SN Herbicide will desiccate weeds that are present in the field at time of application when used as a harvest aid.

MEY Canada GlufoSupreme 150 SN Herbicide may also be used for primocane control in established Raspberries.

SAFETY AND HANDLING

Section 2: Precautions and Protective Clothing and Equipment

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Harmful or fatal if absorbed through the skin.

Harmful if swallowed. Causes eye irritation. DO NOT get in eyes, on skin or on clothing.

Avoid breathing spray mist. Wash thoroughly after using and before eating, drinking or smoking.

If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact Dell Tech Laboratories Ltd. at 1-519-858-5024.

Protective Clothing and Equipment

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes, and protective eyewear (goggles or face shield) during mixing, loading, application, clean-up and repair. Gloves and protective eyewear (goggles or face shield) are not required during application within a closed cab and/or cockpit.

When applied as a tank-mix combination, read and observe all label directions, including rates, personal protective equipment, restrictions and precautions for each product used

in the tank-mix. Always use in accordance with the most restrictive label restrictions and precautions.

Restricted-entry Restriction

~Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

Apply only when the potential for drift to areas of human habitation and areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

Section 3: First Aid and Toxicological Information

FIRST AID: Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION:

Medical personnel should contact 911. **Note to Physician:** If this product is ingested, endotracheal intubation and gastric lavage should be performed as soon as possible, followed by charcoal and sodium sulfate administration.

Section 4: Environmental Precautions

~TOXIC to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under DIRECTIONS FOR USE.

~Avoid spray drift to susceptible plants and USE ONLY FOR RECOMMENDED PURPOSES AND AT RECOMMENDED RATES.

~To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

~Avoid application when heavy rain is forecast.

~Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

Section 5: Storage

- ✗CANNOT be stored below freezing.
- ✗If stored for 1 year or longer, shake well before using.
- ✗Store the tightly closed container away from feeds, seeds, fertilizers, plants and foodstuffs.
- ✗Do not use or store in or around the home.
- ✗Keep in original container during storage.
- ✗To prevent contamination, store this product away from food or feed.

Section 6: Disposal

Recyclable Container Disposal (less than 23 litres):

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

Returnable Container Disposal:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable Container Disposal:

For disposal, this empty container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

Disposal of Unused, Unwanted Product:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for cleanup of spills.

Disposal of Unused Spray Solution:

If any spray solution remains in the tank after spraying is finished, it should be sprayed on the perimeter of the area just sprayed, away from water supplies, ditches, and irrigation canals.

DIRECTIONS FOR USE

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

POST EMERGENT USE

Section 7: Crops, Weeds, Timing and Rates

CANOLA – EASTERN CANADA AND COASTAL BC

- MEY Canada GlufosSupreme 150 SN Herbicide is registered for aerial and ground application to glufosinate ammonium tolerant Canola varieties or hybrids [
- To assure crop safety and optimal herbicide performance, only use MEY Canada GlufosSupreme 150 SN Herbicide on glufosinate ammonium tolerant Canola grown from certified seed.**
- Application of MEY Canada GlufosSupreme 150 SN Herbicide to non-tolerant Canola varieties or hybrids or other non-tolerant crops will result in severe crop injury or death of the crop.**
- Apply MEY Canada GlufosSupreme 150 SN Herbicide from the cotyledon stage up until, but prior to, the early bolting stage of Canola.
- Slight discoloration of the Canola may be visible after application. This effect is temporary and will not influence crop growth, maturity or yield.
- For application to glufosinate ammonium and glyphosate tolerant canola varieties or hybrids** MEY Canada GlufosSupreme 150 SN Herbicide should be applied in a sequential program with glyphosate for effective weed control. Tank-mixing MEY Canada GlufosSupreme 150 SN Herbicide with glyphosate may result in reduced control on certain weed species.
- MEY Canada GlufosSupreme 150 SN Herbicide may also be applied to glufosinate ammonium tolerant Canola lines or varieties grown for seed production.
- For hybrid seed production** - Two applications are required to remove the segregating wild type plants. The first application should occur when the Canola is in the 2-4 leaf stage, the second application when the Canola plants are in the 4-6 leaf stage. All applications of MEY Canada GlufosSupreme 150 SN Herbicide for hybrid seed production should be made at 3.33 L/ha.
- DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.
- Follow use instructions for canola and general instructions in this label.

CANOLA - PRAIRIE PROVINCES AND INTERIOR OF BRITISH COLUMBIA ONLY

MEY Canada GlufosSupreme 150 SN Herbicide is registered for aerial and ground application to glufosinate ammonium tolerant Canola varieties or hybrids grown in the Prairie Provinces and Interior of British Columbia only.

- To assure crop safety and optimal herbicide performance, only use MEY Canada GlufosSupreme 150 SN Herbicide on glufosinate ammonium tolerant Canola grown from certified seed.**

- **Application of MEY Canada GlufosSupreme 150 SN Herbicide to non-tolerant Canola varieties or hybrids or other non-tolerant crops will result in severe crop injury or death of the crop.**
- Apply MEY Canada GlufosSupreme 150 SN Herbicide from the cotyledon stage up until, but prior to, the early bolting stage of Canola.
- Slight discoloration of the Canola may be visible after application. This effect is temporary and will not influence crop growth, maturity or yield.
- MEY Canada GlufosSupreme 150 SN Herbicide may also be applied to glufosinate ammonium tolerant Canola lines or varieties grown for seed production.
- **For hybrid seed production** - Two applications are required to remove the segregating wild type plants. The first application should occur when the Canola is in the 2-4 leaf stage, the second application when the Canola plants are in the 4-6 leaf stage. All applications of MEY Canada GlufosSupreme 150 SN Herbicide for hybrid seed production should be made at 3.33 L/ha.
- DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.
- Follow use instructions for canola and general instructions in this label.

SOYBEAN

- MEY Canada GlufosSupreme 150 SN Herbicide is registered for aerial and ground application to glufosinate ammonium tolerant Soybean varieties
- **Application of MEY Canada GlufosSupreme 150 SN Herbicide to non-tolerant Soybean varieties or other non-tolerant crops will result in severe crop injury or death of the crop.**
- Apply MEY Canada GlufosSupreme 150 SN Herbicide at 2 L to 3.33 L/ha from the cotyledon to the flowering stage of the crop.

SUSCEPTIBLE WEEDS: MEY Canada GlufosSupreme 150 SN Herbicide has an effect on all weeds and crops except for those crops which are developed to be tolerant to applications of MEY Canada GlufosSupreme 150 SN Herbicide . The following weeds are susceptible to application of MEY Canada GlufosSupreme 150 SN Herbicide . Best control will be obtained when MEY Canada GlufosSupreme 150 SN Herbicide is applied in the recommended leaf stages.

WEED	RECOMMENDED WEED LEAF STAGE								RATE
	1	2	3	4	5	6	7	8	
Cow Cockle-----									1.33 L/ha
Green Foxtail -----									Maximum 3 tillers

WEED	RECOMMENDED WEED LEAF STAGE				RATE			
	Barnyard Grass				1	2	3	4

2026-0660
2026-04-02

5 6 8 2 L/ha
7

-|

Lady's-thumb-----|

2026-0660
2026-04-02

Lamb's-quarters	
Russian Thistle	Up to 8 cm height
Smartweed	
Stinkweed	
Volunteer Flax	Up to 6 cm height
Wild Mustard	-----

WEED	RECOMMENDED WEED LEAF STAGE								RATE
	1	2	3	4	5	6	7	8	
Canada Thistle ¹	Up to 10 cm height								2.67 L/ha
Common Chickweed	----- Leaf Pairs								
Hemp-nettle	----- Leaf Pairs								
Kochia	Up to 8 cm height								
Perennial Sow Thistle	-----								
Quackgrass ¹	-----								
Redroot Pigweed	-----								
Round-leaved Mallow	-----								
Scentless Chamomile	Up to 10 cm height								
Shepherd's-purse	-----								
Volunteer Barley ²	----- Maximum 2 tillers								
Volunteer Wheat	----- Maximum 2 tillers								
Wild Buckwheat	-----								

WEED	RECOMMENDED WEED LEAF STAGE								RATE
	1	2	3	4	5	6	7	8	
Cleavers	-----	Whorls							3.33 L/ha
Dandelion	1-15 cm rosette								
Flixweed	Up to 10 cm height								
Hemp-nettle	-----				Leaf Pairs				
Quackgrass ³	-----								
Stork's-bill	-----								
Wild Oats	-----				Maximum 2 tillers				

WEED	RECOMMENDED WEED LEAF STAGE								RATE
	1	2	3	4	5	6	7	8	
Heavy Populations									
Canada Thistle ¹	Up to 10 cm height								3.33 L/ha
Quackgrass ¹	-----								
Volunteer Barley ²	-----				Maximum 2 tillers				
Volunteer Wheat	-----				Maximum 2 tillers				
Wild Buckwheat	-----								

WEED	RECOMMENDED WEED LEAF STAGE								RATE
	1	2	3	4	5	6	7	8	
Canada Thistle ⁴	Up to 10 cm height								4 L/ha
Quackgrass ⁶	-----								

¹ Top Growth Suppression Only

² Suppression Only

³ Improved Top Growth Control

⁴ Better Top Growth Suppression

⁵ Improved Control

⁶ Season Long Control for Heavy Populations

SECOND APPLICATION - CANOLA

~A second application of MEY Canada GlufosSupreme 150 SN Herbicide can be made to Canola treated initially with up to 4.0 L/ha if new weed germination or growth is present.

~A first application of up to 4 L/ha may be followed by a second application of up to 3.33 L/ha,

OR

A first application of up to 3.33 L/ha may be followed by a second application of up to 4 L/ha.

~Do not apply more than a total of 7.33 L/ha per year.

- ~Apply when the new weed growth is in the correct leaf stage and up until, but prior to, the bolting stage of Canola.

SECOND APPLICATION - SOYBEAN

- ~A second application of MEY Canada GlufosSupreme 150 SN Herbicide can be made to Soybeans treated initially with up to 3.33 L/ha if new weed germination or growth is present. Apply when the new weed growth is in the correct leaf stage and up to the maximum leaf stage of the crop.
- ~Do not apply beyond the flowering stage of Soybeans.
- ~Do not apply more than a total of 6.67 L/ha in one season.

Use of Corn separator rows in hybrid Canola seed production

- ~MEY Canada GlufosSupreme 150 SN Herbicide is registered for use on glufosinate ammonium tolerant Corn varieties or hybrids when used as separator rows in hybrid Canola seed production.
- ~Corn should be seeded at 20,000 – 30,000 plants/ha.
- ~Row spacing for glufosinate ammonium tolerant Corn should be 18-23 cm.
- ~**Note:** Corn must be destroyed and not used for food or feed.

Section 8: Tank Mixes

- ~Do not tank mix MEY Canada GlufosSupreme 150 SN Herbicide with herbicides, fertilizers or chemical additives unless recommended on this label.
- ~Consult the label of the tank mix partner for further directions for use, restrictions and precautions. Tank mix partners may have additional restrictions for leaf staging.
- ~For enhanced activity MEY Canada GlufosSupreme 150 SN Herbicide may be tank mixed with the following products:

CANOLA

TANK MIX PRODUCT	RATE	DIRECTIONS
Facet L + Merge®	0.28 L/ha + 0.5 – 1.0 L/ha	For enhanced and more consistent control of cleavers, apply MEY Canada GlufosSupreme 150 SN Herbicide at a rate of 3.33 L/ha plus Facet L at 0.28 L/ha. Apply in 100 L/ha of water with Merge Adjuvant at a rate of 0.5 to 1 L/ha. Application should be made from the 2 to 6 true leaf stage of the canola crop.

Section 9: Primocane Control in Raspberry

- ~ Apply MEY Canada GlufosSupreme 150 SN Herbicide in a minimum of 330 L/ha of water.

2026-0660
2026-04-02

Use a combination of spray pressures and nozzles, which will provide small droplet sizes and thorough coverage.

CROP	RECOMMENDATIONS	RATE
RASPBERRY (established Raspberries only)	MEY Canada Glufosupreme 150 SN Herbicide is recommended for the control of primocanes in established Raspberries. Apply when shoots are	BROADCAST RATE 6.67 L/ha BANDED RATE Adjust the Broadcast application rate

	about 10-20 cm in height. DO NOT apply to immature or weak plantings.	per hectare proportional to the width of the band applied. Use the formula to calculate the banded rate per hectare:
	FORMULA TO CALCULATE THE BANDED RATE PER HECTARE: Banded rate per hectare = (Band width ÷ Row width) x Broadcast Rate Example: 3.05 metre row spacing and 0.61 metre band width (0.61 M ÷ 3.05 M) x 6.67 L/ha = 1.33 L/ha band rate	

Caution: Avoid contact of the spray with desirable canes, plants and vegetation.

DESICCATION USE

Section 10: Crops, Instructions and Rate

- ~MEY Canada GlufosSupreme 150 SN Herbicide may be applied for desiccation, by ground or air, to Alfalfa (grown for seed).
- ~MEY Canada GlufosSupreme 150 SN Herbicide will also desiccate weeds, which are present in the field at time of application (Wild Buckwheat may not be completely desiccated).
- ~Desiccation of crops and weeds will be best when environmental conditions are favourable (warm temperatures, good moisture conditions, high humidity)

CROP	INSTRUCTIONS	RATE
ALFALFA	Apply MEY Canada GlufosSupreme 150 SN Herbicide	2.7 L/ha
(grown for seed)	at 50-75% pod turn (brown) stage. Do NOT apply more than once per year.	

Note: Do NOT tank mix MEY Canada GlufosSupreme 150 SN Herbicide with pesticides, fertilizers or chemical additives unless recommended on this label.

GENERAL INSTRUCTIONS

Section 11: Herbicide Resistance Management Recommendations

For resistance management, MEY Canada GlufosSupreme 150 SN Herbicide is a Group 10 herbicide. Any weed population may contain or develop plants naturally resistant to MEY Canada GlufosSupreme 150 SN Herbicide and other Group 10 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

2026-0660
2026-04-02

To delay herbicide resistance:

~Where possible, rotate the use of MEY Canada GlufosSupreme 150 SN Herbicide or

other Group 10 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.

- ✓Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- ✓Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- ✓Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- ✓Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- ✓Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- ✓For further information or to report suspected resistance, contact Dell Tech Laboratories Ltd. at 1-519-858-5024

Section 12: Application Instructions and Precautions

APPLICATION INSTRUCTIONS:

- ✓The speed of action of MEY Canada GlufosSupreme 150 SN Herbicide is influenced by environmental factors. At cool temperatures (below 10° C), poor moisture and low humidity, the speed of action may be reduced. Generally, visual symptoms appear 2-4 days after application.
- ✓For best results, apply to emerged, young, actively growing weeds. Weeds that emerge after application will not be controlled.
- ✓MEY Canada GlufosSupreme 150 SN Herbicide will have an effect on weeds that are larger than the recommended leaf stage; however speed of activity and control may be reduced.
- ✓MEY Canada GlufosSupreme 150 SN Herbicide works primarily as a contact herbicide. Thorough coverage of the weeds to be controlled, or the crop to be desiccated, is essential.
- ✓When a rate range is given the higher rate should be utilized:
 1. when the weed or crop growth is dense;
 2. when the weeds are large and/or mature - i.e., advanced leaf stages and plant height;
 3. when the environmental conditions are cool and dry.
- ✓Refer to Section 7 for weeds controlled.

Ground Application Instructions:

- Ensure that all circuits (pipes, booms etc.) have the correct MEY Canada Glufosupreme 150 SN Herbicide /water concentration before application is started.
- Apply MEY Canada Glufosupreme 150 SN Herbicide in a minimum of 110 L/ha of water, at a pressure of 275 kPa and at a ground speed of 6-8 kph. If check valves are used, apply at 310 kPa. The use of 80° or 110° flat fan nozzles is highly recommended for optimum spray coverage and canopy penetration.
- Application of the spray at a 45° angle forward will result in better spray coverage. Refer to Sections 7, 8 (Post-emergent) and 10 (Desiccation) for the correct rate and timing of application.
- For desiccation uses only**, where the crop canopy is dense or weed growth is heavy, achieve better spray coverage with higher spray volumes. Under these conditions, apply in a minimum of 170-220 L/ha of water. Refer to Aerial Application for appropriate aerial water volumes.

Aerial Application Instructions:

- Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.
- Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.
- Exercise extreme caution during the aerial application of any insecticide, herbicide or fungicide. Drift of pesticides is not always visible with the human eye. Small droplets may drift into sensitive areas without obvious signs of danger. Follow these directions precisely.
- When applying MEY Canada Glufosupreme 150 SN Herbicide by aircraft, uniform spray coverage is essential. Applicators are required to use the correct combination of spray nozzle tips, nozzle placement and spray pressures which will provide a coarse droplet size distribution with a volume mean diameter greater than 350 microns. Do not use raindrop nozzles. To avoid streaked, uneven or overlapped application, use appropriate marking devices.
- Post-emergent Use in Glufosinate Ammonium Tolerant Canola and Soybean Water Volume:** Apply MEY Canada Glufosupreme 150 SN Herbicide in a minimum of 55 L/ha of water only.
- Desiccation Use Only Water Volume Range:** If the crop canopy is dense or weed growth is heavy, achieve better spray coverage with higher spray volumes. Under these conditions, if using aerial application, apply MEY Canada Glufosupreme 150 SN Herbicide in a minimum of 33-55 L/ha of water.

APPLICATION PRECAUTIONS

- ⚠ Weed control may be reduced when heavy dew, fog or mist is present at the time of application.
- ⚠ Uniform, thorough spray coverage is important to achieve consistent weed control.

Crop injury may result if MEY Canada GlufosSupreme 150 SN Herbicide is applied to a crop stressed by severe weather conditions, frost, drought, water-saturated soil, low fertility, disease or insect damage.

MEY Canada GlufosSupreme 150 SN Herbicide is a non-selective herbicide and has the potential to desiccate, defoliate or kill all green plants. Avoid contact with other desirable plants or crops by direct application or from spray drifts as severe damage may occur.

Application of MEY Canada GlufosSupreme 150 SN Herbicide to non-tolerant Canola or Soybean varieties, hybrids or other non-tolerant crops will result in severe crop injury or death of the crops.

To assure crop safety and optimal herbicide performance, only use MEY Canada GlufosSupreme 150 SN Herbicide on glufosinate ammonium tolerant Canola or Soybean grown from certified seed.

Plant-back Intervals:

- 70 days for buckwheat, barley, millet, oats, rye, sorghum, triticale.
- 120 days for all other crops except field corn, canola and soybeans, dry common beans (not grown for seed), alfalfa, carrot, lettuce, onion and potato.

Ground Application Precautions:

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.

Do NOT apply if winds exceed 16 kph when using open boom sprayers for ground application.

Do NOT apply if winds exceed 25 kph when using hooded sprayer for ground application.

Do not use flood jet nozzles, controlled droplet application equipment or air-assisted spray equipment.

Aerial Application Precautions:

Aerial application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 16 km/h at flying height at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) coarse classification. To reduce drift caused by turbulent wingtip vortices, the nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wingspan or rotorspan.

Read and understand the entire label before opening this product. If you have questions, call Dell Tech Laboratories Ltd. at 1-519-858-5024 or obtain technical advice from the distributor or your provincial agricultural representative.

Aerial Application Operator Precautions:

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

- ~The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.
- ~All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Rainfast Period:

If rainfall occurs within 4 hours of application, effectiveness may be reduced.

Spray Buffer Zones:

Spot treatments using hand-held equipment do not require a spray buffer zone.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), and sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) .

Method of application	Crop		Buffer Zones (metres) Required for the Protection of:		
			Freshwater Habitat of Depths:		Terrestrial Habitat
			Less than 1 m	Greater than 1 m	
Field sprayer	Alfalfa, Canola, Corn, Raspberry, Soybean		1	0	1
Aerial	Alfalfa, soybean (fixed wing and rotary)		1	0	20
	Canola	Fixed-wing	1	0	35
		Rotary-wing	1	0	30

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

Section 13: Mixing Instructions

MEY Canada GlufosSupreme 150 SN Herbicide must be applied with properly calibrated, clean equipment.

- ~MEY Canada GlufosSupreme 150 SN Herbicide is specially formulated to mix readily in water.
- ~Prior to adding MEY Canada GlufosSupreme 150 SN Herbicide to the spray tank,

ensure that the spray tank is thoroughly clean (see Section 14, “Sprayer Cleanup”).

1. Fill the tank three-quarters full with clean water.
2. Add the correct amount of MEY Canada GlufosSupreme 150 SN Herbicide .
3. Add the remaining amount of water, begin agitation, and spray out immediately.

Section 14: Sprayer Cleanup

~Before and after using MEY Canada GlufosSupreme 150 SN Herbicide always complete a thorough cleaning of the spray tank, lines, and filter. Spray equipment should be thoroughly rinsed using a strong detergent solution.

Section 15: Livestock Feeding and Preharvest Intervals

LIVESTOCK FEEDING:

Canola:

- ~Grain and meal from treated crop can be fed to livestock.
- ~Do not graze the treated crop or cut for hay; sufficient data are not available to support such use.

Corn (used as separator rows in Hybrid Canola Seed Production):

- ~Do not feed treated crop to livestock; sufficient data are not available to support such use.

Soybean:

- ~Do not graze the treated fields within 20 days of application.

PRE-HARVEST INTERVALS (PHI)

Canola:

- ✓ ~When MEY Canada GlufosSupreme 150 SN Herbicide is tank mixed with Facet L, observe a PHI of 60 days from the date of treatment.

Soybean:

- ~Do not apply within 70 days of harvesting Soybean seed.

Section 16: Notices

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.