

2026-2522
2026-08-07

ENRICH Glufosinate Ammonium 200 SN

GROUP	10	HERBICIDE
-------	----	-----------

**FOR SALE FOR USE ON GLUFOSINATE AMMONIUM TOLERANT FIELD CORN
AND ON GLUFOSINATE AMMONIUM TOLERANT SOYBEANS IN CANADA AND
ON GLUFOSINATE AMMONIUM TOLERANT CANOLA IN EASTERN CANADA AND
BRITISH COLUMBIA (INCLUDING THE PEACE RIVER REGIONS)**

**SOLUTION
COMMERCIAL**

POISON



**WARNING
EYE AND SKIN IRRITANT**

REGISTRATION NUMBER: 36040 PEST CONTROL PRODUCTS ACT

ACTIVE INGREDIENT:
glufosinate-ammonium 200 g/L
NET CONTENTS: 1L - 1,000 L

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

Product Information: +1-514-578-1459

Enrich Agrochemical Company Limited

**9015 Crois Rubens Brossard, QC Canada J4X2H7
1-514-578-1459**

In case of spills, poisoning or fire, telephone emergency response number: +1-514-578-1459

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed. Harmful if absorbed through the skin. Causes eye irritation. DO NOT get in eyes. May cause skin irritation. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist.

DO NOT eat, drink or smoke/vape while using this product. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum/tobacco, smoking/vaping, or using the toilet. Vehicle cabs must be decontaminated regularly.

Protective Clothing and Equipment: Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks, shoes and a respirator with a NIOSH-approved organic-vapour-removing cartridge (with a prefilter) approved for pesticides, or a NIOSH-approved canister approved for pesticides during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair. Gloves are not required during application within a closed cab.

If the product comes in contact with clothing/PPE, remove all contaminated clothing/PPE, wash with soap and water and dress in clean clothing. Wash the outside of gloves with soap and water before removing. Launder contaminated clothing/PPE regularly and separately from other laundry.

Restricted-Entry Interval: DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

If this pest control product is to be used on a commodity that may be exported and you require information regarding Maximum Residue Limits for an importing country, please contact Enrich Agrochemical Company Limited at +1-514-578-1459.

FIRST AID:

If swallowed, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15– 20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Note to Physician: If this product is ingested, endotracheal intubation and gastric lavage should be performed as soon as possible, followed by charcoal and sodium sulfate administration.

ENVIRONMENTAL PRECAUTIONS:

Toxic to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under Directions for Use.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

STORAGE: Store this product away from food or feed.

CANNOT be stored below freezing. If stored for 1 year or longer, shake well before using. Store the tightly closed container away from feeds, seeds, fertilizers, plants and foodstuffs. **DO NOT** store in and around the home. Keep in original container during storage.

DISPOSAL:

Recyclable Container: DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

Returnable Container: DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable Container: For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

Disposal of Unused, Unwanted Product: For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for cleanup of spills.

Disposal of Unused Spray Solution: If any spray solution remains in the tank after spraying is finished, it should be sprayed on the perimeter of the area just sprayed, away from water supplies, ditches, and irrigation canals.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

ENRICH Glufosinate Ammonium 200 SN

GROUP	10	HERBICIDE
-------	-----------	-----------

**FOR SALE FOR USE ON GLUFOSINATE AMMONIUM TOLERANT FIELD CORN
AND ON GLUFOSINATE AMMONIUM TOLERANT SOYBEANS IN CANADA AND
ON GLUFOSINATE AMMONIUM TOLERANT CANOLA IN EASTERN CANADA AND
BRITISH COLUMBIA (INCLUDING THE PEACE RIVER REGIONS)**

**SOLUTION
COMMERCIAL**



EYE AND SKIN IRRITANT

REGISTRATION NUMBER: 36040 PEST CONTROL PRODUCTS ACT

**ACTIVE INGREDIENT: glufosinate ammonium 200 g/L
NET CONTENTS: 1L – 1,000L**

READ THE LABEL AND THIS BOOKLET BEFORE USING

Product Information: +1-514-578-1459

Enrich Agrochemical Company Limited

**9015 Crois Rubens Brossard, QC Canada J4X2H7
1-514-578-1459**

In case of spills, poisoning or fire, telephone emergency response number: +1-514-578-1459

ENRICH Glufosinate Ammonium 200 SN		<i>Section Number:</i>
<i>Table of Contents:</i>		
GENERAL INFORMATION	The Product	1
SAFETY AND HANDLING	Precautions, Protective Clothing and Equipment, and Restricted-Entry Interval	2
	First Aid and Toxicological Information	3
	Environmental Precaution	4
	Storage	5
	Disposal	6
DIRECTIONS FOR USE	Crops, Weeds, Rates, Timing	7
	Application Instructions and Cautions	8
	Tank Mixes	9
	Mixing Instructions	10
	Sprayer Cleanup	11
	Livestock Feeding and Preharvest Intervals	12
	Resistance Management	13
	Notices	14

GENERAL INFORMATION

Section 1: The Product

ENRICH Glufosinate Ammonium 200 SN provides control of a broad spectrum of grassy and broadleaf weeds in Canola varieties, Corn hybrids and Soybean varieties that are specially developed to be tolerant to glufosinate ammonium.

ENRICH Glufosinate Ammonium 200 SN may also be applied to glufosinate ammonium tolerant inbred Corn lines grown for seed production.

For Canola hybrid seed production, ENRICH Glufosinate Ammonium 200 SN may be used to remove segregating wild type plants within the female population. See Section 7 (Crops, Weeds, Rates, Timing - Canola Hybrid Seed Production).

SAFETY AND HANDLING

Section 2: Precautions, Protective Clothing and Equipment, and, Re-entry Restrictions

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed. Harmful if absorbed through the skin. Causes eye irritation. DO NOT get in eyes. May cause skin irritation. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist.

DO NOT eat, drink or smoke/vape while using this product. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum/tobacco, smoking/vaping, or using the toilet. Vehicle cabs must be decontaminated regularly.

Protective Clothing and Equipment: Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks, shoes and a respirator with a NIOSH-approved organic-vapour-removing cartridge (with a prefilter) approved for pesticides, or a NIOSH-approved canister approved for pesticides during mixing, loading, application, clean-up and repair. In addition, wear protective eyewear (goggles or face shield) during mixing, loading, clean-up and repair. Gloves are not required during application within a closed cab.

If the product comes in contact with clothing/PPE, remove all contaminated clothing/PPE, wash with soap and water and dress in clean clothing. Wash the outside of gloves with soap and water before removing. Launder contaminated clothing/PPE regularly and separately from other laundry.

Restricted-Entry Interval:

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

Section 3: First Aid and Toxicological Information

FIRST AID:

If swallowed, call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing, take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled, move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes, hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Note to Physician: If this product is ingested, endotracheal intubation and gastric lavage should be performed as soon as possible, followed by charcoal and sodium sulfate administration.

Section 4: Environmental Precautions

Toxic to aquatic organisms and non-target terrestrial plants . Observe spray buffer zones specified under Directions for Use.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

Section 5: Storage

Store this product away from food or feed.

CANNOT be stored below freezing.

If stored for 1 year or longer, shake well before using.

Store the tightly closed container away from feeds, seeds, fertilizers, plants and foodstuffs.

DO NOT store in and around the home.

Keep in original container during storage.

Section 6: Disposal

Recyclable Container:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

Returnable Container: DO NOT reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

Refillable Container: For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. DO NOT reuse this container for any other purpose.

Disposal of Unused, Unwanted Product:

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for cleanup of spills.

Disposal of Unused Spray Solution:

If any spray solution remains in the tank after spraying is finished, it should be sprayed on the perimeter of the area just sprayed, away from water supplies, ditches, and irrigation canals.

DIRECTIONS FOR USE

Section 7: Crops, Weeds, Rates, Timing

Best control will be obtained when ENRICH Glufosinate Ammonium 200 SN is applied at the recommended leaf stages of the weeds.

In Canola grown in Eastern Canada and British Columbia, apply ENRICH Glufosinate Ammonium 200 SN from the cotyledon stage up until, but prior to, the bolting stage of Canola.

In Corn grown in Canada, apply ENRICH Glufosinate Ammonium 200 SN from the 1-8 leaf stage of the

Corn plant or 5-6 visible collars (the leaf is counted once the next leaf is visible in the whorl).

In Soybeans grown in Canada, apply ENRICH Glufosinate Ammonium 200 SN from the cotyledon to the flowering stage of the crop.

Crop	Weed	Recommended Time (Leaf Stage)								Rate
		1	2	3	4	5	6	7	8	
CANOLA, CORN, and SOYBEANS tolerant to ENRIC H Glufosina te Ammoniu m 200 SN	Crabgrass-----					x				1.5 L/ha
	Giant Foxtail -----					x				
	Green Foxtail -----					x				
	Proso Millet-----					x				
	Chickweed -----								x	
	Cocklebur -----				x					
	Common Ragweed-----					x				
	Redroot Pigweed -----						x			
	Shepherd's- purse -----						x			

Crop	Weed	Recommended Time (Leaf Stage)								Rate
		1	2	3	4	5	6	7	8	
CANOLA, CORN, and SOYBEANS tolerant to ENRIC H Glufosina te Ammoniu m 200 SN	Barnyard Grass -----					x				2 L/ha
	Bristly Foxtail -----				x					
	Fall Panicum-----					x				
	Old Witchgrass -----					x				
	Wild Oats -----					x				
	Yellow Foxtail-----					x				
	Eastern Black Nightshade -----						x			
	Lady's-thumb-----								x	
	Lamb's-quarters -----								x	
	Perennial Sowthistle -----				x					
	Common Ragweed-----							x		
	Stinkweed -----								x	
	Velvetleaf -----					x				
	Wild Buckwheat-----					x				

Wild Mustard -----x
Wormseed Mustard ----- x

Canada Thistle and Field Bindweed: For season long suppression of Canada Thistle and Field Bindweed, apply ENRICH Glufosinate Ammonium 200 SN at a rate of 2 L/ha when the weeds are in the 1-6 leaf stage.

Quackgrass: For season long suppression of Quackgrass, apply ENRICH Glufosinate Ammonium 200 SN at a rate of 2.5 L/ha plus ammonium sulphate at a rate of 6 L/ha (49% solution) OR 3 kg/ha (99%) when the Quackgrass is in the 1-4 leaf stage.

Voluntary Crop Weeds: ENRICH Glufosinate Ammonium 200 SN will not control volunteer crop weeds that emerge from previous plantings of glufosinate ammonium tolerant crops.

ADDITIVES: For enhanced control of Velvetleaf, add ammonium sulphate to the tank at a rate of 6 L/ha (49% solution) OR 3 kg/ha (99%).

SECOND APPLICATION: A second application of ENRICH Glufosinate Ammonium 200 SN can be made to fields treated initially with up to 2.5 L/ha if new weed germination or growth is present.

Apply when the new weed growth is in the correct leaf stage and up to the maximum leaf stage of the crop.

Application after the 8-leaf stage of Corn must be made with drop nozzles as a directed spray below the whorl of the Corn plant.

Application in Canola must be made prior to the bolting stage of the Canola.

Do not apply beyond the flowering stage of Soybeans.

Do not apply more than a total of 4.5 L/ha in one season.

SPLIT APPLICATION PROGRAM: For season long control of the above listed weeds a split application program with ENRICH Glufosinate Ammonium 200 SN can be utilised. The first application must be a minimum of 2 L/ha and at the proper weed stage. For the second application, the rate of

1.25 L/ha may be utilized provided ENRICH Glufosinate Ammonium 200 SN is applied as soon as the second flush of weeds occurs and before the maximum leaf stage of the crop.

FOR CANOLA HYBRID SEED PRODUCTION - Two applications are required to remove the segregating wild type plants. The first application should occur when the Canola is in the 2-4 leaf stage, the second application when the Canola plants are in the 4-6 leaf stage and the third application, if necessary, may be applied to the Canola up until, but prior to, the time of bolting. All applications of ENRICH Glufosinate Ammonium 200 SN for hybrid seed production should be made at 2.5 L/ha.

Section 8: Application Instructions and Cautions

Application Instructions:

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

For best results, apply to emerged, young, actively growing weeds. Weeds that emerge after application will not be controlled.

ENRICH Glufosinate Ammonium 200 SN will have an effect on weeds that are larger than the recommended leaf stage, however speed of activity and control may be reduced.

Apply ENRICH Glufosinate Ammonium 200 SN in a minimum of 110 L/ha of water, at a pressure of 175-

275 kPa and at a ground speed of 6-10 kph. The use of 80° or 110° flat fan nozzles is highly recommended for optimum spray coverage and canopy penetration.

Application of the spray at a 45° angle forward will result in better spray coverage. Follow directions elsewhere on the label for the correct rate and timing of application.

The speed of action of ENRICH Glufosinate Ammonium 200 SN is influenced by environmental factors.

At cool temperatures (below 10°C), poor moisture and low humidity, speed of action may be reduced. Generally, visual symptoms appear 2-4 days after application.

IF RAINFALL OCCURS WITHIN 4 HOURS OF APPLICATION, EFFECTIVENESS MAY BE REDUCED.

Weed control may be reduced when heavy dew, fog or mist are present at the time of application.

ENRICH Glufosinate Ammonium 200 SN works primarily as a contact herbicide. Thorough coverage of the weeds to be controlled is essential.

Application Cautions:

Field sprayer application: DO NOT apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. DO NOT apply with sprays finer than the American Society of Agricultural and Biological Engineers (ASABE) S572 (572.1 to 572.3) Medium classification. Boom height must be 60 cm or less above the crop or ground.

DO NOT apply by air.

Do not use flood jet nozzles, controlled droplet application equipment or air-assisted spray equipment. Uniform, thorough spray coverage is important to achieve consistent weed control.

DO NOT apply if winds exceed 16 kph when using open boom sprayers. DO NOT apply if winds exceed 25 kph when using hooded sprayers.

Avoid contact with desirable plants or crops either from direct application or from spray drift as severe damage may occur.

Plant back Intervals:

- 70 days for Buckwheat, Barley, Millet, Oats, Rye, Sorghum, Triticale and Wheat
- 120 days for all other crops, except Field Corn, Canola and Soybeans
- No plant back interval is required for Field Corn, Canola or Soybeans.

Application of ENRICH Glufosinate Ammonium 200 SN to non-tolerant crops will result in severe crop injury or death of the crop.

SUSCEPTIBLE WEEDS: ENRICH Glufosinate Ammonium 200 SN has an effect on all weeds and crops except for those crops which are developed to be tolerant to applications of ENRICH Glufosinate Ammonium 200 SN, including when they become voluntary weeds.

Spray Buffer zones:

A spray buffer zone is NOT required for low-clearance hooded or shielded sprayers that prevent spray contact with crop, fruit or foliage.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

Method of Application	Crop	Spray Buffer Zones (meters) Required for the Protection of:		
		Freshwater Habitat of Depths:		Terrestrial Habitat
		Less than 1 m	Greater than 1 m	
Field Sprayer	Corn, Canola, Soybean	1	0	1

When tank mixes are permitted, consult the labels of the tank mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASABE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product for conventional application equipment can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Drift Mitigation portion of the Canada.ca website.

Section 9: Tank Mixes

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Enrich Agrochemical Company Limited at +1- 514-578-1459 for information before applying any tank mix that is not specifically recommended on this label.

For control of weeds present and/or control of later flushes of annual broadleaf weeds and/or annual grasses (see Tank Mix Directions), ENRICH Glufosinate Ammonium 200 SN may be tank mixed with the following herbicides when used in Eastern Canada and British Columbia:

CROP	TANK MIX PRODUCT	RATE	DIRECTIONS
CANOLA	Centurion® Emulsifiable Concentrate Post Emergence Herbicide or Select® Emulsifiable Concentrate Post Emergence Herbicide + Amigo Adjuvant at 0.5% v/v	63 mL/ha	For control of Volunteer Barley and improved control of Wild Oats, apply ENRICH Glufosinate Ammonium 200 SN at a rate of 2.0 – 2.5 L/ha plus Centurion Emulsifiable Concentrate Post Emergence Herbicide or Select Emulsifiable Concentrate Post Emergence Herbicide at 63 mL/ha with adjuvant as recommended on the Centurion Emulsifiable Concentrate Post Emergence Herbicide or Select Emulsifiable Concentrate Post Emergence Herbicide label. Apply when the weeds are in the 1-4 leaf stage with a maximum of 2 tillers.
CORN	AAtrex Liquid 480 Herbicide*	1.75 – 2.33 L/ha	Mix with the appropriate rate of ENRICH Glufosinate Ammonium 200 SN for the targeted weed species. Adjust atrazine rate to the level of residual activity desired. This tank mix will not control later flushes of triazine resistant weeds.
CORN	Distinct® Herbicide	285 g/ha	Mix with the appropriate rate of ENRICH Glufosinate Ammonium 200 Herbicide for the targeted weed species. This tank mix will provide enhanced control of labeled annual broadleaf weeds.
CORN	Marksman® Herbicide	2.5 - 3.7 L/ha	Mix with the appropriate rate of ENRICH Glufosinate Ammonium 200 SN for the targeted weed species. Adjust Marksman Herbicide rate to the level of residual activity desired. This tank mix will provide residual control of annual grasses and annual broadleaf weeds.
SOYBEAN	Pursuit®Herbicide	312 mL/ha	Mix with the appropriate rate of ENRICH Glufosinate Ammonium 200 SN for the targeted weed species. This tank mix will provide residual control of annual grasses and annual broadleaf weeds.

SOYBEA N	Basagran® Forte Herbicide	1.75 L/ha	Mix with the appropriate rate of ENRICH Glufosinate Ammonium 200 SN for the targeted weed species. This tank mix will provide enhanced control of annual broadleaf weeds.
----------	------------------------------	-----------	---

*for atrazine products with different concentration, adjust the rate accordingly.

Note: Do not tank mix ENRICH Glufosinate Ammonium 200 SN with fertilizers or any other chemical additives unless recommended on this label.

Section 10: Mixing Instructions

ENRICH Glufosinate Ammonium 200 SN must be applied with properly calibrated, clean equipment. ENRICH Glufosinate Ammonium 200 SN is specially formulated to mix readily in water.

Prior to adding ENRICH Glufosinate Ammonium 200 SN to the spray tank, ensure that the spray tank is thoroughly clean (see Section 12 “SPRAYER CLEANUP”).

1. Fill tank to three-quarters full with clean water prior to adding ENRICH Glufosinate Ammonium 200 SN.
2. Add the correct amount of ENRICH Glufosinate Ammonium 200 SN.
3. Add the remaining amount of water, begin agitation, and spray out immediately.

Note: With the exception of use with CENTURION Emulsifiable Concentrate Post-Emergence Herbicide or SELECT Emulsifiable Concentrate Post-Emergence Herbicide, when tank mixing ENRICH Glufosinate Ammonium 200 SN always add the tank mix partner to the tank first, ensure that it is thoroughly mixed, THEN add ENRICH Glufosinate Ammonium 200 SN. When using ammonium sulphate, ALWAYS add it to the tank first, prior to the addition of the tank mix partner and ENRICH Glufosinate Ammonium 200 SN. Refer to the following specific instructions for mixing with CENTURION Emulsifiable Concentrate Post-Emergence Herbicide or SELECT Emulsifiable Concentrate Post-Emergence Herbicide.

ENRICH Glufosinate Ammonium 200 SN and CENTURION Emulsifiable Concentrate Post-Emergence Herbicide or SELECT Emulsifiable Concentrate Post-Emergence Herbicide: Glufosinate Ammonium

Tolerant Canola only

When tank mixing ENRICH Glufosinate Ammonium 200 SN and CENTURION Emulsifiable Concentrate Post-Emergence Herbicide or SELECT Emulsifiable Concentrate Post-Emergence Herbicide, always add AMIGO adjuvant to the tank first, **then** add ENRICH Glufosinate Ammonium 200 SN to the tank followed by the CENTURION Emulsifiable Concentrate Post-Emergence Herbicide or SELECT Emulsifiable Concentrate Post-Emergence Herbicide.

The addition of an anti-foaming agent may reduce foaming, especially when using soft water.

1. Thoroughly clean the sprayer by flushing the system with water containing detergent (see Section 12, “Sprayer Cleanup”).
2. Fill clean spray tank half full with clean water. Start agitation system.
3. Add the correct amount of AMIGO. Continue to agitate until AMIGO is thoroughly mixed.
4. STOP agitation. Add the required amount of ENRICH Glufosinate Ammonium 200 SN to the spray tank. Start agitation system.
5. Add the correct amount of CENTURION Emulsifiable Concentrate Post-Emergence Herbicide or SELECT Emulsifiable Concentrate Post-Emergence Herbicide along with the remaining amount of water necessary to fill the spray tank.
6. Continue to agitate or run the by-pass system and spray out immediately.
7. After any break in the spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to remix the spray materials. Do not allow the mixture to sit overnight.
8. If an oil film starts to build up in the tank, drain it and clean tank with strong detergent solution.
9. Immediately after use, thoroughly clean the sprayer by flushing the system with water containing detergent (see Section 12, “Sprayer Cleanup”).

Note: The addition of an anti-foaming agent may reduce foaming, especially when using soft water.

Note: Ensure that all circuits (pipes, booms, etc.) have the correct ENRICH Glufosinate Ammonium 200 SN/water concentration before application is started.

Section 11: Sprayer Cleanup

Before and after using ENRICH Glufosinate Ammonium 200 SN always complete a thorough cleaning of the spray tank, lines and filter. Spray equipment should be thoroughly rinsed using a strong detergent solution.

Section 12: Livestock Feeding and Preharvest Intervals (PHIs)

CANOLA:

Grain and meal from treated Canola can be fed to livestock.

DO NOT graze or feed other portions of the treated Canola to livestock.

When ENRICH Glufosinate Ammonium 200 SN is tank mixed with CENTURION Emulsifiable Concentrate Post-Emergence Herbicide or SELECT Emulsifiable Concentrate Post-Emergence Herbicide, observe a PHI of 60 days from the date of treatment (or last treatment when a second application has been made).

CORN:

DO NOT apply within 86 days of harvesting Corn grain.

DO NOT graze the treated fields within 20 days of application.

SOYBEAN:

DO NOT apply within 70 days of harvesting Soybean seed.

DO NOT graze the treated fields within 20 days of application.

Section 13: Resistance Management Recommendations

For resistance management, ENRICH Glufosinate Ammonium 200 SN is a Group 10 herbicide. Any weed population may contain or develop plants naturally resistant to ENRICH Glufosinate Ammonium 200 SN and other Group 10 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

Where possible, rotate the use of ENRICH Glufosinate Ammonium 200 SN or other Group 10 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.

Use tank mixtures with herbicides from a different group when such use is permitted.

To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.

Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.

Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible, by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.

Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.

For further information or to report suspected resistance, contact Enrich Agrochemical Company Limited at +1-514-578- 1459.

Section 14: Notices**NOTICE TO USER:**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.