

2026-2286
2026-07-23

GROUP	2	HERBICIDE
-------	---	-----------

Rainbow Imazethapyr 240 SL

COMMERCIAL (AGRICULTURAL)

**SOLUTION
HERBICIDE**

ACTIVE INGREDIENT: Imazethapyr, present as ammonium salt - 240 g/L

REGISTRATION NO. 36063 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND ATTACHED BROCHURE BEFORE USING

KEEP OUT OF REACH OF CHILDREN

CAUTION- EYE AND SKIN IRRITANT

**For Chemical Emergency:
Spill, Leak, Fire, Exposure, or Accident, Call CHEMTREC: 1-800-424-9300**

NET CONTENTS: 1 Litre – Bulk

Rainbow CropSciences Inc.
535 E Plainfield Road, Suite E
Willowbrook, IL 60527
(630) 828-5088

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

DO NOT get in eyes. May irritate eyes.

May cause skin irritation. Avoid contact with skin and clothing.

DO NOT contaminate food or feed products.

Avoid breathing vapour or spray mist. Use with adequate ventilation.

DO NOT eat, drink or smoke/vape while using this product. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum/tobacco, smoking/vaping, using the toilet or handling mobile devices. Vehicle cabs MUST be decontaminated regularly.

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, protective eyewear (goggles or face shield), socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab.

If the product comes in contact with clothing/PPE, remove all contaminated clothing/PPE, wash the outside of gloves with soap and water before removing, wash contaminated skin with soap and water, and then dress in clean clothing. Launder contaminated clothing/PPE regularly and separately from other laundry. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Heavily contaminated or drenched clothing and other absorbent materials contaminated with this product should be discarded. DO NOT reuse them.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

DO NOT apply by air.

ENVIRONMENTAL PRECAUTIONS

TOXIC to non-target terrestrial and aquatic plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

STORAGE

1. Store the leftover product in original tightly closed container.
2. Keep product from freezing. DO NOT store below 0°C. If the product is exposed to temperatures below 0°C during shipment or storage, make sure the product has thawed completely, and shake the container vigorously.
3. DO NOT ship or store the product near food, feed, seed and fertilizers. Store this product away from food or feed.

4. Store the product in cool, dry, locked, well-ventilated areas without floor drain.

DISPOSAL

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for cleanup of spills.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. DO NOT induce vomiting unless told to do so by a poison control centre or doctor. DO NOT give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

GROUP

2

HERBICIDE

Rainbow Imazethapyr 240 SL**COMMERCIAL (AGRICULTURAL)****SOLUTION
HERBICIDE****ACTIVE INGREDIENT:** Imazethapyr, present as ammonium salt - 240 g/L**REGISTRATION NO. 36063 PEST CONTROL PRODUCTS ACT****READ THE LABEL AND ATTACHED BROCHURE BEFORE USING****KEEP OUT OF REACH OF CHILDREN****CAUTION- EYE AND SKIN IRRITANT**

For Chemical Emergency:
Spill, Leak, Fire, Exposure, or Accident, Call CHEMTREC: 1-800-424-9300

NET CONTENTS: 1 Litre – Bulk

Rainbow CropSciences Inc.
535 E Plainfield Road, Suite E
Willowbrook, IL 60527
(630) 828-5088

Booklet

PRECAUTIONS**KEEP OUT OF REACH OF CHILDREN.**

DO NOT get in eyes. May irritate eyes.

May cause skin irritation. Avoid contact with skin and clothing.

DO NOT contaminate food or feed products.

Avoid breathing vapour or spray mist. Use with adequate ventilation.

DO NOT eat, drink or smoke/vape while using this product. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum/tobacco, smoking/vaping, using the toilet or handling mobile devices. Vehicle cabs MUST be decontaminated regularly.

Wear a long-sleeved shirt, long pants, chemical-resistant gloves, protective eyewear (goggles or face shield), socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab.

If the product comes in contact with clothing/PPE, remove all contaminated clothing/PPE, wash the outside of gloves with soap and water before removing, wash contaminated skin with soap and water, and then dress in clean clothing. Launder contaminated clothing/PPE regularly and separately from other laundry. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Heavily contaminated or drenched clothing and other absorbent materials contaminated with this product should be discarded. DO NOT reuse them.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

DO NOT apply by air.

ENVIRONMENTAL PRECAUTIONS

TOXIC to non-target terrestrial and aquatic plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

STORAGE

1. Store the leftover product in original tightly closed container.
2. Keep product from freezing. DO NOT store below 0°C. If the product is exposed to temperatures below 0°C during shipment or storage, make sure the product has thawed completely, and shake the container vigorously.
3. DO NOT ship or store the product near food, feed, seed and fertilizers. Store this product away from food or feed.
4. Store the product in cool, dry, locked, well-ventilated areas without floor drain.

Booklet

DISPOSAL

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for cleanup of spills.

Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. DO NOT induce vomiting unless told to do so by a poison control centre or doctor. DO NOT give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

DIRECTIONS FOR USE**GENERAL INFORMATION**

Rainbow Imazethapyr 240 SL is a selective herbicide that can be applied as an early pre-plant, pre-plant incorporated, pre-emergent or post-emergent treatment in various crops. The application method depends upon the crop, anticipated weed spectrum and the preference of the applicator. With early pre-plant and

Booklet

pre-emergent treatments, susceptible weeds emerge, are present as stunted plants and then die. When Rainbow Imazethapyr 240 SL is applied post-emergence, absorption may occur through both the roots and foliage. Susceptible weeds stop growing and eventually die.

Use Rainbow Imazethapyr 240 SL at 312 mL (75 g active) – 420 mL (100 g active) per hectare. For post-emergent application to glyphosate tolerant soybeans, 210 mL (50 g active) – 312 mL (75 g active) per hectare may be applied in tank-mix with glyphosate. See appropriate rate tables for specific application rates for each crop.

APPLICATION INSTRUCTIONS

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

Field sprayer application: DO NOT apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. DO NOT apply with sprays finer than the American Society of Agricultural and Biological Engineers (ASABE) S572 (572.1 to 572.3) Medium classification. Boom height must be 60 cm or less above the crop or ground.

DO NOT apply by air.

SPRAY BUFFER ZONES

A spray buffer zone is NOT required for:

- uses with hand-held application equipment permitted on this label,
- low-clearance hooded or shielded sprayers that prevent spray contact with crop, fruit or foliage

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of Application	Crop/Site	Spray Buffer Zones (metres) Required for the Protection of:				
		Freshwater Habitat of Depths:		Estuarine/Marine Habitat of Depths:		Terrestrial Habitat
		Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer	All crops	1	1	0	0	1

When tank mixes are permitted, consult the labels of the tank mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASABE) category indicated on the labels for those tank mix partners.

Booklet

The spray buffer zones for this product for conventional application equipment can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Drift Mitigation portion of the Canada.ca website.

TANK MIXING

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. DO NOT tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Rainbow CropSciences Inc. at (630) 828-5088 for information before applying any tank mix that is not specifically recommended on this label.

FOR USE IN THE PRAIRIE PROVINCES (i.e., Manitoba, Saskatchewan, Alberta) & INTERIOR OF BRITISH COLUMBIA (INCLUDING THE PEACE RIVER REGION) ONLY

REGISTERED CROPS

Use as an early post-emergence in the following crops.

Alfalfa (newly seeded purestand for forage or seed production)
 Alfalfa (established for seed production)
 Chickling Vetch (seed production only)
 Dry Beans (pinto, pink, red)
 Field Peas
 Soybeans (Manitoba only)

REGISTERED SOIL ZONES

Rainbow Imazethapyr 240 SL is only registered in the black and grey wooded soil zones of the Prairie Provinces. Rainbow Imazethapyr 240 SL is not registered for use in the brown and dark brown soil zones with the exception of dry beans and alfalfa (newly seeded purestand for forage or seed production) under irrigation.

CROP USE RATES

Crop	Use Rate (mL/ha)	Water Volume (L/ha)
Alfalfa (newly seeded purestand for forage or seed production)	210	100-400
Alfalfa (established for seed production)	210	100-400
Chickling vetch (seed production only)	210	100-400
Dry beans (pinto, pink, red)	210	100-400
Field peas	210	100-400
Soybeans (Manitoba only)	210	100-400

For all applications, a non-ionic surfactant containing at least 80% active ingredient (such as Agral® 90 Non-Ionic Wetting & Spreading Agent Liquid, IPCO Ag-surf® Original) at the rate of 0.25% (v/v) must be

Booklet

added to the spray solution (e.g., 2.5 L of surfactant per 1000 L of spray solution).

See crop specific section for additional details.

WEEDS CONTROLLED**EARLY POST- EMERGENCE WEED CONTROL (up to and including the 4 leaf stage)**

Weeds	Application Timing	
	Field Peas	Newly seeded pure stand alfalfa for forage or seed production
Broadleaf weeds		
Common groundsel	-	PC
Chickweed*	C	-
Cleavers*	C	-
Green smartweed	-	C
Hemp-nettle	C	-
Redroot pigweed	C	C
Shepherd's purse*	C	PC
Smartweed	C	-
Stinkweed*	C	C
Volunteer canola	C	C
Wild buckwheat	S	-
Wild mustard*	C	C
Grasses		
Green foxtail	C	PC
Wild oats	S	-

C = Control PC = Partial Control and Reduced Competition S = Suppression

* Certain populations of these weed are known to be resistant to group 2. If you suspect this potential on your farm, follow recommendations in Resistance-Management Recommendations section.

CROP SPECIFIC RECOMMENDATIONS**FIELD PEAS**

Timing	Early post-emergence
Soil Zones	Rainbow Imazethapyr 240 SL is registered for use in black and grey wooded soils ONLY.
Rate	210 mL/ha + adjuvant
Weeds Controlled	See above for a complete list of weeds controlled.

Booklet

Remarks	Rainbow Imazethapyr 240 SL, by early post-emergence application up to the 4-true leaf stage of the weeds, will control weeds listed in the Weeds Controlled section. Field peas up until the sixth (6th) trifoliate leaf stage. An interval of 60 days must follow the Rainbow Imazethapyr 240 SL application before field peas are harvested. Field peas treated with Rainbow Imazethapyr 240 SL may be fed to livestock 30 days after application.
---------	--

ALFALFA (NEWLY SEEDED PURESTAND FOR FORAGE OR SEED PRODUCTION)

Timing	Early post-emergence
Soil Zones	Black, grey wooded and irrigated brown soil zones.
Rate	210 mL/ha + adjuvant
Weeds Controlled	See above for a complete list of weeds controlled.
Remarks	Rainbow Imazethapyr 240 SL can be applied as an early post-emergence treatment in newly seeded purestand alfalfa in the year of establishment. Rainbow Imazethapyr 240 SL may be applied when the crop has developed at least one (1) fully expanded trifoliate leaf and up to and including the 4 leaf stage of susceptible weeds. Rainbow Imazethapyr 240 SL is intended for use on purestand alfalfa in the year of establishment, on stands that will remain for 3 or more years. Apply only in the year of establishment and only after the crop has one fully developed trifoliate leaf. ONLY ONE (1) application of Rainbow Imazethapyr 240 SL may be made during the life of the alfalfa stand. DO NOT graze or harvest for forage until 14 days after treatment. Mature seed from treated plants should not be used for human consumption.

ALFALFA (ESTABLISHED FOR SEED PRODUCTION PURPOSES)

Timing	Early post-emergence
Soil Zones	Black and grey wooded soil zones
Rate	210 mL/ha + adjuvant
Weeds Controlled	Stinkweed, wild mustard, volunteer canola, redroot pigweed and suppression of green foxtail
Remarks	One application per year; post-emergent; ground application; apply before alfalfa reaches 30 cm. DO NOT use Rainbow Imazethapyr 240 SL in the last year of the alfalfa stand.

DRY BEANS (PINTO, PINK, RED)

Timing	Early post-emergence
Soil Zones	Black, grey wooded and irrigated brown soil zones only
Rate	210 mL/ha + adjuvant
Weeds Controlled	Hairy nightshade
Remarks	One application per year; ground sprayer; apply in 100 - 400 L water/ha post-emergence; apply up to the 2nd trifoliate leaf stage of the dry bean; up to 6 leaf stage of hairy nightshade; PHI of 75 days.

CHICKLING VETCH / GRASS PEA (FOR SEED PRODUCTION ONLY)

Timing	Early post-emergence
Soil Zones	Black and grey wooded soil zones
Rate	210 mL/ha + adjuvant
Weeds Controlled	Labelled weeds
Remarks	One application per year; post-emergent; ground application; PHI 60 days; apply at the 5-7 leaf stage of chickling vetch/grass pea.

SOYBEANS (MANITOBA ONLY)

Timing	Early post-emergence
Soil Zones	Black and grey wooded soil zones
Rate	210 mL/ha + adjuvant
Weeds Controlled	Annual grass and broadleaf weeds
Remarks	Apply as post-emergent application once per season at 1 to 3 leaves of soybean growth; ground application only; PHI 85 days.

FOLLOW CROPPING

There is the possibility of residual soil activity from Rainbow Imazethapyr 240 SL the year following application. Research studies have shown the following crops can be safely grown in black and grey wooded soil zones the year following a Rainbow Imazethapyr 240 SL application.

0 - 12 months after application In case of crop failure, the following crops can be planted in the same season:	1 year after application The following crops can be planted in the following season:
field peas Clearfield® Canola¹	spring barley spring wheat lentils alfalfa field peas Clearfield Canola¹
Conduct a field bioassay (a test strip grown to maturity) the year BEFORE growing any other crop than those listed above.	

WARNINGS

- (1) DO NOT APPLY Rainbow Imazethapyr 240 SL on brown & dark brown soil zones unless under irrigation as the follow crop label ONLY pertains to the black and grey wooded soil zones. Use in non-irrigated brown/dark brown soil zones could result in rotational crop injury.
- (2) DO NOT over-apply Rainbow Imazethapyr 240 SL. Over application may result in injury particularly if the crop is under stress.
- (3) DO NOT let spray drift contaminate crops other than those being sprayed.
- (4) DO NOT apply 2 applications of Rainbow Imazethapyr 240 SL within the same year. ONLY ONE (1) application of Rainbow Imazethapyr 240 SL may be made during the season.
- (5) In cases of crop failure, only field peas or **Clearfield Canola¹** (e.g. canola varieties with the **Clearfield** trait) may be re-planted in the year of application.
- (6) **DO NOT APPLY BY AERIAL APPLICATION - GROUND APPLICATION ONLY.**

NOT FOR USE IN PRAIRE PROVINCES

REGISTERED CROPS

Use as an early pre-plant, pre-plant incorporated, pre-emergent, or post-emergent in the following crops.

Adzuki Beans
Alfalfa Grown for Seed Production
Dry Common Beans of the species *Phaseolus vulgaris*^{1, 2}
Imazethapyr Tolerant Corn* (**Clearfield** Brands)
Lima Beans (Ontario only)
Processing Peas
Snap beans
Snow Peas
Soybeans

¹ Dry common bean varieties may vary in their tolerance to herbicides, including Rainbow Imazethapyr 240 SL. Since not all dry common bean varieties have been tested for tolerance to Rainbow Imazethapyr 240 SL, first use of Rainbow Imazethapyr 240 SL should be limited to a small area of each variety to confirm tolerance prior to adoption as a general field practice. Additionally, consult your seed supplier for information on the tolerance of specific varieties of dry common beans to Rainbow Imazethapyr 240 SL.

² Including, but not limited to: kidney, cranberry, yellow eye, Dutch brown, black and white beans.

MOISTURE REQUIREMENTS

As with most soil-applied herbicides, pre-emergent applications of Rainbow Imazethapyr 240 SL require moisture for activation. Soil-applied Rainbow Imazethapyr 240 SL requires sufficient water within 7 days of application to moisten the soil to a depth of 5 cm for activation. If adequate moisture is not received within 7 to 10 days of application, perform a shallow inter-row cultivation 5-8 cm deep using a roller or S-tine cultivator to control escaped weeds until the field receives adequate moisture. For early pre-plant applications (soybeans only), more than 7-10 days may elapse before the receipt of adequate precipitation to activate the herbicide and reduce the risk of weed escapes. Growers preferring surface applications of herbicides may choose this type of application of Rainbow Imazethapyr 240 SL.

PLANT BACK RESTRICTIONS AND ROTATIONAL CROPS

In cases of crop failure, replant only soybeans, all types of dry common beans of the species *Phaseolus vulgaris*, lima beans, adzuki beans, imazethapyr tolerant corn, and processing peas in the year of application. Winter wheat may also be re-planted in cases of crop failure or as a rotational crop 100 days following a Rainbow Imazethapyr 240 SL application. Soil preparation for re-planting should be no deeper than 10 cm.

Field corn, imazethapyr tolerant corn, soybeans, winter wheat, spring wheat, spring barley, all types of dry common beans of the species *Phaseolus vulgaris*, lima beans, adzuki beans, and processing peas may be planted the season following a Rainbow Imazethapyr 240 SL application. Conduct a field bioassay (a test strip grown to maturity) the year BEFORE growing any other crop.

CROP SPECIFIC RECOMMENDATIONS

CROP: SOYBEANS

Apply Rainbow Imazethapyr 240 SL as an early pre-plant, pre-plant incorporated, pre-emergent or post-emergent treatment in soybeans.

For fields that contain weeds other than those listed in the "Weed Control in Soybeans: Rainbow Imazethapyr 240 SL Alone" table, tank mix for broad-spectrum weed control (see "Herbicide Tank Mix Options – Soybeans"). The choice of product for tank mixing will depend on the specific weed(s) to be controlled. Consult the labels of the tank mix products to determine which product will provide control of the specific weeds present in the field.

Weed Control in Soybeans: Rainbow Imazethapyr 240 SL Alone

Weeds	Application Timing					
	Early Pre-plant (prior to weed emergence)	Early Pre-plant (emerged weeds prior to 2 true leaf stage)	Pre-plant Incorporated	Pre-Emergent	Early Post-Emergent (before weeds reach 2 true leaf stage)	Late Post-Emergent (maximum leaf stage in parenthesis)
Broadleaf Weeds						
lamb's-quarters	C	PC	C ²	C	PC	
redroot pigweed	C	C	C	C	C	C (12)
smartweed	C			C		
lady's-thumb	C			C ³		
wild mustard	C	C	C	C	C	
velvetleaf	C ¹	C ¹	C ¹	C ¹	C ¹	C (8)
ragweed, common	C	PC	PC ²	C ³	C ⁴	
ragweed, giant					PC ⁵	PC (10)

Eastern black nightshade	PC	C	C	C	C	
wild buckwheat		PC			C	
cocklebur		C ¹			C ^{1, 4}	
Grasses						
foxtail, green and yellow	C	C	C	C	C	C (4)
barnyard grass	C	PC	PC ²	C ³	C ⁴	C (6)
old witchgrass	C			C ³	C	
proso millet	PC	PC	PC	PC	PC	
crabgrass, large					PC	
Perennials						
yellow nutsedge					PC	

C = Control PC = Partial Control and Reduces Competition

¹ Some plants of velvetleaf and/or cocklebur that germinate deeper in the soil and emerge late may escape treatment.

² Tank mixing is recommended for fields with a history of heavy infestations of this weed species.

³ The higher label rate or tank mixing is recommended for fields with a history of heavy infestations of this weed species.

⁴ The higher label rate is required for heavy infestations of this weed species.

⁵ The higher label rate is required.

Herbicide Tank Mix Options – Soybeans

Tank Mix Option	Application Timing			
	Early Pre-plant	Pre-plant Incorporated	Pre- emergent	Early and Late Post-emergent
Roundup® ⁴ or Vantage™ Plus MAX Herbicide Solution	X ¹			X ^{1, 3}
Roundup® Transorb HC Liquid Herbicide + FirstRate® Herbicide	X ²			
Sencor® 75 DF / Sencor® 480 F	X ¹	X ^{1, 2}	X ¹	
Conquest® 480	X ^{1, 2}	X ^{1, 2}	X ^{1, 2}	
Conquest® 480 + glyphosate	X ^{1, 2}		X ^{1, 2}	
Conquest® 480 + ZIDUA® SC	X ^{1, 2}		X ^{1, 2}	
Conquest® 480 + ZIDUA SC + glyphosate	X ^{1, 2}		X ^{1, 2}	
Conquest® 480 + ZIDUA SC + ERAGON® LQ + MERGE® Adjuvant + glyphosate	X ^{1, 2}		X ^{1, 2}	
Conquest® 480 + PROWL® H2O Herbicide + glyphosate	X ^{1, 2}			
Conquest® 480 + PROWL H2O Herbicide + ERAGON LQ + MERGE Adjuvant + glyphosate	X ^{1, 2}			
Conquest® 480 + FRONTIER® MAX Herbicide + glyphosate	X ^{1, 2}		X ^{1, 2}	
Conquest® 480 + ERAGON LQ + MERGE Adjuvant + glyphosate	X ^{1, 2}		X ^{1, 2}	

Conquest® 480 + INTEGRITY® + MERGE Adjuvant + glyphosate	x1, 2		x1, 2	
Conquest® 480 + PROWL H2O Herbicide		x1, 2		
Conquest® 480 + FRONTIER MAX Herbicide		x1, 2	x1, 2	
Tricor LQ Herbicide		x1, 2		
Lexone® DF Herbicide Dispersible Granules		x1, 2	x1	
Treflan TM Liquid EC Herbicide / Rival® Emulsifiable Concentrate Liquid Herbicide / Trifluralin® 480 EC Herbicide / RHI Trifluralin 480 EC Liquid Herbicide		x1		
Liberty® 200 SN Herbicide				x2, 5
Basagran® Liquid Herbicide / Basagran® Forte Herbicide				x1, 2

¹ Refer to the label of the specific tank mix product for information regarding rates, recommendations, precautions and restrictions.

² Refer to Tank Mix Options under the appropriate Application Timing in this label.

³ Glyphosate Tolerant Soybeans only (i.e., varieties with the Roundup Ready®). Refer to CROP: GLYPHOSATE TOLERANT gene SOYBEANS section.

⁴ Roundup Transorb® HC Liquid Herbicide or Roundup WeatherMAX® with Transorb 2 Technology Liquid Herbicide.

⁵ For use on soybean varieties that are specially developed to be tolerant to glufosinate ammonium (e.g. LibertyLink® seeds).

Timing	Early Pre-plant Application - Soybeans Apply Rainbow Imazethapyr 240 SL up to 30 days before planting in conventional, reduced tillage or no-till soybeans. Rainbow Imazethapyr 240 SL alone may be applied as a surface application using this technique. Only one additional working of the soil to prepare the seedbed is recommended following the application. This final seedbed preparation should not work the soil deeper than 10 cm. Deeper tillage will result in reduced concentration of herbicide in the weed germination zone and reduction in weed control. DO NOT plow following the application.	
Rate	Early Pre-plant – Prior to Weed Emergence	420 mL/ha
	Early Pre-plant – to Emerged Weeds (before the weeds reach the 2 true leaf stage).	420 mL/ha + non-ionic surfactant. Non-ionic surfactant MUST BE ADDED. Liquid fertilizer added to the spray solution will provide quicker burndown of weeds.
Water Volume	100-400 L/ha	
Surfactant/Adjuvant	For Early Pre-plant – to Emerged Weeds: Non-ionic surfactant – 0.25% v/v (e.g. 2.5 L/1000 L of spray solution) Liquid fertilizer solution (10-34-0, 28-0-0 or 32-0-0) - 2 L/ha	
Weeds Controlled	See “Weed Control in Soybeans” table above.	
Remarks	Plant only soybeans during the season of application.	

Tank Mix Options	Rainbow Imazethapyr 240 SL may also be tank mixed with the herbicides listed in the “Herbicide Tank Mix Options – Soybeans” table. Always follow the most conservative rates, recommendations, precautions and restrictions on the tank mix labels.							
	For the combined tank mix with ROUNDUP TRANSORB HC LIQUID Herbicide and FIRSTRATE Herbicide, use the following:							
	<table><tr><td>Rates</td><td>Rainbow Imazethapyr 240 SL – 312 mL/ha ROUNDUP TRANSORB HC LIQUID Herbicide – 2.5 L/ha (0.900 kg ai/ha) FIRSTRATE Herbicide – 21 g/ha (0.0175 kg ai/ha)</td></tr><tr><td>Water Volume</td><td>100-400 L/ha</td></tr></table>		Rates	Rainbow Imazethapyr 240 SL – 312 mL/ha ROUNDUP TRANSORB HC LIQUID Herbicide – 2.5 L/ha (0.900 kg ai/ha) FIRSTRATE Herbicide – 21 g/ha (0.0175 kg ai/ha)	Water Volume	100-400 L/ha		
	Rates	Rainbow Imazethapyr 240 SL – 312 mL/ha ROUNDUP TRANSORB HC LIQUID Herbicide – 2.5 L/ha (0.900 kg ai/ha) FIRSTRATE Herbicide – 21 g/ha (0.0175 kg ai/ha)						
	Water Volume	100-400 L/ha						
	Comments: This tank mix provides control of emerged weeds found on the ROUNDUP TRANSORB HC LIQUID Herbicide label and residual control of germinating weeds found on the Rainbow Imazethapyr 240 SL and the FIRSTRATE Herbicide labels. Temporary crop injury may occur however, yield will not normally be affected. Avoid sprayer overlap. Severe crop injury will occur.							
	For other tank mixes with Conquest 480 applied in No or Minimum Tillage systems, use the following:							
		Products	Rates per hectare and soil types					
			COARSE	MEDIUM	FINE			
		Rainbow Imazethapyr 240 SL + Conquest 480 ¹	Do not use	312 – 420 mL 800 mL – 1.5 L	312 – 420 mL 960 mL – 2.0 L			
		Rainbow Imazethapyr 240 SL + Conquest 480 ¹ + glyphosate	Do not use	312 – 420 mL 800 mL – 1.5 L See glyphosate label	312 – 420 mL 960 mL – 2.0 L See glyphosate label			
		Rainbow Imazethapyr 240 SL + Conquest 480 ¹ + ZIDUA SC	Do not use	312 – 420 mL 800 mL – 1.5 L 120 – 240 mL OR 332 – 417 mL ²	312 – 420 mL 960 mL – 2.0 L 120 – 240 mL OR 493 mL			
		Rainbow Imazethapyr 240 SL + Conquest 480 ¹ + ZIDUA SC + glyphosate	Do not use	312 – 420 mL 800 mL – 1.5 L 120 – 240 mL OR 332 – 417 mL ² See glyphosate label	312 – 420 mL 960 mL – 2.0 L 120 – 240 mL OR 493 mL See glyphosate label			
		Rainbow Imazethapyr 240 SL + Conquest 480 ¹ + ZIDUA SC + ERAGON LQ ³ + MERGE Adjuvant + glyphosate	Do not use	312 – 420 mL 800 mL – 1.5 L 120 – 240 mL OR 332 – 417 mL ² 73 mL 1 L 900 g ae	312 – 420 mL 960 mL – 2.0 L 120 – 240 mL OR 493 mL 73 mL 1L 900 g ae			
	Rainbow Imazethapyr 240 SL + Conquest 480 ¹ + PROWL H2O Herbicide + ERAGON LQ ³ + MERGE Adjuvant + glyphosate	Do not use	312 mL 800 mL – 1.5 L 2.2 L 73 mL 1 L 900 g ae	312 mL 960 mL – 2.0 L 2.2 L 73 mL 1L 900 g ae				
	Rainbow Imazethapyr 240 SL + Conquest 480 ¹ + FRONTIER MAX	Do not use	312 – 420 mL 800 mL – 1.5 L 756 – 963 mL 900 g ae	312 – 420 mL 960 mL – 2.0 L 756 – 963 mL 900 g ae				

	Herbicide + glyphosate				
	Rainbow Imazethapyr 240 SL + Conquest 480 ¹ + ERAGON LQ ³ + MERGE Adjuvant + glyphosate	Do not use	312 – 420 mL 800 mL – 1.5 L 73 mL 1 L 900 g ae	312 – 420 mL 960 mL – 2.0 L 73 mL 1L 900 g ae	
	Rainbow Imazethapyr 240 SL + Conquest 480 ¹ + INTEGRITY ³ + MERGE Adjuvant + glyphosate	Do not use	312 – 420 mL 800 mL – 1.5 L 370 mL 1 L See glyphosate label	312 – 420 mL 960 mL – 2.0 L 370 mL 1L See glyphosate label	
	1 Use the higher rates when weeds are dense and/or soils with high organic matter (over 4%) and at the heavy end of the soil texture class. 2 The rate of ZIDUA SC for medium to fine soil depends on soil organic matter levels, refer to ZIDUA SC label for rates. 3 Apply pre-plant from 21 days prior to planting up to the day of planting.				

Timing	Pre-plant Incorporated Application - Soybeans Incorporate Rainbow Imazethapyr 240 SL evenly throughout the top 5 cm of the soil profile. Incorporation may be achieved with a double pass using discs or cultivator operated at 8 to 12 kph with the second pass at an angle to the first. Cultivators must have 3 or 4 rows of flexible sweeps staggered and spaced less than 15 cm apart followed by a drag or rolling basket to ensure no soil is left unturned.
Rate	312 mL/ha
Water Volume	100-400 L/ha
Weeds Controlled	See “Weed Control in Soybeans” table above.
Remarks	DO NOT apply Rainbow Imazethapyr 240 SL as a pre-plant incorporated application more than one year in sequence. Allow at least 24 months between pre-plant incorporated applications.

Timing	Pre-emergent Application - Soybeans Pre-emergent applications of Rainbow Imazethapyr 240 SL may be applied before the crop and weeds emerge.			
Rate	312 - 420 mL/ha			
Water Volume	100-400 L/ha			
Weeds Controlled	See “Weed Control in Soybeans” table above.			
Tank Mix Options	For fields that contain heavy infestations of common ragweed, old witchgrass, barnyard grass, lady's- thumb or weeds other than those listed under “Rainbow Imazethapyr 240 SL Alone”, tank mixing may be required to provide broad-spectrum weed control (see “Herbicide Tank Mix Options – Soybeans” table). Follow the most conservative rates, recommendations, precautions and restrictions on the tank mix labels. For other tank mixes with Conquest 480 applied in No or Minimum Tillage systems, use the following:			
	Products	Rates per hectare and soil types		
		COARSE	MEDIUM	FINE
	Rainbow Imazethapyr 240 SL + Conquest 480 ¹	Do not use	312 – 420 mL 800 mL – 1.5 L	312 – 420 mL 960 mL – 2.0 L
	Rainbow Imazethapyr 240 SL + Conquest 480 ¹ + glyphosate	Do not use	312 – 420 mL 800 mL – 1.5 L See glyphosate label	312 – 420 mL 960 mL – 2.0 L See glyphosate label
	Rainbow Imazethapyr 240 SL + Conquest 480 ¹ + ZIDUA SC	Do not use	312 – 420 mL 800 mL – 1.5 L 120 – 240 mL OR 332 – 417 mL ²	312 – 420 mL 960 mL – 2.0 L 120 – 240 mL OR 493 mL
	Rainbow Imazethapyr 240 SL + Conquest 480 ¹ + ZIDUA SC + glyphosate	Do not use	312 – 420 mL 800 mL – 1.5 L 120 – 240 mL OR 332 – 417 mL ² See glyphosate label	312 – 420 mL 960 mL – 2.0 L 120 – 240 mL OR 493 mL See glyphosate label
	Rainbow Imazethapyr 240 SL + Conquest 480 ¹ + ZIDUA SC + ERAGON LQ ³ + MERGE Adjuvant + glyphosate	Do not use	312 – 420 mL 800 mL – 1.5 L 120 – 240 mL OR 332 – 417 mL ² 73 mL 1 L 900 g ae	312 – 420 mL 960 mL – 2.0 L 120 – 240 mL OR 493 mL 73 mL 1L 900 g ae
	Rainbow Imazethapyr 240 SL + Conquest 480 ¹ + FRONTIER MAX Herbicide ⁴	Do not use	312 mL 800 mL – 1.5 L 756 – 963 mL	312 mL 960 mL – 2.0 L 756 – 963 mL
	Rainbow Imazethapyr 240 SL + Conquest 480 ¹ + FRONTIER MAX Herbicide ⁴ + glyphosate	Do not use	312 mL 800 mL – 1.5 L 756 – 963 mL See glyphosate label	312 mL 960 mL - 2.0 L 756 – 963 mL See glyphosate label
	Rainbow Imazethapyr 240 SL + Conquest 480 ¹ + ERAGON LQ ³ + MERGE Adjuvant + glyphosate	Do not use	312 – 420 mL 800 mL – 1.5 L 73 mL 1 L 900 g ae	312 – 420 mL 960 mL – 2.0 L 73 mL 1L 900 g ae

	Rainbow Imazethapyr 240 SL + Conquest 480 ¹ + INTEGRITY ³ + MERGE Adjuvant	Do not use	312 – 420 mL 800 mL – 1.5 L 370 mL 1 L	312 – 420 mL 960 mL – 2.0 L 370 mL 1L
--	---	------------	---	--

	+ glyphosate		See glyphosate label	See glyphosate label
--	--------------	--	----------------------	----------------------

¹ Use the higher rates when weeds are dense and/or soils with high organic matter (over 4%) and at the heavy end of the soil texture class.
² The rate of ZIDUA SC for medium to fine soil depends on soil organic matter levels, refer to ZIDUA SC label for rates.
³ For Pre-emergent timing, apply prior to when the soybeans cause the ground to crack and no more than 3 days after planting
⁴ The rate of FRONTIER MAX Herbicide depends on soil type and soil organic matter levels, refer to FRONTIER MAX Herbicide label for rates.

For other tank mixes with Conquest 480 applied in Conventional Tillage systems, use the following:

Products	Rates per hectare and soil types		
	COARSE	MEDIUM	FINE
Rainbow Imazethapyr 240 SL + Conquest 480 ¹	312 – 420 mL 850 mL – 1.1 L	312 – 420 mL 1.1 – 1.75 L	312 – 420 mL 1.75 – 2.25 L
Rainbow Imazethapyr 240 SL + Conquest 480 ¹ + ZIDUA SC	312 – 420 mL 850 mL – 1.1 L 120 – 240 mL OR 250 mL	312 – 420 mL 1.1 – 1.75 L 120 – 240 mL OR 332 – 417 mL ²	312 – 420 mL 1.75– 2.25 L 120 – 240 mL OR 493 mL ²
Rainbow Imazethapyr 240 SL + Conquest 480 ¹ + FRONTIER MAX Herbicide ³	312 – 420 mL 850 mL – 1.1 L 756 – 963 mL	312 – 420 mL 1.1 – 1.75 L 756-963 mL	312 – 420 mL 1.2 – 1.75 L 756-963 mL

¹ The rate of Conquest 480 depends on soil organic matter levels. See rates for Conquest 480 alone in Conquest 480 label.
² The rate of ZIDUA SC for medium to fine soil depends on soil organic matter levels, refer to ZIDUA SC label for rates.
³ The rate of FRONTIER MAX Herbicide depends on soil type and soil organic matter levels, refer to FRONTIER MAX Herbicide label for rates.

Timing	Early and Late Post-emergent Application - Soybeans Apply Rainbow Imazethapyr 240 SL early post-emergent after the crop has emerged and before the weeds reach the 2 true leaf stage. Rainbow Imazethapyr 240 SL may also be applied late post-emergent after the crop has emerged for control of certain weed species up to the growth stages indicated in the “Weed Control in Soybeans: Rainbow Imazethapyr 240 SL Alone” table. To minimize weed competition with the crop, application should be made as early as possible after weed emergence.	
Rate	Early Post-emergent – (before the weeds reach the 2 true leaf stage)	312-420 mL/ha + non-ionic surfactant + liquid fertilizer
	Late Post-emergent	420 mL/ha + non-ionic surfactant + liquid fertilizer
Water Volume	100-400 L/ha	

Surfactant/ Adjuvant	Non-ionic surfactant – 0.25% v/v (e.g. 2.5 L/1000 L of spray solution) Liquid fertilizer solution (10-34-0, 28-0-0 or 32-0-0) - 2 L/ha
Weeds Controlled	See “Weed Control in Soybeans” table above.
Remarks	Addition to the spray solution of a non-ionic surfactant plus liquid fertilizer is essential for post- emergent application, improving uptake of the product by weeds resulting in improved herbicidal activity. Avoid applications when weeds and/or crop are under growth stress. Stunting of the crop may result following post-emergence application. This condition is the result of stem inter-node shortening but will not result in yield reduction.
Tank Mix Options	Rainbow Imazethapyr 240 SL may also be tank mixed with the herbicides listed in the “Herbicide Tank Mix Options – Soybeans” table. Always follow the most conservative rates, recommendations, precautions and restrictions on the tank mix labels. Comments: Use the 312 mL/ha rate of Rainbow Imazethapyr 240 SL when tank mixed with BASAGRAN LIQUID Herbicide or BASAGRAN FORTE Herbicide. For Rainbow Imazethapyr 240 SL plus BASAGRAN LIQUID Herbicide, a non-ionic surfactant at the rate of 0.25% (v/v) and fertilizer solution (10-34-0, 28-0-0 or 32-0-0) at the rate of 2 L/ha MUST BE ADDED to the spray solution. For Rainbow Imazethapyr 240 SL plus BASAGRAN FORTE Herbicide, only fertilizer solution MUST BE ADDED at the rate of 2 L/ha.

CROP: GLYPHOSATE TOLERANT SOYBEANS
(i.e., varieties with the Roundup Ready® gene)

Timing	Post-emergent Application For broadleaf and grass weeds other than those listed in “Weed Control in Soybeans: Rainbow Imazethapyr 240 SL Alone” table, Rainbow Imazethapyr 240 SL may be tank mixed with Roundup¹ or Vantage Plus MAX Herbicide Solution (glyphosate) and applied to Roundup Ready soybeans. Follow the rates, application timings, recommendations, precautions and restrictions on the glyphosate label. Refer to the glyphosate label for adjuvant recommendations. ¹ Roundup Transorb HC Liquid Herbicide or Roundup WeatherMax with Transorb 2 Technology Liquid Herbicide.
Rainbow Imazethapyr 240 SL Rate	312 mL/ha
Water Volume	100-400 L/ha

Alternative Rainbow Imazethapyr 240 SL Rate	Rainbow Imazethapyr 240 SL at 160 - 210 mL/ha Tank-mixing Rainbow Imazethapyr 240 SL at 160 - 210 mL/ha with glyphosate will provide residual control of late germinating eastern black nightshade, common lamb's quarters, redroot pigweed, velvetleaf, fall panicum and wild proso millet. Use 900 g ae/ha of glyphosate¹ and apply up to and including the 3rd trifoliate leaf stage of Roundup Ready soybeans in 100-200 litres per hectare of clean water. Use higher rate for heavier infestations. Consult the glyphosate label for weeds controlled by glyphosate, as well as for further recommendations, precautions and restrictions. ¹ 900 g ae glyphosate/ha is equivalent to 2.5 L/ha of Roundup Transorb HC Liquid Herbicide or 1.67 L/ha of Roundup WeatherMax with Transorb 2 Technology Liquid Herbicide, or 1.88 L/ha of Vantage Plus MAX Herbicide Solution.
Remarks	WARNING Apply Rainbow Imazethapyr 240 SL + glyphosate herbicide tank mix ONLY to glyphosate tolerant soybeans, i.e. varieties with the Roundup Ready® gene. SOYBEANS WHICH ARE NOT DESIGNATED AS GLYPHOSATE TOLERANT WILL BE DAMAGED OR DESTROYED BY THIS TREATMENT.

CROP: GLUFOSINATE AMMONIUM TOLERANT (LIBERTYLINK®) SOYBEANS
(British Columbia and Eastern Canada Only)

Timing	Post-emergent Application For control of broadleaf and grass weeds other than those listed in the "Weed Control in Soybeans: Rainbow Imazethapyr 240 SL Alone" table, Rainbow Imazethapyr 240 SL may be tank mixed with Liberty 200 SN Herbicide and applied to glufosinate ammonium tolerant (LibertyLink) soybeans from the cotyledon to the flowering stage of the crop. Consult the Liberty 200 SN Herbicide label for the recommended leaf stage of weeds at application for best weed control.
Rates	312 mL/ha Rainbow Imazethapyr 240 SL + 1.5 - 2.5 L/ha Liberty 200 SN Herbicide Tank-mix Rainbow Imazethapyr 240 SL with the appropriate rate of Liberty 200 SN Herbicide for the targeted weed species. Refer to the Liberty 200 SN Herbicide label for rates, weeds controlled, further recommendations, precautions and restrictions.
Water Volume	Minimum of 110 L/ha
Remarks	WARNING Apply Rainbow Imazethapyr 240 SL plus Liberty 200 SN Herbicide tank-mix ONLY to glufosinate ammonium tolerant soybeans (e.g. LibertyLink seeds). Application of Liberty 200 SN Herbicide to non- tolerant soybeans will result in severe crop injury or death of the crop.

CROP: EDIBLE BEANS (including dry common beans of the species *Phaseolus vulgaris*^{1, 2}, lima beans (Ontario only) and adzuki beans)

Apply Rainbow Imazethapyr 240 SL as a pre-emergent treatment in lima beans (Ontario only), dry common beans of the species *Phaseolus vulgaris* and adzuki beans, and as a pre-plant incorporated treatment in dry common beans of the species *Phaseolus vulgaris* and adzuki beans.

¹ Dry common bean varieties may vary in their tolerance to herbicides, including Rainbow Imazethapyr 240 SL. Since not all dry common bean varieties have been tested for tolerance to Rainbow Imazethapyr 240 SL, first use of Rainbow Imazethapyr 240 SL should be limited to a small area of each variety to confirm tolerance prior to adoption as a general field practice. Additionally, consult your seed supplier for information on the tolerance of specific varieties of dry common beans to Rainbow Imazethapyr 240 SL.

² Including, but not limited to: kidney, cranberry, yellow eye, Dutch brown, black and white beans.

Weed Control in Edible Beans: Rainbow Imazethapyr 240 SL Alone

Weeds	Application Timing	
	Pre-emergent	Pre-plant Incorporated (dry common beans of the species <i>Phaseolus vulgaris</i> and adzuki beans only)
Broadleaf Weeds		
lamb's-quarters	C	C
redroot pigweed	C	C
smartweed	C	C
lady's-thumb	C	C
wild mustard	C	C
velvetleaf	C ¹	C ¹
ragweed, common	PC	PC
Eastern black nightshade	C	C
Grasses		
foxtail, green and yellow	C	C
proso millet	PC	PC

C = Control PC = Partial Control and Reduced Competition

¹ Some plants of velvetleaf that germinate deeper in the soil and emerge late may escape treatment.

Timing	Pre-emergent Application – Edible Beans
Rate	312 mL/ha
Water Volume	100-400 L/ha
Weeds Controlled	See “Weed Control in Edible Beans” table above.
Tank Mix Options	For fields that contain heavy infestations of broadleaf weeds other than those listed under “Rainbow Imazethapyr 240 SL Alone”, tank mixing may be required to provide broad-spectrum weed control. Cranberry and Kidney Beans Tank mix Rainbow Imazethapyr 240 SL with 1.15 – 1.75 L/ha (1.05-1.60 kg ai/ha) of DUAL® MAGNUM Herbicide to control labelled broadleaf weeds and grasses in cranberry beans and kidney beans. Refer to the DUAL MAGNUM Herbicide label for additional recommendations, precautions and restrictions not specified on this label.

Timing	Pre-plant Incorporated Application (dry common beans of the species <i>Phaseolus vulgaris</i> and adzuki beans only)
Rate	312 mL/ha
Water Volume	100-400 L/ha
Weeds Controlled	See “Weed Control in Edible Beans” table above.

Tank Mix Options	White Beans Tank mix Rainbow Imazethapyr 240 SL with 1.25-2.4 L/ha of TREFLAN LIQUID EC Herbicide to control a broader spectrum of weeds in white beans only. Follow recommendations, precautions and restrictions on the TREFLAN LIQUID EC Herbicide label. Consult the tank mix partner label for weeds controlled other than those listed for Rainbow Imazethapyr 240 SL alone. Dry Common Beans of the Species <i>Phaseolus vulgaris</i> ^{1,2} Tank mix Rainbow Imazethapyr 240 SL with 1.15 – 1.75 L/ha (1.05 - 1.60 kg ai/ha) of DUAL® Magnum Herbicide or DUAL II MAGNUM Herbicide to control labelled broadleaf weeds and grasses in dry common beans. Refer to the DUAL Magnum Herbicide or DUAL II MAGNUM Herbicide label for additional recommendations, precautions and restrictions not specified on this label. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels. ¹ NOTE: Dry common bean varieties may vary in their tolerance to herbicides, including the tank-mix of Rainbow Imazethapyr 240 SL plus DUAL MAGNUM Herbicide or DUAL II MAGNUM Herbicide. Since not all dry common bean varieties have been tested for tolerance to the tank-mix of Rainbow Imazethapyr 240 SL plus DUAL MAGNUM Herbicide or DUAL II MAGNUM Herbicide, first use of this tank-mix should be limited to a small area of each variety to confirm tolerance prior to adoption as a general field practice. Additionally, consult your seed supplier for information on the tolerance of specific varieties of dry common beans to the tank-mix of Rainbow Imazethapyr 240 SL plus DUAL MAGNUM Herbicide or DUAL II MAGNUM Herbicide. ² CAUTION: Beans should be seeded at least 4 cm deep or crop injury may result.
------------------	---

CROP: IMAZETHAPYR TOLERANT CORN* (i.e., Clearfield® BRANDS)

Apply Rainbow Imazethapyr 240 SL as a pre-plant incorporated, pre-emergent or early post-emergent treatment on imazethapyr tolerant corn.

Tank-mixing may be required to provide broad-spectrum weed control for fields that contain weeds other than those listed in the “Weed Control in Imazethapyr Tolerant Corn: Rainbow Imazethapyr 240 SL Alone” table below. Selection of product for tank mixing will depend on the specific weed(s) that are to be controlled. Consult the labels of the tank mix products to determine which will provide control of the specific weeds present in the field.

Weed Control in Imazethapyr Tolerant Corn: Rainbow Imazethapyr 240 SL Alone

Weeds	Application Timing		
	Pre-plant Incorporated	Pre-emergent	Early Post-emergent
Broadleaf Weeds			
lamb's-quarters	C ²	PC	PC
redroot pigweed	C	C	C
velvetleaf	C ¹	C ^{1, 2}	C ^{1, 2}
Eastern black nightshade	C	C	C
Grasses			
foxtail, green and yellow	C	PC	C
barnyard grass	PC	C	C ²

C = Control PC = Partial Control and Reduced Competition

¹ Some plants of velvetleaf that germinate deeper in the soil and emerge late may escape treatment.

² Tank mixing is required for fields with a history of heavy infestations of this weed species.

Timing	Pre-plant Incorporated Application – Imazethapyr Tolerant Corn Incorporate Rainbow Imazethapyr 240 SL evenly throughout the top 5 cm of the soil profile. Incorporation may be achieved with a double pass using discs or cultivator operated at 8 to 12 kph with the second pass at an angle to the first. Cultivators must have 3 or 4 rows of flexible sweeps staggered and spaced less than 15 cm apart followed by a drag or rolling basket to ensure no soil is left unturned.
Rate	312 mL/ha
Water Volume	100-400 L/ha
Weeds Controlled	See “Weed Control in Imazethapyr Tolerant Corn” table above.
Remarks	DO NOT apply Rainbow Imazethapyr 240 SL as a pre-plant incorporated application more than one year in sequence. Allow at least 24 months between pre-plant incorporated applications.
Tank Mix Options	Tank mix Rainbow Imazethapyr 240 SL with atrazine to provide a broader spectrum of weed control. Follow rates, recommendations, precautions and restrictions on the atrazine label.

Timing	Pre-emergent Application – Imazethapyr Tolerant Corn Pre-emergent applications of Rainbow Imazethapyr 240 SL may be applied before the crop and weeds emerge.
Rate	312 mL/ha
Water Volume	100-400 L/ha

Weeds Controlled	See “Weed Control in Imazethapyr Tolerant Corn” table above.
Tank Mix Options	Tank-mix Rainbow Imazethapyr 240 SL with either PROWL® H2O Herbicide or atrazine herbicide to provide a broader spectrum of weed control. Follow rates, recommendations, precautions and restrictions on the PROWL® H2O Herbicide or atrazine herbicide label.

Timing	Early Post-emergent Application – Imazethapyr Tolerant Corn Apply Rainbow Imazethapyr 240 SL before or after the crop has emerged and before the weeds reach the 2 true leaf stage.
Rate	312 mL/ha + non-ionic surfactant + liquid fertilizer
Water Volume	100-400 L/ha
Surfactant/ Adjuvant	Non-ionic surfactant – 0.25% v/v (e.g. 2.5 L/1000 L of spray solution) Liquid fertilizer solution (10-34-0, 28-0-0 or 32-0-0) - 2 L/ha
Weeds Controlled	See “Weed Control in Imazethapyr Tolerant Corn” table above.
Remarks	Addition to the spray solution of a non-ionic surfactant plus liquid fertilizer is essential for post- emergent application, improving uptake of the product by weeds resulting in improved herbicidal activity.
Tank Mixing	Tank mix Rainbow Imazethapyr 240 SL with either PROWL H2O Herbicide, atrazine or MARKSMAN® Herbicide to provide a broader spectrum of weed control. Follow rates, recommendations, precautions and restrictions on the PROWL H2O Herbicide, atrazine or MARKSMAN Herbicide label.

CROP: PROCESSING PEAS

Timing	Pre-emergent or Pre-plant Incorporated Application
Rate	312 mL/ha
Water Volume	200 L/ha
Weeds Controlled	Weeds found in the “Rainbow Imazethapyr 240 SL Alone: Weed Control in Soybeans” table.

CROP: SNOW PEAS

Timing	Pre-emergent or Pre-plant Incorporated Application
Rate	312 mL/ha
Water Volume	100-400 L/ha
Weeds Controlled	Weeds found in the “Rainbow Imazethapyr 240 SL Alone: Weed Control in Soybeans” table.

CROP: SNAP BEANS

Timing	Pre-emergent Application
Rate	312 mL/ha
Water Volume	200 L/ha

Weeds Controlled	Weeds found in the "Rainbow Imazethapyr 240 SL Alone: Weed Control in Soybeans" table.
------------------	--

CROP: ALFALFA GROWN FOR SEED PRODUCTION

Timing	Post-emergent Application Apply early post-emergent when weeds are less than 7.5 cm (3 in.) tall. Apply only after the crop has one fully developed trifoliate leaf.
Rate	312 – 420 mL/ha
Water Volume	200 L/ha
Surfactant/Adjuvant	A non-ionic surfactant at the rate of 0.25% (v/v) MUST BE ADDED to the spray solution (e.g., 2.5 L of surfactant per 1000 L of spray solution). Fertilizer solutions, 10-34-0, 28-0-0 or 32-0-0 MUST BE ADDED to the spray solution at the rate of 2 L/ha to provide quicker burndown of weeds.
Weeds Controlled	Weeds found in the "Rainbow Imazethapyr 240 SL Alone: Weed Control in Soybeans" table.

MIXING INSTRUCTIONS

1. Ensure the spray tank is clean before use. Follow the clean-out recommendations stated on the label of the product that was previously used.
2. Fill the spray tank one-half full to three-quarters full of water and start agitation.
3. Using a calibrated measuring device, add the required amount of tank mix partner (refer to the tank mixture section of each crop for tank mixtures).
4. Mix thoroughly.
5. Using a separate calibrated measuring device, add the required amount of Rainbow Imazethapyr 240 SL to the tank while agitating the spray solution.
6. While the solution remains agitating, add the required amount of non-ionic surfactant if required.
7. If required, add the required amount of liquid fertilizer (28-0-0, 10-34-0 or 32-0-0) to the spray solution.
8. Continue agitation while filling the remainder of the spray tank with water.
9. Clean the spray tank after use.

RESTRICTIONS AND LIMITATIONS

1. DO NOT over apply Rainbow Imazethapyr 240 SL. Over application may result in injury particularly if the crop is under stress.
2. Crop Pre-harvest Interval

Crop	Application to Harvest Interval (days)
Soybeans	100
Dry Common Beans of the species <i>Phaseolus vulgaris</i> and Adzuki Beans	100
Imazethapyr Tolerant Corn	100
Lima Beans	90
Processing Peas	50
Snow Peas	60
Snap Beans	40

3. CAUTION: DO NOT graze treated crops or cut for hay, sufficient data are not available to support such use.
4. DO NOT let spray drift contaminate crops in adjacent fields.
5. ONLY ONE (1) application of Rainbow Imazethapyr 240 SL may be made during the season.
6. DO NOT apply Rainbow Imazethapyr 240 SL as a pre-plant incorporated application in all crops more than one year in sequence. Allow at least 24 months between pre-plant incorporated applications.
7. Emerged weeds in pre-emergent applications which reach the 2 - 3 true leaf stage might be considered as escapes. Shallow cultivation or application of a post-emergent herbicide is recommended. DO NOT cultivate deeply.
8. DO NOT apply Rainbow Imazethapyr 240 SL when crop is under stress conditions because crop injury may result.
9. Post-emergent application of Rainbow Imazethapyr 240 SL to soybeans may cause stunting. This condition is the result of stem inter-node shortening and does not cause yield reductions if Rainbow Imazethapyr 240 SL has been used at label rates and following label recommendations.
10. Rainbow Imazethapyr 240 SL may cause stunting or delayed maturity in white beans and kidney beans. Stunting is the result of stem inter-node shortening and should not cause yield reductions if Rainbow Imazethapyr 240 SL has been used at label rates and following label recommendations.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, Rainbow Imazethapyr 240 SL is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to Rainbow Imazethapyr 240 SL and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of Rainbow Imazethapyr 240 SL or other Group 2 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area, if possible, by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Rainbow CropSciences Inc. at (630) 828-5088.