2026-2791
2026-08-12

Container label

GROUP 2 HERBICIDE

RAINBOW METSULFURON-METHYL 60 WG

60% WETTABLE GRANULES

COMMERCIAL

ACTIVE INGREDIENT: Metsulfuron-Methyl 60%

REGISTRATION NO.: **36073** PEST CONTROL PRODUCTS ACT

NET CONTENTS: 30 g to Bulk

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

For Chemical Emergency: Spill, Leak, Fire, Exposure, or Accident, Call CHEMTREC: 1-800-424-9300

Rainbow CropSciences Inc.
535 E Plainfield Road, Suite E
Willowbrook, IL 60527
(630) 828-5088

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

- Avoid breathing spray mist.
- DO NOT eat, drink or smoke/vape while using this product. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum/tobacco, smoking/vaping, using the toilet or handling mobile devices. Aircraft cockpit and vehicle cabs MUST be decontaminated regularly.
- Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab and/or cockpit.
- DO NOT allow the pilot to mix chemicals to be loaded onto the aircraft. The pilot is allowed to load premixed chemicals with a closed system. It is desirable that the pilot have communication capabilities at each treatment site at the time of application.
- If the product comes in contact with clothing/personal protective equipment (PPE), remove all contaminated clothing/PPE, wash the outside of gloves with soap and water before removing, wash contaminated skin with soap and water, and then dress in clean clothing. Launder contaminated clothing/PPE regularly and separately from other laundry. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Heavily contaminated or drenched clothing and other absorbent materials contaminated with this product should be discarded. DO NOT reuse them.
- Do not re-enter treated fields before 12 hours after application.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IN CASE OF EMERGENCY, CALL CHEMTREC: 1-800-424-9300 (24 HOURS)

TOXICOLOGICAL INFORMATION

Treat symptomatically.

ENVIRONMENTAL HAZARDS

Toxic to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application of this product when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the

Canada.ca website.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination

STORAGE

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed.

DISPOSAL:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

For information on disposal of unused, unwanted product, contact the registrant and the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill and for cleanup of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

GROUP 2 HERBICIDE

Booklet

RAINBOW METSULFURON-METHYL 60 WG

60% WATER DISPERSIBLE GRANULES

COMMERCIAL

ACTIVE INGREDIENT: Metsulfuron-Methyl 60%

REGISTRATION NO.: **36073** PEST CONTROL PRODUCTS ACT

NET CONTENTS: 30g to Bulk

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

For Chemical Emergency: Spill, Leak, Fire, Exposure, or Accident, Call CHEMTREC: 1-800-424-9300

Rainbow CropSciences Inc.
535 E Plainfield Road, Suite E
Willowbrook, IL 60527
(630) 828-5088

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

- Avoid breathing spray mist.
- DO NOT eat, drink or smoke/vape while using this product. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum/tobacco, smoking/vaping, using the toilet or handling mobile devices. Aircraft cockpit and vehicle cabs MUST be decontaminated regularly.
- Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab and/or cockpit.
- DO NOT allow the pilot to mix chemicals to be loaded onto the aircraft. The pilot is allowed to load premixed chemicals with a closed system. It is desirable that the pilot have communication capabilities at each treatment site at the time of application.
- If the product comes in contact with clothing/PPE, remove all contaminated clothing/PPE, wash the outside of gloves with soap and water before removing, wash contaminated skin with soap and water, and then dress in clean clothing. Launder contaminated clothing/PPE regularly and separately from other laundry. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Heavily contaminated or drenched clothing and other absorbent materials contaminated with this product should be discarded. DO NOT reuse them.
- Do not re-enter treated fields before 12 hours after application.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Treat symptomatically.

ENVIRONMENTAL HAZARDS

Toxic to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application of this product when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body. Additional guidance can be found on the Runoff Mitigation portion of the Canada.ca website.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

FOR POSTEMERGENCE USE ON WHEAT (SPRING AND DURUM), BARLEY AND ESTABLISHED CREEPING RED FESCUE, ORCHARD GRASS, CRESTED AND INTERMEDIATE WHEAT GRASS (SEED OR FORAGE) (SOIL PH 7.9 OR LOWER), AND FOR CONTROL OF UNDESIRABLE BRUSH/WOODY PLANTS IN PASTURE, RANGELAND AND NON-CROP AREAS AND BROADLEAF WEEDS IN ROUGH TURF AND NON-CROP AREAS. FOR CONTROL OF WESTERN SNOWBERRY AND WILD ROSE IN PASTURE AND RANGELAND In Manitoba, Saskatchewan, Alberta, Peace River Region and Interior of British Columbia only

IMPORTANT

Injury to or loss of desirable trees or vegetation may result from failure to observe the following. Do not apply, drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts or similar areas. Prevent drift of spray to desirable plants..

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.
As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

APPLY RAINBOW METSULFURON-METHYL 60 WG WITH RECOMMENDED SURFACTANT.

Carefully observe sprayer clean-up instructions, as spray tank residue may damage crops other than wheat, barley or creeping red fescue.

PRIOR TO USE REFER TO “MINIMUM CROP ROTATION GUIDELINES” SECTION OF LABEL.

GENERAL INFORMATION

RAINBOW METSULFURON-METHYL 60 WG is recommended for postemergence use on wheat (spring and durum), barley and established creeping red fescue, orchard grass, crested and intermediate wheat grass (seed or forage) inManitoba, Saskatchewan, Alberta, Peace River Region and Interior of British Columbia only (soil pH of 7.9 or lower). In addition, RAINBOW METSULFURON-METHYL 60 WG is recommended for use on pasture, non-crop areas, rough turf and rangeland in the same geographic region (no soil pH restriction).

RAINBOW METSULFURON-METHYL 60 WG is to be mixed in water with a recommended surfactant and applied as a uniform broadcast spray. Use for selective control or suppression of certain broadleaf weeds in spring wheat (including durum), barley and established creeping red fescue (seed or forage) and for control of Western Snowberry and Wild Rose in pasture and rangeland. Application as a foliar spray can be used to control undesirable brush/woody plants in pasture, rangeland, and non-crop areas, and for control of broadleaf weeds in rough turf and non-crop areas. RAINBOW METSULFURON-METHYL 60 WG is non-corrosive, non-flammable, non-volatile and does not freeze. Wheat, barley or creeping red fescue may be grazed by or fed to livestock any time after treatment.

RAINBOW METSULFURON-METHYL 60 WG is a 60% active ingredient herbicide formulated as water dispersible granules.

Prior to using RAINBOW METSULFURON-METHYL 60 WG, careful consideration should be given to crop rotation plans. Crops other than those listed in the "Minimum Crop Rotation Guidelines" section can be very sensitive to low concentrations of RAINBOW METSULFURON-METHYL 60 WG in the soil. Refer to the "Minimum Crop Rotation Guidelines" section of this label to plan your RAINBOW METSULFURON-METHYL 60 WG use and crop rotations.

Rainfall, soil temperature and soil pH are important factors affecting RAINBOW METSULFURON-METHYL 60 WG breakdown in the soil. RAINBOW METSULFURON-METHYL 60 WG breakdown is more rapid under conditions of moist soil, warm soil temperatures and low soil pH. The breakdown process is slower under conditions of dry soil, cool soil temperatures and high soil pH.

RAINBOW METSULFURON-METHYL 60 WG must be applied with a recommended surfactant early postemergence to the main flush of actively growing broadleaf weeds. For pasture and rangeland applications, RAINBOW METSULFURON-METHYL 60 WG should be applied between mid-June and mid-August after the target species have leafed out, but before fall colouration begins. Cattle may graze the treated areas on the day of treatment.

Warm, moist growing conditions promote active weed growth and enhance the activity of RAINBOW METSULFURON-METHYL 60 WG by allowing maximum foliar uptake and contact activity. If cold, dry conditions prevail, delay treatment until active weed growth resumes. Brush and weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and re-growth may occur. RAINBOW METSULFURON-METHYL 60 WG may only be applied with ground spray equipment. For best results, ensure good spray coverage of target weeds.

RAINBOW METSULFURON-METHYL 60 WG rapidly inhibits the growth of susceptible weeds, however, typical symptoms (discolouration) of dying weeds may not be noticeable for 1 to 3 weeks after application depending upon growing conditions and weed susceptibility. Degree of control and duration of effect depend on weed sensitivity, weed size, growing conditions, soil pH and spray coverage.

TANK MIXES

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Rainbow CropSciences Inc. at (630) 828-5088 for information before applying any tank mix that is not specifically recommended on this label.

SURFACTANTS FOR USE WITH RAINBOW METSULFURON-METHYL 60 WG

IPCO AG-SURF® ORIGINAL, WEEDAWAY AG-SURF LIQUID SPRAY ADJUVANT, Agral® 90 Non-Ionic Wetting & Spreading Agent Liquid, AGRAL*90 NON-IONIC LIQUID WETTING & SPREADING AGENT, CITOWETT® PLUS, SUPER SPREADER ADJUVANT, and Liberate Adjuvant are recommended for use with RAINBOW METSULFURON-METHYL 60 WG, RAINBOW METSULFURON-METHYL 60 WG + 2,4-D, and RAINBOW METSULFURON-METHYL 60 WG + MCPA. Use the surfactant at the rate of 2 L per 1000 L spray solution (0.2% v/v). See section on MIXING INSTRUCTIONS.

CROP LEAF STAGING: WHEAT, BARLEY AND CREEPING RED FESCUE

Apply RAINBOW METSULFURON-METHYL 60 WG or RAINBOW METSULFURON-METHYL 60 WG tank mixtures to wheat, barley, or creeping red fescue when the crop is between the stages listed in the table below.

Post-Emergent Treatment	Crop Leaf Staging	Crop
RAINBOW METSULFURON-METHYL 60 WG	2 leaf to flag leaf (shot blade)	Wheat, barley, Creeping red fescue
RAINBOW METSULFURON-METHYL 60 WG + 2,4-D (amine or ester)	Full 3 leaf to just before flag leaf	Wheat, barley
RAINBOW METSULFURON-METHYL 60 WG + MCPA (amine or ester)	Full 3 leaf to just before flag leaf	Wheat, barley

HOW TO APPLY TO WHEAT (SPRING OR DURUM), BARLEY OR ESTABLISHED CREEPING RED FESCUE (SEED OR FORAGE)

Apply RAINBOW METSULFURON-METHYL 60 WG to the first main flush of actively growing weeds, when the crop is in the leaf stage specified in the section CROP STAGE. Apply before the crop canopy is dense enough to prevent thorough coverage of the weeds. Apply with ground equipment only. Make only one application per growing season. For further information on control of Canada thistle, sow thistle, Russian thistle, lamb's quarters or wild buckwheat, refer to the section SPECIFIC WEED CONTROL RECOMMENDATIONS.

DO NOT apply by air.

RATES TO APPLY:
RAINBOW METSULFURON-METHYL 60 WG 7.5 g/ha (plus surfactant)

HOW TO APPLY TO WHEAT (SPRING), OR BARLEY

Apply RAINBOW METSULFURON-METHYL 60 WG or RAINBOW METSULFURON-METHYL 60 WG tank mixtures to the first main flush of actively growing weeds, when the crop is in the leaf stage specified in the section CROP STAGE. Apply before the crop canopy is dense enough to prevent thorough coverage of the weeds. Apply with ground equipment only. Make only one application per growing season. For further information on control of Canada thistle, sow thistle, Russian thistle, lamb's quarters or wild buckwheat, refer to the section SPECIFIC WEED CONTROL RECOMMENDATIONS.

DO NOT apply by air.

RATES TO APPLY:
RAINBOW METSULFURON-METHYL 60 WG 7.5 g/ha (plus surfactant)

RAINBOW METSULFURON-METHYL 60 WG + 2,4-D (use one of the following, plus surfactant):

RAINBOW METSULFURON-METHYL 60 WG	5.0 or 7.5 g/ha
Plus	
2,4-D Acid equivalent	420 – 560 g/ha

Refer to the 2,4-D label for information on crop variety restrictions, precautions, and application information

RAINBOW METSULFURON-METHYL 60 WG + MCPA (use one of the following, plus surfactant):

RAINBOW METSULFURON-METHYL 60 WG	5.0 or 7.5 g/ha
Plus	
MCPA Amine 500 HERBICIDE	0.7 – 1.1 L/ha
Or	
MPOWER MCPA Ester 500	0.7 – 1.1 L/ha

Refer to the MCPA label for information on crop variety restrictions, precautions, and application information.

WEEDS CONTROLLED OR SUPPRESSED†

† Weed suppression is a visual reduction in weed competition (reduced population or vigour) as compared to an untreated area. Degree of suppression will vary with size of weed and environmental conditions prior to and following treatment.

RAINBOW METSULFURON-METHYL 60 WG 7.5 grams/ha + surfactant

Ball mustard	Prostrate pigweed
Blue bur	Redroot pigweed
Canada thistle†	Russian thistle†
Chickweed	Scentless chamomile
Common groundsel	Shepherd's purse
Corn spurry	Sow thistle (annual; perennial)†
Cow cockle	Stinkweed
Flixweed	Stork's bill
Green smartweed	Tartary buckwheat
Hemp-nettle	Toadflax†
Kochia	Volunteer canola °
Lady's thumb	Wild buckwheat†
Lamb's-quarters†	Wild mustard

° Volunteer canola: RAINBOW METSULFURON-METHYL 60 WG alone will not control imazethapyr tolerant canola varieties, eg. canola varieties with the Pursuit Smart* trait.

RAINBOW METSULFURON-METHYL 60 WG 5 grams/ha + 2,4-D (amine or ester) + Surfactant

Annual sunflower	Prickly lettuce
Ball mustard	Redroot pigweed
Canada thistle†	Russian pigweed
Chickweed	Russian thistle
Cow cockle	Shepherd's purse
Flixweed	Sow thistle†
Green smartweed	Stinkweed
Hemp-nettle	Sweetclover
Kochia	Toadflax†
Lamb's-quarters	Volunteer canola°
Narrow-leaved hawksbeard	Wild buckwheat†
(spring seedlings)	Wild mustard
Plantain	Wormseed mustard

° Including imazethapyr tolerant canola varieties, eg. Canola varieties with the Pursuit* Smart trait. Consult the 2,4-D label for appropriate use rates.

RAINBOW METSULFURON-METHYL 60 WG 7.5 grams/ha + 2,4-D (amine or ester) + Surfactant:

All of the weeds controlled or suppressed by RAINBOW METSULFURON-METHYL 60 WG 5 grams/ha + 2,4-D plus:

Bluebur	Prostrate pigweed
Common groundsel	Scentless chamomile
Corn spurry	Stork's bill
Lady's-thumb	Tartary buckwheat
Toadflax†	

RAINBOW METSULFURON-METHYL 60 WG 5.0 grams/ha + MCPA (amine or ester) + Surfactant:

Annual sunflower
Ball mustard
Canada thistle†
Chickweed
Cow cockle
Flixweed
Green smartweed
Hemp-nettle
Kochia
Lamb's-quarters
Plantain
Prickly lettuce
Redroot pigweed
Russian pigweed
Russian thistle
Shepherd's purse
Sow thistle†
Stinkweed
Sweet clover
Toadflax†
Tumble mustard
Volunteer canola°
Wild buckwheat†
Wild mustard
Wormseed mustard

° Including imazethapyr tolerant canola varieties, eg. canola varieties with the Pursuit* Smart trait. Use the highest rate of MCPA (1.1 litre/ha) when volunteer imazethapyr tolerant canola is the target weed.

RAINBOW METSULFURON-METHYL 60 WG 7.5 grams/ha + MCPA (amine or ester) + Surfactant:
All of the weeds controlled or suppressed by RAINBOW METSULFURON-METHYL 60 WG 5.0 grams/ha + MCPA, plus:

Bluebur
Common groundsel
Corn spurry
Lady's-thumb
Prostrate pigweed
Scentless chamomile
Stork's bill
Tartary buckwheat
Toadflax†

SPECIFIC WEED RECOMMENDATIONS:
Weeds must be actively growing at time of application or control may be reduced. Application should be made before the crop canopy prevents thorough coverage of the weeds.

CANADA THISTLE, SOW THISTLE – Apply when the majority of thistles have emerged. RAINBOW METSULFURON-METHYL 60 WG + surfactant should be applied before the thistles are more than 15 cm tall. RAINBOW METSULFURON-METHYL 60 WG + 2,4-D or RAINBOW METSULFURON-METHYL 60 WG + MCPA + surfactant may be applied up to the bud stage of the thistles. Thistles that emerge after application will not be controlled. A single application will prevent emerged thistles from competing with the crop.

WILD BUCKWHEAT – Apply RAINBOW METSULFURON-METHYL 60 WG + surfactant at the cotyledon to 3-leaf stage. RAINBOW METSULFURON-METHYL 60 WG + 2,4-D or RAINBOW METSULFURON-METHYL 60 WG + MCPA + surfactant should be applied at the 1-3 leaf stage. Under dry conditions, control may be reduced. Large weeds may re-grow after treatment.

LAMB'S-QUARTERS, RUSSIAN THISTLE – Apply RAINBOW METSULFURON-METHYL 60 WG + surfactant before these weeds are more than 8 cm tall.

HOW TO APPLY TO PASTURE, ROUGH TURF AND NON-CROP AREAS SUCH AS UTILITY RIGHTS OF WAY, ROADSIDES, INDUSTRIAL SITES, FENCE LINES):

FOR CONTROL OF BROADLEAF WEEDS

RAINBOW METSULFURON-METHYL 60 WG may be applied as a broadcast treatment to rough turf areas at 20, 25 and 30 grams per hectare with a recommended surfactant for season long weed control. RAINBOW METSULFURON-METHYL 60 WG may be tank mixed with 2,4-D to increase the spectrum of weeds controlled. Add a recommended surfactant at 0.2% v/v (2.0 litres per 1000 litres spray solution). Temporary chlorosis, height reduction or growth suppression of grass species may occur.

For best results apply RAINBOW METSULFURON-METHYL 60 WG post-emergence to young (less than 10 cm tall or across) annual weeds; biennial and perennial weeds up to the early bud stage. Weeds should be actively growing at time of application.

Apply to brush species as a full coverage spray to foliage and stems using equipment that will ensure uniform coverage.

Note: Severe injury may occur to desirable broadleaf species such as alfalfa and red clover.

WEEDS CONTROLLED OR SUPPRESSED†

† Weed suppression is a visual reduction in weed competition (reduced population or vigour) as compared to an untreated area. Degree of suppression will vary with size of weed and environmental conditions prior to and following treatment.

RAINBOW METSULFURON-METHYL 60 WG AT 20 g/ha:

Canada thistle†
Common tansy
Dandelion†
Kochia
Russian thistle
Scentless chamomile
Sow thistle†
Sweet clover

RAINBOW METSULFURON-METHYL 60 WG AT 25 g/ha:

Weeds controlled/suppressed at 20 g/ha, plus:

Western Snowberry

RAINBOW METSULFURON-METHYL 60 WG AT 30 g/ha:

Weeds controlled/suppressed at 25 grams per hectare, plus:

Dandelion (control)

RAINBOW METSULFURON-METHYL 60 WG at 20 or 30 g/ha may be tankmixed with 2,4-D amine or ester at 1.0 kg ai/ha to control the weeds above and the weeds that are controlled by this rate of 2,4-D (refer to the 2,4-D label). Add a recommended surfactant at 0.2% v/v (2.0 litres per 1000 litres spray solution).

PASTURE, RANGELAND & NON-CROP AREAS such as Utility Rights of Way, roadsides, industrial sites, fence lines.

FOR CONTROL OF UNDESIRABLE BRUSH/WOODY PLANT SPECIES

For control of Balsam Poplar and Willow, apply 100 grams of RAINBOW METSULFURON-METHYL 60 WG per hectare.

For control of Cherry and Trembling Aspen, apply 150 grams of RAINBOW METSULFURON-METHYL 60 WG per hectare.

Add a recommended surfactant such as IPCO AG-SURF* ORIGINAL, WEEDAWAY AG-SURF LIQUID SPRAY ADJUVANT, AGRAL* 90 NON-IONIC WETTING & SPREADING AGENT, AGRAL 90 NON-IONIC LIQUID WETTING & SPREADING AGENT, or CITOWETT* PLUS at 0.2% v/v (2 litres per 1000 litres of spray solution).

For best results, applications of RAINBOW METSULFURON-METHYL 60 WG should be made when brush species and weeds are actively growing. Complete coverage of all foliage and stems is required for brush control. Applications should be made after the target species have leafed out, but before fall colouration has begun.

Do not treat brush species that exceed 2.5 m in height, or control may be decreased. For woody plants exceeding this height, cut and spray regrowth.

Note: Where dense stands exist or where growth of the target species is advanced, use of the high volume foliar application method is recommended.

PASTURE AND RANGELAND

FOR CONTROL OF WESTERN SNOWBERRY AND WILD ROSE in the Prairie Provinces and British Columbia

For control of Western Snowberry apply 25 grams of RAINBOW METSULFURON-METHYL 60 WG per hectare.

For control of Wild Rose apply 30 grams of RAINBOW METSULFURON-METHYL 60 WG per hectare.

Add a recommended surfactant at 0.2% v/v (2.0 litres per 1000 litres spray solution).

Apply as a ground application only in 100 - 225 litres spray volume per hectare. Apply as a full coverage spray to foliage and stems using equipment that will assure uniform coverage.

GROUND APPLICATIONS:

Low Volume Foliar Broadcast
Apply to brush species as full coverage spray to foliage and stems in 100-500 litres spray volume per hectare, using properly calibrated equipment that will ensure uniform coverage. Direct the spray solution to thoroughly wet the foliage of the target plants, but not to the point of runoff.

High Volume Foliar Broadcast
For control of brush/woody plants up to 2.5 metres in height, use RAINBOW METSULFURON-METHYL 60 WG at rates of 100-150 grams per hectare in enough water to make 1000 litres of spray solution. Apply as a full coverage spray to foliage, stems and limbs using up to 2000 litres of total spray per hectare depending on plant species, height and density of growth, and on the type of spray equipment used. Do not apply more than 150 grams of RAINBOW METSULFURON-METHYL 60 WG per hectare.

Spray brush to the point of runoff. Coverage should be thorough to wet all foliage. To minimize spray drift, do not use pressures exceeding 1400 kPa at the spray nozzle. Use of a drift control system is suggested to minimize spray drift.

NOTE TO BUYER/USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR SPECIAL USE APPLICATIONS.

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Rainbow CropSciences Inc. under the User Requested Minor Use Label Expansion program. For these uses, Rainbow CropSciences Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

FORAGE GRASSES FOR FEEDS OR SEED PRODUCTION:

For post-emergent application to creeping red fescue, orchard grass, crested and intermediate wheat grass. These forage grasses should be established for a minimum of one year before applying RAINBOW METSULFURON-METHYL 60 WG. Optimum control of narrow-leaved hawk's beard and scentless chamomile is obtained if application is made to small, actively growing weeds.

Apply RAINBOW METSULFURON-METHYL 60 WG as a broadcast application at 7.5 g/ha post-emergent. Application should be made from the 2 leaf, up to the shot blade stage of the crop, and before canopy closure to permit thorough coverage of actively growing weeds.

Make only one application per growing season. Application should be made using ground equipment only.

Rotation options are determined by soil pH, the crop to be planted and a minimum interval. Refer to the recropping recommendation section as well as other application instructions.

ESTABLISHED CREEPING RED FESCUE FOR SEED:

For post-emergent broadleaf and grassy weed control, apply the following tankmix:

RAINBOW METSULFURON-METHYL 60 WG + ASSURE II Herbicide + Adjuvant:

RAINBOW METSULFURON-METHYL 60 WG	7.5 g/ha
Plus	
ASSURE® II Herbicide	0.5 – 0.75 L/ha
Plus	
SURE-MIX™ SURFACTANT	5.0 L/1000 L spray solution

This tankmix will control all label weeds listed on the respective RAINBOW METSULFURON-METHYL 60 WG and ASSURE II Herbicide labels.

Apply RAINBOW METSULFURON-METHYL 60 WG + ASSURE® II Herbicide tank mixtures post-emergent to the first main flush of actively growing weeds, when established creeping red fescue is in the 2 leaf to flag leaf (shot blade) stage. Apply before the crop canopy is dense enough to prevent thorough coverage of the weeds.

Apply with ground equipment only. Make only one application per growing season.

Refer to other sections of this label, and to the ASSURE II Herbicide labels for additional application and/or use precautions instructions.

PASTURE, RANGELAND, ROUGH TURF AND NON-CROP AREAS IN BRITISH COLUMBIA:

For weeds listed in the RAINBOW METSULFURON-METHYL 60 WG Directions for Use and for control of field scabious and perennial pepperweed:

RAINBOW METSULFURON-METHYL 60 WG	20 - 30 g/ha
Plus	
IPCO AG-SURF* ORIGINAL or WEEDAWAY AG-SURF LIQUID SPRAY ADJUVANT	2.0 L/1000 L spray solution (0.2% v/v)
Or	
AGRAL* 90 NON-IONIC WETTING & SPREADING AGENT LIQUID or AGRAL*90 NON-IONIC LIQUID WETTING & SPREADING AGENT	2.0 L/1000 L spray solution (0.2% v/v)
Or	
CITOWETT* PLUS	2.0 L/1000 L spray solution (0.2% v/v)

Apply the spray mixture post-emergent, only one application per growing season. Apply with ground equipment only. Apply a minimum of 100 litres/ha of total spray volume. For mixing instructions and use precautions, refer the RAINBOW METSULFURON-METHYL 60 WG Directions for Use.

AERIAL APPLICATION:

Apply a minimum of 25 L of spray volume per hectare. Apply a maximum of 50 L of spray volume per hectare. Use 50 L/ha for better coverage in pasture, rangeland and non-crop areas.

Apply only by aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label. Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate specified for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment. Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices or a Global Positioning System (GPS).

Use Precautions:
Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.
Product Specific Precautions:
Read and understand the entire label before opening this product. If you have questions, call Rainbow CropSciences Inc. at (630) 828-5088 or obtain technical advice from the distributor or your provincial agricultural representative.
Application of this specific product must meet and/or conform to the following:
Volume: Apply the specified rate in a minimum spray volume of 25 litres per hectare for conventionally piloted aircraft.

SPRAY DRIFT MANAGEMENT:

DO NOT re-enter treated fields until 12 hours after application.

Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

Field sprayer application: DO NOT apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. DO NOT apply with sprays finer than the American Society of Agricultural and Biological Engineers (ASABE) S572 (572.1 to 572.3) Medium classification for application to crops. DO NOT apply with spray droplets smaller than the American Society of Agricultural and Biological Engineers (ASABE) Coarse for application to pasture and rangeland. Boom height must be 60 cm or less above the crop or ground.

Conventionally piloted aircraft application: DO NOT apply when wind speed is less than 1 km/h. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with sprays finer than the American Society of Agricultural and Biological Engineers (ASABE) S572 (572.1 to 572.3) Coarse classification. Reduce drift caused by turbulent wingtip vortices. Nozzle distribution along the spray boom length MUST NOT exceed 65% of the wing- or rotorspan.

The potential for spray drift with ground broadcast applications can be reduced by:
- Applying a coarse spray using large droplet producing nozzle tips.
- The use of Radiarc* or an equivalent drift control system or additive.
- Keep the spray boom as low as possible.
- Apply with minimum wind velocity.
- When using a power sprayer and handgun, direct sprays no higher than the tops of the target plants.

SPRAY BUFFER ZONES:

A spray buffer zone is NOT required for:
- uses with hand-held application equipment permitted on this label,
- low-clearance hooded or shielded sprayers that prevent spray contact with crop, fruit or foliage,

For application to rights-of-way, spray buffer zones for protection of sensitive terrestrial habitats are not required; however, the best available application strategies which minimize off-site drift, including meteorological conditions (e.g. wind direction, low wind speed) and spray equipment (e.g. coarse droplet sizes, minimizing the height above canopy), should be used. Applicators must, however, observe the specified spray buffer zones for protection of sensitive aquatic habitats.

The spray buffer zones specified in the tables below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrub lands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

Ground Application Rate (and use)*	Spray Buffer Zones (in meters) Required for the Protection of :		
	Freshwater Habitat of Depths :		Terrestrial Habitat*
	Less than 1 m	Greater than 1 m	
4.5 g a.i./ha (crops)	1	0	15
12 g a.i./ha (pasture and rangeland)	1	1	10
15 g a.i./ha (pasture and rangeland)	1	1	10
18 g a.i./ha (pasture and rangeland)	1	1	10
60 g a.i/ha (non-crop areas)	2	1	35
90 g a.i/ha (non-crop areas)	3	1	45

*Spray buffer zones for the protection of terrestrial habitats are not required for use on rights-of-way including railroad ballast, rail and hydro rights-of-way, utility easements, roads, and training grounds and firing ranges on military bases.

Method of application	Crop	Aircraft	Droplet size	Spray Buffer Zones (metres) Required for the Protection of:		
				Freshwater Habitat of Depths:		Terrestrial* habitat
				Less than 1 m	Greater than 1 m	
Aerial	Pasture, range land and rough turf	Fixed wing	Coarse	20	10	175
			Coarse to very Coarse	15	10	150
			Very Coarse	15	10	125
			Very Coarse to Extremely coarse	15	10	125
		Rotary wing	Coarse	15	10	150
			Coarse to very Coarse	15	10	125
			Very Coarse	15	10	100
			Very Coarse to Extremely coarse	10	10	95
	Non-crop areas	Fixed wing	Coarse	70	35	800
			Coarse to very Coarse	60	30	800
			Very Coarse	50	25	800
			Very Coarse to Extremely coarse	40	20	750
		Rotary wing	Coarse	55	30	800
			Coarse to very Coarse	45	25	525
			Very Coarse	40	20	475
			Very Coarse to Extremely coarse	35	15	425

*Spray buffer zones for the protection of terrestrial habitats are not required for use on rights-of-way including railroad ballast, rail and hydro rights-of-way, utility easements, roads, and training grounds and firing ranges on military bases.

When tank mixes are permitted, consult the labels of the tank mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASABE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product for conventional application equipment can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Drift Mitigation portion of the Canada.ca website.

MIXING INSTRUCTIONS:

1. Fill the spray tank 1/4 to 1/3 full with clean water. Prepare mixture volume to apply a minimum of 100 litres spray volume per hectare.

2. With the agitator running, add the required amount of RAINBOW METSULFURON-METHYL 60 WG.

3. Continue to agitate until RAINBOW METSULFURON-METHYL 60 WG is completely dispersed.

4. If another herbicide is to be added, do this now.

5. Add the rest of the required water.

6. If required in the mixture, add the surfactant now.

7. On repeat tank loads, draw down the tank contents to less than 10% of the original volume, and repeat from step 1. If more than 10% of the spray volume remains, pre-slurry the RAINBOW METSULFURON-METHYL 60 WG in a bucket of water before adding to the spray tank.

8. If an antifoam agent is required, add it last.

NOTE:
Continuous agitation is required to keep RAINBOW METSULFURON-METHYL 60 WG in suspension in the spray tank for a uniform application.

Use spray preparation of RAINBOW METSULFURON-METHYL 60 WG within 48 hours or product degradation may occur. If spray preparation is left standing without agitation, thoroughly agitate before spraying.

Avoid over-filling of spray tank. Mix only enough product for the job at hand. During spray operation, make scheduled checks of spray equipment to help ensure proper application.

Do not mix RAINBOW METSULFURON-METHYL 60 WG with liquid fertilizers, substances that contain boron or substances that release free chlorine. Mixing RAINBOW METSULFURON-METHYL 60 WG with any of these compounds will result in an insoluble substance in the tank.

APPLICATION INSTRUCTIONS:
Apply the spray mixture with ground equipment only, using a minimum of 100 litres spray volume per hectare. Use a properly calibrated sprayer that will ensure thorough coverage and a uniform spray pattern. Flat fan nozzles are recommended. Use 50-mesh screens or larger.

Continuous agitation is required to keep RAINBOW METSULFURON-METHYL 60 WG in suspension. Avoid overlapping, and shut off spray booms while starting, turning, slowing or stopping to prevent over-application.

NOTE:
Extreme care must be taken to prevent drift to desirable plants or non-target agricultural land.

SPRAYER CLEANUP:
Immediately after spraying, thoroughly remove all traces of RAINBOW METSULFURON-METHYL 60 WG from mixing and spray equipment in order to avoid subsequent injury to sensitive crops. Follow these instructions:

1. Empty the tank and drain the sump completely. Remove any contamination on the outside of the spraying equipment by washing with clean water.

2. Spray the interior tank walls (including the lid) with clean water using a minimum volume of 10% of the tank volume. Add household ammonia (containing a minimum 3% ammonia) at a solution rate of 1 L/100 L water or ALLCLEAN Cleaner (or other similarly approved cleaner) to the tank. Circulate the water through the lines, including all by-pass lines, for at least two minutes. Flush the boom well and empty the sprayer. Completely drain the sump.

3. Repeat step 2. For this rinse, the addition of household ammonia or other cleaner is not required.
4. Remove the strainers, nozzles, tips and screens and clean separately in a bucket containing water and ammonia solution.

If only ammonia is used as a cleaner, the rinsate solution may be applied to the crop(s) listed on this label. Do not exceed the maximum-labeled use rate. If other cleaners are used consult the cleaner label for rinsate disposal instructions. If no instructions are given, dispose of the rinsate on site or at an approved waste disposal facility.

CAUTION
DO NOT use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty chlorine odour which may cause eye, nose, throat and lung irritation. DO NOT clean equipment in an enclosed area.

MINIMUM CROP ROTATION GUIDELINES:
Rotation options are determined by soil pH, the crop to be planted and a minimum interval. Minimum interval is the time from the last application of RAINBOW METSULFURON-METHYL 60 WG to date of planting the rotational crop.

BLACK AND GREY WOODED SOILS INTERVAL PRIOR TO PLANTING (MONTHS AFTER APPLICATION)									
Soil pH	**SW**	**DW**	**BR**	**OT**	**RS**	**FL**	**LN**	**CS**	**YM**
6.9 or Lower	10	10	10	10	10	10	34	48	48
7.0 to 7.9	10	10	10	10	22	34	48	48	48

SW = spring wheat; DW = durum wheat; BR = barley; OT = oats; RS = canola; FL = flax; LN = lentils; CS = canary seed; YM = yellow mustard
On black and grey wooded soils of pH 7.5 or lower, FESCUE may be planted in 10 months; ALFALFA, RED CLOVER, PEAS, and FLAX 22 months following application of RAINBOW METSULFURON-METHYL 60 WG.

Do not use on soils with pH greater than 7.9. All other crops FIELD BIOASSAY*

BROWN AND DARK BROWN SOILS INTERVAL PRIOR TO PLANTING (MONTHS AFTER APPLICATION)									
Soil pH	**SW**	**DW**	**BR**	**OT**	**RS**	**FL**	**LN**	**CS**	**YM**
6.9 or Lower	10	10	10	10	22	22	34	48	48
7.0 to 7.9	10	10	10	22	34	34	48	48	48

SW = spring wheat; DW = durum wheat; BR = barley; OT = oats; RS = canola; FL = flax; LN = lentils; CS = canary seed; YM = yellow mustard

Do not use on soils with pH greater than 7.9.

All other crops “FIELD BIOASSAY” section

IMPORTANT:
Land previously treated with RAINBOW METSULFURON-METHYL 60 WG cannot be rotated to crops other than those listed until a FIELD BIOASSAY confirms that residues of RAINBOW METSULFURON-METHYL 60 WG are not present. A FIELD BIOASSAY involves growing test strips of the crop(s) intended for production the following year in fields previously treated with RAINBOW METSULFURON-METHYL 60 WG. Crop response will indicate whether or not to rotate to the crop(s) used in the test strip. See “Field Bioassay” section. A successful bioassay would be anticipated in the 3rd or 4th season after application (soil pH 7.0 – 7.9) dependent on the rotational crop selected. Failure to follow these instructions could result in injury to subsequent crops.

NOTE:

Wherever RAINBOW METSULFURON-METHYL 60 WG is used on land previously treated with DuPont GLEAN® Herbicide, read the rotational guidelines on both labels and follow the one with the longest interval stated for your situation.

NOTE:
When considering interval prior to re-cropping broadleaf crops after a RAINBOW METSULFURON-METHYL 60 WG application, extend the rotational interval one year if rainfall was less than 130 mm in the Brown & Dark brown Soil Zones or 250 mm in the Black and Grey Wooded Soil Zones in any year within a stated interval prior to planting.

FIELD BIOASSAY:
Select a representative area or areas of the field previously treated with RAINBOW METSULFURON-METHYL 60 WG to plant your bioassay crop(s). Be sure to consider factors such as size of field, soil texture, drainage and turn-around areas when selecting the site(s) that are most representative of the soil conditions in the field. On large fields, more than one site may be needed in order to obtain reliable results.

Plant the test strips perpendicular to the direction in which the field was sprayed. The strips should be long enough to cross the width of several spray swaths. Large test strip areas are more reliable than small ones.

Use standard tillage and seeding equipment to plant the bioassay.

Prepare a seed bed and plant the crops and varieties you want the option of growing the following year. It is important to use the same planting time, conditions, techniques and cultural practices you normally use to plant and grow the bioassay crop(s). Also plant into an adjacent area not treated with RAINBOW METSULFURON-METHYL 60 WG to use as a comparison.

As the crop(s) emerges and grows, examine these key points in RAINBOW METSULFURON-METHYL 60 WG treated and non-treated areas:

- crop stand
- root development
- rate of growth
- plant colour and vigour
- yield

Allow the bioassay crop(s) to grow to maturity while making your observations.

Do not overspray the test strips with herbicides that may damage the bioassay crop(s).

If the bioassay indicates that RAINBOW METSULFURON-METHYL 60 WG residues are still present, continue cropping only to those crops listed on the label, and do not rotate to other crops until bioassay results indicate that susceptible crops are growing normally.

RESISTANCE-MANAGEMENT RECOMMENDATIONS:
For resistance management, RAINBOW METSULFURON-METHYL 60 WG is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to RAINBOW METSULFURON-METHYL 60 WG and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.
To delay herbicide resistance:

- Where possible rotate the use of RAINBOW METSULFURON-METHYL 60 WG or other Group 2 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from different groups when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone

partner.

- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor treated weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Rainbow CropSciences Inc. at (630) 828-5088 for further information.

USE PRECAUTIONS:

- Do not use on soils above pH 7.9.
- Do not use on highly variable soils that have large gravelly or sandy areas, eroded knolls, or calcium deposits.
- Do not use more than 7.5 grams per hectare per year on agricultural crops.
- Do not apply to irrigated land where tail water will be used to irrigate other crop land.

Because crop varieties differ in their tolerance to herbicides, limit first use of RAINBOW METSULFURON-METHYL 60 WG to a small area of each variety prior to adoption as a field practice.

Do not apply RAINBOW METSULFURON-METHYL 60 WG to a crop that is stressed by severe weather conditions, drought, low fertility, water saturated soil, disease or insect damage as crop injury may result. Drought, disease or insect damage following application also may result in crop injury.

Do not apply RAINBOW METSULFURON-METHYL 60 WG during periods of intense rainfall or to soils saturated with water. Do not apply directly to standing or running water. Do not apply in areas where surface water from the treatment site can run off to adjacent cropland, or into streams, irrigation water or wells.

Under certain conditions (such as heavy rainfall, prolonged cool weather, frost conditions, wide fluctuations in day/night temperatures) temporary lightening in crop colour and occasionally a slight reduction in crop height may occur.

Do not apply to a crop undersown with legumes or grasses as injury to the forage may result. Do not remove soil from treated fields for use on lawns, in gardens or for backfill.

Applications should only be made when there is no hazard of spray drift contaminating non-target land areas since very small quantities of the RAINBOW METSULFURON-METHYL 60 WG spray solution may severely injure susceptible crops or other non-target plants.

Aerial application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) coarse classification.

To reduce drift caused by turbulent wingtip vortices, the nozzle distribution along the spray boom length MUST NOT exceed 65% of the wing- or rotorspan.

The additive effect of soil residues from the use of RAINBOW METSULFURON-METHYL 60 WG and imazamethabenz-methyl on the same land has not been determined. Crop rotation guidelines and minimum rotation intervals are not known and injury to rotational crops other than wheat (excluding durum) may occur. Wherever land will be treated with RAINBOW METSULFURON-METHYL 60 WG and has been or will be treated with imazamethabenz-methyl, plant only wheat (excluding durum) after both herbicides have been used, until a bioassay (see "Bioassay" section of the RAINBOW METSULFURON-METHYL 60 WG label) demonstrates that other crops can be successfully grown.

STORAGE

Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed.

DISPOSAL:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

For information on disposal of unused, unwanted product, contact the registrant and the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill and for cleanup of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.