2025-2093
2026-10-06

{Booklet Label}

GROUP	**4**	**FUNGICIDE**
GROUP	**7**	**FUNGICIDE**
GROUP	**22**	**FUNGICIDE**

V-10536 FS

SEED TREATMENT FUNGICIDE
Suspension
COMMERCIAL

A fungicide seed treatment product providing contact and systemic control or suppression of labeled seed and seedling diseases.

An INDIFLIN® brand

ACTIVE INGREDIENTS:
Ethaboxam……..…….. 57.5 g/L
Inpyrfluxam……..…… 38.4 g/L
Metalaxyl……..…… 30.7 g/L

Contains 1,2-benzisothiazolin-3-one at 0.09% as a preservative.

REGISTRATION NO. 36089 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND BOOKLET BEFORE USING
KEEP V-10536 FS AND TREATED SEED OUT OF REACH OF CHILDREN AND ANIMALS

For use with closed transfer commercial seed treaters (facilities and mobile treaters). Closed transfer includes closed mixing, loading, calibrating and closed treatment equipment.

This product contains no colourant. An appropriate colourant must be added when the product is applied to seed, according to regulations pertaining to the *Seeds Act*.

NET CONTENTS: 50 mL to bulk

Valent Canada, Inc.
201-230 Hanlon Creek Blvd.,
Guelph, Ontario, N1C 0A1
(519) 767-9262

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

FIRST AID:

IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.
IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.
IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.
IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL OR POISONING CALL 1-800-682-5368

TOXICOLOGICAL INFORMATION:

No specific antidote is available. Treat the patient symptomatically.

PRECAUTIONS:

KEEP V-10536 FS AND TREATED SEED OUT OF REACH OF CHILDREN AND ANIMALS.

DO NOT eat, drink or smoke/vape while using this product. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum/tobacco, smoking/vaping, using the toilet or handling mobile devices. Vehicle cabs MUST be decontaminated regularly.

Apply only in a way that this product will not contact workers or other persons, either directly or through drift. Only workers wearing the personal protective equipment may be in the area where treating, bagging, sewing and/or stacking is occurring.

Follow the personal protective equipment, engineering controls and restriction requirements for the appropriate task and scenario. These restrictions are required to minimize exposure to the worker.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Seed Type	Tasks	PPE/Engineering Controls
For Commercial Seed Treatment Facilities and Mobile Treaters		

All seed types	Mixing, loading, treating, calibrating, bagging, sewing seed bags, stacking and driving a forklift	Closed transfer only[1]. Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes. In addition, when bagging or sewing bags of treated seeds or when transferring seeds to a storage bin, workers MUST wear a NIOSH-approved N95 filtering face piece respirator (dust mask) that is properly fit tested.
	Cleaning and repairing	Wear chemical-resistant coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes.
For Handling and Planting Commercially Treated Seeds		
All seed types	Handling and planting	Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes. Use a closed cab tractor when planting treated seed. Gloves are not required within a closed cab.

[1] Closed transfer includes closed mixing, loading, calibrating, and closed treatment. No open transfer is permitted.

If the product comes in contact with clothing/PPE, remove all contaminated clothing/PPE, wash the outside of gloves with soap and water before removing, wash contaminated skin with soap and water, and then dress in clean clothing. Launder contaminated clothing/PPE regularly and separately from other laundry. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Heavily contaminated or drenched clothing and other absorbent materials contaminated with this product should be discarded. DO NOT reuse them.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., contact Valent Canada, Inc.

ENVIRONMENTAL PRECAUTIONS

Toxic to aquatic organisms, birds, small wild mammals and non-target terrestrial plants. Any spilled seeds must be incorporated into the soil or otherwise cleaned up from the soil surface.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

SEED QUALITY AND STORAGE OF TREATED SEED:

Use proper storage conditions for treated seed. Due to seed quality and seed storage conditions beyond the control of Valent Canada, Inc., no claims are made to guarantee the germination of carry-over treated seed.

USE RESTRICTIONS:

- DO NOT use treated seed for food, feed, or oil processing.

- Due to seed quality conditions beyond the control of Valent Canada, Inc., no claims are made to guarantee germination of carry-over seed.

LABELLING OF TREATED SEED:
All containers or packages containing treated corn, legume vegetables and soybean seeds for sale or use in Canada must be labeled or tagged as follows:

- KEEP TREATED SEED OUT OF REACH OF CHILDREN AND ANIMALS.
- This seed has been treated with *V-10536* FS containing ethaboxam, metalaxyl, and inpyrfluxam.
- DO NOT graze or feed livestock on seeded area for four weeks after planting.
- DO NOT use for feed, food or oil processing.
- Excess seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practices.
- Store treated seed away from food or feed.
- During handling and planting of treated seed, wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes. Use a closed cab tractor when planting treated seeds. Gloves are not required within a closed cab.
- Dispose of all excess treated seed. Leftover treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. DO NOT contaminate water bodies when disposing of planting equipment washwaters.
- Dispose of seed packaging in accordance with local requirements.
- Toxic to birds and small wild mammals. Any spilled seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface

USE INFORMATION:
V-10536 FS (containing ethaboxam metalaxyl, and inpyrfluxam) is a fungicide seed treatment product offering both contact and systemic activity in providing broad-spectrum seed and seedling protection against labeled seedling diseases. The fungicide components in *V-10536 FS* are effective for protection against *Pythium* spp., *Phytophthora sojae, Aphanomyces euteiches* and *Rhizoctonia solani*.

APPLICATION DIRECTIONS:
Thoroughly shake product before opening container.

V-10536 FS must be used only in accordance with use instructions on this label utilizing mechanical, slurry or mist-type seed treating equipment, provided that the equipment can be calibrated to accurately and uniformly apply the product to seed. This product is for commercial application. DO NOT apply this product in a hopper-box or planter-box at planting time. Always mix product thoroughly before use.

Accurate application of *V-10536 FS* to seed is required for best performance. Thoroughly mix *V-10536 FS* in the desired seed slurry mixture, prior to application to seed.

Tank mixing
This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. DO NOT tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Valent Canada, Inc. at 519-767-9262 for information before applying any tank mix that is not specifically recommended on this label. Pre-test tank mixtures with other seed treatment products to evaluate formulation compatibility and to ensure proper physical compatibility of products.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

The following tables provide use rates of *V-10536 FS* by labeled crop, with disease protection guidelines.

Corn (sweet and field); popcorn	
Diseases Controlled	**Application Rate**
Seed rot, seedling blight and damping off caused by *Rhizoctonia solani* Seed rot/pre-emergence damping-off, post-emergence damping-off and seedling blight caused by *Fusarium* spp. (suppression) Seed rot/pre-emergence damping-off caused by *Pythium* spp., including metalaxyl-resistant *Pythium* spp.	130 mL/100 kg seed* *The following active ingredient components will be applied: ethaboxam: 7.5 g a.i./100 kg seed metalaxyl: 4 g a.i./100 kg seed inpyrfluxam: 5 g a.i./100 kg seed **Field Corn:** Apply 26.6 mL per 80,000 seed (equals 0.042 mg ai/seed) based on 3968 seed/kg count **Sweet Corn:** Apply 19 mL per 100,000 seed (equals 0.03 mg ai/seed) based on 6938 seed/kg count **Popcorn:** Apply 19.2 mL per 100,000 seed (equals 0.03 mg ai/seed) based on 6742 seed/kg count

Crop Group 6	
Legume Vegetables, Succulent or Dried (except soybean) Bean (*Lupinus*) (includes grain lupin, sweet lupin, white lupin, and white sweet lupin); bean (*Phaseolus*) (includes field pea, cowpea, field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean); bean (*Vigna*) (includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean); broad bean (fava); chickpea (garbanzo); guar; jackbean; lablab bean; lentil; pea (*Pisum*) (includes dwarf pea, edible-podded pea, English pea, garden pea, green pea, snowpea, sugar snap pea); pigeon pea; sword bean	
Diseases Controlled	**Application Rate**

Seed rot, seedling blight and damping off caused by *Rhizoctonia solani* Seed rot/pre-emergence damping-off, post-emergence damping-off and seedling blight caused by *Fusarium* spp. (suppression) Seed rot/pre-emergence damping-off caused by *Pythium* spp., including metalaxyl-resistant *Pythium* spp. Early season root rot caused by *Aphanomyces euteiches* (suppression only)	130 mL/100 kg seed* * The following active ingredient components will be applied: ethaboxam: 7.5 g a.i./100 kg seed metalaxyl: 4 g a.i./100 kg seed inpyrfluxam: 5 g a.i./100 kg seed **Bean:** Apply 37 mL per 100,000 seed (equals 0.03 mg ai/seed) based on 3524 seed/kg count **Pea:** Apply 18.6 mL per 100,000 seed (equals 0.017 mg ai/seed) based on 7007 seed/kg count
Soybean	
Diseases Controlled	**Application Rate**
Seed rot, seedling blight and damping off caused by *Rhizoctonia solani* Seed rot/pre-emergence damping-off, post-emergence damping-off and seedling blight caused by *Fusarium* spp. (suppression) Seed rot/pre-emergence damping-off caused by *Pythium* spp., including metalaxyl-resistant *Pythium* spp. Early season root rot caused by *Phytophthora sojae*	130 mL /100 kg seed* * The following active ingredient components will be applied: ethaboxam: 7.5 g a.i./100 kg seed metalaxyl: 4 g a.i./100 kg seed inpyrfluxam: 5 g a.i./100 kg seed **Soybean:** Apply 29.6 mL per 140,000 seed (equals 0.027 mg ai/seed) based on 6173 seed/kg count

RESISTANCE MANAGEMENT RECOMMENDATIONS:
For resistance management, please note that *V-10536 FS* contains Group 4, 7, and 22 fungicides. Any fungal population may contain individuals naturally resistant to V-10536 FS and other Group 4, 7, and 22 fungicides. A gradual or total loss of pest control may occur over time if this fungicide is used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

- Where possible, rotate the use of *V-10536 FS* or other Group 4, 7, and 22 fungicides with different chemical groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that is effective on the target pathogen when such use is permitted.
- Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and

considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications.
- Monitor treated fungal populations for resistance development. Notify Valent Canada, Inc. if reduced sensitivity of the pathogen to *V-10536 FS* is suspected. If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different site of action, if available.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.

Due to the minimal amount of *V-10536 FS* applied to the seed that relocates to the upper parts of the plant and the multiple modes of action, the potential for development of resistance is extremely low. Seed treatments of *V-10536 FS* can therefore be followed by foliar use fungicides containing the same modes of action in accordance with resistance management guidelines as specified on the respective labels.

For further information or to report suspected resistance contact Valent Canada, Inc. at 1-800-682-5368.

STORAGE:
Keep pesticide in its original container. Store in a cool place. DO NOT store in direct sunlight. Protect from freezing temperatures. Store this product away from food or feed.

DISPOSAL:

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for cleanup of spills.

®INDIFLIN is a registered trademark of Sumitomo Chemical Company, Ltd.

{Base Container Label}

GROUP	**4**	**FUNGICIDE**
GROUP	**7**	**FUNGICIDE**
GROUP	**22**	**FUNGICIDE**

V-10536 FS

SEED TREATMENT FUNGICIDE
Suspension
COMMERCIAL

ACTIVE INGREDIENTS:
Ethaboxam……..……..57.5 g/L
Inpyrfluxam……..……38.4 g/L
Metalaxyl……..……30.7 g/L

Contains 1,2-benzisothiazolin-3-one at 0.09% as a preservative.
.

REGISTRATION NO. 36089 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND BOOKLET BEFORE USING
KEEP V-10536 FS AND TREATED SEED OUT OF REACH OF CHILDREN AND ANIMALS

NET CONTENTS: 50 mL to bulk

Valent Canada, Inc.
201-230 Hanlon Creek Blvd.,
Guelph, Ontario, N1C 0A1
519-767-9262

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

FIRST AID:
IF SWALLOWED: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.
IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.
IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.
IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:
No specific antidote is available. Treat the patient symptomatically.

PRECAUTIONS: KEEP V-10536 FS AND TREATED SEED OUT OF REACH OF CHILDREN AND ANIMALS.

Follow the personal protective equipment and restriction requirements for the appropriate worker scenario as described in the booklet label.

ENVIRONMENTAL PRECAUTIONS:
Toxic to aquatic organisms, birds, small wild mammals and non-target terrestrial plants. Any spilled seeds must be incorporated into the soil or otherwise cleaned up from the soil surface.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

STORAGE:
Keep pesticide in its original container. Store in a cool place. Do not store in direct sunlight. Protect from freezing temperatures. Store this product away from food or feed.

DISPOSAL:

DO NOT reuse this container for any purpose. This is a recyclable container and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial or territorial requirements.

For information on disposal of unused, unwanted product, contact the registrant or the provincial or territorial regulatory agency. Contact the registrant and the provincial or territorial regulatory agency in case of a spill, and for cleanup of spills.

See full booklet label for further restrictions and precautions.